

DELLA PROTEIN FUNCTION DURING  
DIFFERENTIAL GROWTH PROCESSES IN  
ARABIDOPSIS

JAVIER GALLEGO BARTOLOME

*Advisors:*

MIGUEL ÁNGEL BLÁZQUEZ

DAVID ALABADÍ DIEGO

VALENCIA, JUNE 2011





El Dr. Miguel Ángel Blázquez Rodríguez, Investigador Científico del CSIC, y el Dr. David Alabadí Diego, Científico Titular del CSIC, ambos pertenecientes al Instituto de Biología Molecular y Celular de Plantas (IBMCP, UPV-CSIC) de Valencia

CERTIFICAN que el Licenciado en Ingeniería Agronómica JAVIER GALLEGO BARTOLOMÉ ha realizado bajo su dirección en el Instituto de Biología Molecular y Celular de Plantas el trabajo que lleva por título “DELLA protein function during differential growth processes in Arabidopsis”, y autorizan su presentación para optar al grado de Doctor.

Y para que así conste, expiden y firman el presente certificado en Valencia, a 1 de Junio de 2011.

Dr. Miguel Ángel  
Blázquez Rodríguez

Dr. David  
Alabadí Diego



En primer lugar quiero agradecer a mis dos tutores, Miguel y David, por todo lo que han hecho por mí estos años de tesis. Os doy las gracias por la formación excelente que he recibido de vosotros. Por haberme guiado, por haber confiado en mí, por haberme hecho sentir uno más y por haberme dado toda la libertad que he sentido en este tiempo. Vosotros sabéis todo lo que hemos trabajado estos años y lo bien que lo hemos pasado en todo momento. Un millón de gracias a los dos.

También a Ramón Serrano por ejercer de tutor de la universidad y mostrarme una sonrisa amable cada vez que necesitaba ayuda.

En siguiente lugar quiero agradecer a mis compañeros de laboratorio que, sin duda, han sido mi segunda familia durante estos años. No hay palabras para expresar todo lo que hemos vivido juntos. De todos me acuerdo y recuerdo grandes momentos. A todos ellos, que bien saben los que son, les agradezco por todo lo que he aprendido de ellos, a nivel científico y humano, por la paciencia que han tenido a veces, y por la generosidad que han mostrado siempre. A todos los deseo lo mejor del mundo. Por supuesto este deseo se extiende a todos los compañeros del Instituto que han compartido un sinfín de experiencias conmigo. En concreto a la gente del 2.08, 2.10 y 2.11 con los cuales he compartido espacios, ciencia y, sobre todo, amistad. En especial quiero nombrar a María Angeles Argomániz, por todo el cariño que me ha dado estos años.

Es imposible no nombrar a algunas personas que han sido clave para mí durante estos años. Jorge (Lozano!), Vero y Marian. Los tres los sabéis, pero nunca está de más: os quiero!

Tampoco me quiero olvidar de los científicos “de fuera”. Mis inicios en el IVIA y todos los grandes amigos que da ahí han salido. Por siempre Paco.

También quiero dar las gracias a los laboratorios que me han hospedado temporalmente y a todas las personas que allí me he encontrado.

Otra pieza clave, ajena a este mundo y que me ha dado alas. María Cabrelles. Gracias.

También quiero nombrar a mis amigos de fuera del laboratorio, que me han recordado que no solo de ciencia vive el hombre. Compartir, escuchar, reír, llorar, el Cedro, festivales, cervezas, música, respeto y mucho cariño. De sobra sabéis quienes sois. Estos años no hubieran sido lo mismo sin vosotros.

A mi madre, a mi padre y a mi hermano, por quererme como soy, por darme todo y por enseñarme tanto y tanto. En especial a mi madre; toda una vida no sería suficiente para agradecerle todo lo que has hecho por mí.

Y por último, quiero dar las gracias a Marian. Por estar ahí, por haberme acompañado en esta última etapa de mi vida que ha sido tan dura, por quererme y dejarme quererla y por darme todo lo mejor. GRACIAS. La etapa del doctorado acaba aquí, ahora empieza una nueva en otras tierras. Espero que sigamos disfrutando tanto como le hemos hecho hasta ahora allá donde sea.

Esta tesis ha sido financiada por una beca JAE del CSIC.



*A mi madre*



## **Summary**





The plant hormones gibberellins (GAs) regulate multiple processes of plant development, such as seed germination, photomorphogenesis, vegetative growth, and flower and fruit development. Most of this regulation occurs at the transcriptional level, through the activity of the DELLAs, which are nuclear-localized proteins subjected to GA-mediated proteolytic degradation. DELLAs do not bind DNA directly, but they have been shown to interact with DNA-binding transcription factors to regulate their targets. Moreover, DELLAs mobilizes distinct set of genes to trigger different GAs responses. In Arabidopsis, DELLAs are encoded by five genes, and genetic studies show that each DELLA displays specific, but also partially overlapping roles with respect to their paralogs. In this Thesis, we have addressed two issues: (1) the contribution of *DELLA* multiplication to the diversification of functions controlled by GAs; and (2) the identification of direct targets regulated by DELLAs in etiolated seedlings with special attention to those involved in differential growth processes.

Using combinations of mutants and transgenic lines expressing two phylogenetically distant *DELLA* genes (*RGA* and *RGL2*), we have found that these two DELLA proteins can perform each other's role as long as they are expressed under the reciprocal promoters, indicating that DELLA subfunctionalization relies mainly on their differential expression patterns. In agreement with this, none of these DELLA proteins displayed significant differences in their ability to interact with several bHLH transcription factors, again suggesting that the function of each DELLA protein probably depends on the set of transcription factors to which they are exposed, and their mutual interactions.

To identify direct DELLA targets, we have performed transcriptomic analyses of dark-grown seedlings expressing an inducible version of *gai-1*, a stable, dominant allele of a *DELLA* gene. This approach rendered a list of over 150 genes differentially expressed between 30 and 240 min after induction of *gai-1*. Further *in silico* analysis of this set of targets has confirmed the functional interaction between DELLAs and bHLH transcription factors, but has also allowed the identification of additional transcription factor families putatively involved in transcriptional regulation by DELLAs.

The presence of several auxin-related genes among the primary targets of DELLA proteins has allowed us to establish a new role for GAs in the modulation of hypocotyl gravitropism through the repression of *IAA19/MASSUGU2* expression by DELLAs. Our results also suggest that this regulatory module fine tunes the auxin-driven gravitropic response, providing flexibility under competing tropic stimuli. Moreover, the repression of *HOOKLESS1* and the auxin efflux carriers *PIN3* and *PIN7* by DELLAs, is proposed as the molecular mechanism to explain the already known physiological regulation of apical hook development by GAs.

Las hormonas vegetales giberelinas (GAs) regulan múltiples procesos del desarrollo de las plantas, como la germinación, la fotomorfogénesis, el crecimiento vegetativo y el desarrollo floral. Esta regulación ocurre principalmente sobre la transcripción a través de las proteínas DELLA, que son proteínas nucleares sometidas a degradación proteolítica inducida por GAs. Las DELLAs no unen DNA directamente pero son capaces de interactuar con factores de transcripción que unen DNA para regular sus dianas. Además, las DELLAs movilizan diferentes grupos de genes según el proceso que estén controlando. En *Arabidopsis*, las DELLAs están codificadas por cinco genes y los estudios genéticos muestran que cada DELLA realiza funciones específicas pero también solapan con otras DELLAs en el control de determinados procesos. En esta Tesis hemos abordado dos cuestiones: (1) la contribución de la multiplicación de las DELLAs a la diversificación de funciones controlada por GAs y (2) la identificación de dianas directas reguladas por las DELLAs en plántulas etioladas, con especial atención a aquellas relacionadas con el crecimiento diferencial.

Usando una combinación de mutantes y líneas transgénicas que expresan dos DELLAs filogenéticamente alejadas (RGA y RGL2), hemos encontrado que estas dos proteínas DELLA pueden suplantar la función de la otra siempre que se expresen bajo el control de los promotores recíprocos, lo que indica que la subfuncionalización de las DELLA reside principalmente en sus diferentes patrones de expresión. En consonancia con esto, ninguna de estas dos proteínas mostró diferencias significativas en su capacidad de interactuar con diferentes factores de transcripción de tipo bHLH, de nuevo sugiriendo que la función de cada proteína DELLA depende del grupo de factores de transcripción al que se vean expuestas, y su interacción con éstos.

Para identificar dianas directas de las DELLAs, hemos llevado a cabo un análisis transcriptómico en plántulas etioladas expresando una versión inducible por choque térmico de *gai-1*, un alelo de GAI resistente a degradación inducida por GAs. Con este abordaje se encontraron 150 genes diferencialmente expresados a tiempos cortos tras la inducción de *gai-1*. El análisis *in silico* de este grupo de genes confirmó la relación funcional entre las DELLA y los bHLH, pero además ha permitido la identificación de nuevas familias de factores de transcripción que podrían estar implicadas en la regulación transcriptional mediada por las DELLAs.

La presencia de varios genes relacionados con auxinas entre las dianas directas de las DELLA nos ha permitido encontrar un nuevo papel de las GAs en la modulación de la respuesta gravitrópica, a través de la represión de *IAA19* por las DELLA. Nuestros resultados también sugieren que este módulo sirve para matizar la respuesta a auxinas durante la respuesta gravitrópica, confiriendo flexibilidad bajo situaciones donde se enfrenten diferentes estímulos trópicos. Además, la represión de *HOOKLESS1*, *PIN3* Y *PIN7* por las DELLA, se propone como el mecanismo molecular que explicaría el papel ya conocido de las GAs en el control del desarrollo del gancho apical.

Les hormones vegetals giberelines (GAs) regulen múltiples processos del desenvolupament de les plantes, com la germinació, la fotomorfogènesis, el creixement vegetatiu i el desenvolupament floral. Aquesta regulació ocorre principalment sobre la transcripció a través de les proteïnes DELLA, que són proteïnes nuclears sotmeses a degradació proteolítica induïda per GAs. Les DELLAs no uneixen DNA directament però són capaces d'interaccionar amb factors de transcripció que uneixen DNA per regular les seves dianes. A més, les DELLAs mobilitzen diferents grups de gens segons el procés que estiguin controlant. En *Arabidopsis*, les DELLAs estan codificades per cinc gens i els estudis genètics mostren que cada DELLA realitza funcions específiques però també solapan amb altres DELLAs en el control de determinats processos. En aquesta Tesi hem abordat dues qüestions: (1) la contribució de la multiplicació de les DELLAs a la diversificació de funcions controlada per GAs i (2) la identificació de dianes directes regulades per les DELLAs en plàntules etiolades, amb especial atenció a aquelles relacionades amb el creixement diferencial.

Usant una combinació de mutants i línies transgèniques que expressen dues DELLAs filogenèticament allunyades (RGA i RGL2), hem trobat que aquestes dues proteïnes DELLA poden suplantar la funció de l'altra sempre que s'expressin sota el control dels promotors recíprocs, la qual cosa indica que la subfuncionalització de les DELLA resideix principalment en els seus diferents patrons d'expressió. D'acord amb això, cap d'aquestes dues proteïnes va mostrar diferències significatives en la seva capacitat d'interaccionar amb diferents factors de transcripció de tipus bHLH, de nou suggerint que la funció de cada proteïna DELLA depèn del grup de factors de transcripció al que es vegin exposades, i la seva interacció amb aquests.

Per identificar dianes directes de les DELLAs, hem dut a terme una anàlisi transcriptòmic en plàntules etiolades expressant una versió inducible per xoc tèrmic de *gai-1*, un al·lel de GAI resistent a degradació induïda per GAs. Amb aquest abordatge es van trobar 150 gens diferencialment expressats a temps curts després de la inducció de *gai-1*. L'anàlisi *in silico* d'aquest grup de gens va confirmar la relació funcional entre les DELLA i els bHLH, però a més ha permès la identificació de noves famílies de factors de transcripció que podrien estar implicades en la regulació transcriptional intervinguda per les DELLAs.

La presència de diversos gens relacionats amb auxines entre les dianes directes de les DELLA ens ha permès trobar un nou paper de les GAs en la modulació de la resposta gravitròpica, a través de la repressió de *IAA19* per les DELLA. Els nostres resultats també suggereixen que aquest mòdul serveix per matisar la resposta a auxines durant la resposta gravitròpica, conferint flexibilitat sota situacions on s'enfrontin diferents estímuls tròpics. A més, la repressió de *HOOKLESS1*, *PIN3* i *PIN7* per les DELLA, es proposa com el mecanisme molecular que explicaria el paper ja conegut de les GAs en el control del desenvolupament del ganxo apical.



## **Table of contents**



<b>INTRODUCTION.....</b>	<b>1</b>
<b>I.1 Plant hormones.....</b>	<b>3</b>
<b>I.2 Gibberellins.....</b>	<b>3</b>
<b>I.3 Gibberellin metabolism.....</b>	<b>5</b>
<b>I.4 Gibberellin signaling.....</b>	<b>7</b>
<b>I.5 DELLA proteins.....</b>	<b>8</b>
<b>I.6 Gibberellin-induced DELLA degradation.....</b>	<b>10</b>
<b>I.7 Transcriptional regulation by DELLA proteins.....</b>	<b>12</b>
<b>I.8 Bibliography.....</b>	<b>15</b>
 <b>OBJECTIVES.....</b>	 <b>23</b>
 <b>CHAPTER 1: Transcriptional diversification and functional conservation between DELLA proteins in Arabidopsis.....</b>	 <b>25</b>
<b>1.1. Abstract .....</b>	<b>27</b>
<b>1.2. Introduction.....</b>	<b>28</b>
<b>1.3. Materials and Methods.....</b>	<b>30</b>
<b>1.4. Results and discussion.....</b>	<b>34</b>
1.4.1. Phylogenetic relationships of DELLA proteins in Angiosperms.....	34
1.4.2. Expression of chimeric versions of DELLA genes.....	34
1.4.3. Functional substitution of RGA by RGL2.....	37
1.4.4. Functional substitution of RGL2 by RGA.....	38
1.4.5. Conservation of DELLA protein interactions.....	40
<b>1.5. Bibliography.....</b>	<b>44</b>
 <b>CHAPTER 2: Transcriptional regulation by DELLA proteins during etiolated development in <i>Arabidopsis thaliana</i> .....</b>	 <b>49</b>
<b>2.1. Introduction.....</b>	<b>51</b>
<b>2.2. Results and discussion.....</b>	<b>52</b>
2.2.1. Identification of genes rapidly regulated by GAI in etiolated seedlings.....	52

2.2.2. Comparison of DELLA-regulated genes in different developmental situations.....	54
2.2.3. GAI regulates target genes in part through PIFs and HY5 transcription factors.....	54
2.2.4. Promoter analysis of GAI regulated targets suggests new transcription factors mediating DELLAactivity.....	57
2.2.5. Gene ontology analysis of GAI-regulated genes.....	59
2.2.6. Direct regulation of the GA pathway by DELLA proteins.....	60
2.2.7. DELLA proteins mediate direct cross-regulation with auxin and ethylene pathways	61
2.2.8. DELLAs impinge on transcriptional networks.....	63
<b>2.3. Conclusions.....</b>	<b>64</b>
<b>2.4. Materials and Methods.....</b>	<b>65</b>
<b>2.5. Bibliography.....</b>	<b>67</b>

## **CHAPTER 3: A hormonal regulatory module that provides flexibility to tropic responses.....**

**71**

<b>3.1. Abstract.....</b>	<b>73</b>
<b>3.2. Introduction.....</b>	<b>74</b>
<b>3.3. Results.....</b>	<b>75</b>
3.3.1. Gibberellin deficiency enhances gravitropic reorientation.....	75
3.3.2. Expression of <i>IAA19/MSG2</i> is repressed by DELLA proteins.....	75
3.3.3. Physiological relevance of the regulation of gravitropism by gibberellins.....	80
<b>3.4. Discussion.....</b>	<b>81</b>
<b>3.5. Materials and Methods.....</b>	<b>84</b>
<b>3.6. Bibliography.....</b>	<b>86</b>

## **CHAPTER 4: Hierarchy of hormone action controlling apical hook development in *Arabidopsis*.....**

**91**

<b>4.1. Abstract.....</b>	<b>93</b>
<b>4.2. Introduction.....</b>	<b>94</b>
<b>4.3. Results.....</b>	<b>95</b>
4.3.1. Dynamics of GA-regulated apical hook development.....	95



4.3.2. GA control on hook development is dependent and independent upon ethylene activity.....	96
4.3.3. The expression of <i>ACS5/ETO2</i> , <i>ACS8</i> , and <i>HLS1</i> genes is regulated by the GA pathway.....	98
4.3.4. GA-regulation of <i>ACS5/ETO2</i> and <i>ACS8</i> gene expression depends on the phase of hook development .....	99
4.3.5. GAs support ethylene production in etiolated seedling.....	99
4.3.6. GAs regulate partly hook development by modulating PIF activity.....	101
4.3.7. HLS1 activity mediates GA effect on hook development.....	102
4.3.8. GAs are needed to sustain differential auxin response during apical hook development.....	103
4.3.9. GAs participate in maintaining <i>PIN3</i> and <i>PIN7</i> expression in the apical hook.....	105
4.3.10. GA activity in the endodermis is required for apical hook development.....	107
<b>4.4. Discussion.....</b>	<b>110</b>
4.4.1. GAs regulate hook formation independently of ethylene activity.....	110
4.4.2. GAs prevent hook opening in cooperation with the ethylene pathway.....	111
4.4.3. GAs regulate hook development by transcriptional regulation of auxin and ethylene pathways.....	111
<b>4.5. Experimental procedures.....</b>	<b>114</b>
<b>4.6. Bibliography.....</b>	<b>117</b>
<b>4.7. Supporting figures.....</b>	<b>122</b>
 <b>GENERAL DISCUSSION.....</b>	 <b>125</b>
 <b>D.1. Contribution of gibberellins to plant plasticity relies partially on DELLA protein subfuncionalization.....</b>	 <b>127</b>
<b>D.2. Regulation of transcriptional networks as a mechanism to improve plastic development.....</b>	<b>128</b>
<b>D.3. DELLA co-expressed as a source of new interactors.....</b>	<b>131</b>
<b>D.4. Gibberellin modulation of differential growth processes.....</b>	<b>133</b>
<b>D.5. Future perspectives.....</b>	<b>135</b>
<b>D.6. Bibliography .....</b>	<b>137</b>

**CONCLUSIONS..... 139**

# Introduction



## **I.1. Plant hormones**

Pioneering studies during the 19th century demonstrated that plant growth processes were regulated by “substances” with a capacity to move from one part of the plant to another. Today, more than one hundred years later, most of these substances have been identified as small molecules derived from secondary metabolic pathways. In general, these compounds are present at very low concentrations and act either locally, at or near the site of synthesis, or in distant tissues. The first few plant regulators identified –the so-called “classical plant hormones”– are abscisic acid (ABA), auxins, gibberellins (GAs), brassinosteroids (BRs), cytokinin (CK), ethylene, jasmonic acid (JA), and salicylic acid (SA). Recently a new hormone called strigolactone has been discovered and it is possible that there still exist a few more unidentified growth regulators of this class. Collectively, these compounds regulate different aspects of plant life, from pattern formation to the response against biotic and abiotic stress. Hormones have been classically divided into two main categories according to their roles: *growth-related* hormones (auxins, gibberellins, brassinosteroids, cytokinin) or *stress-related* hormones (ethylene, jasmonic acid, salicylic acid), but more recent studies have established the notion that many of these hormones in fact have an impact in both development and stress responses (Jaillais and Chory, 2010; Bari and Jones, 2009). Although the physiological function of these compounds has been studied for decades, the last 15 years have seen a dramatic increase in our understanding of the molecular mechanisms underlying hormone homeostasis, transport and response. As explained below, GAs constitute an excellent study model to investigate how plants integrate environmental and endogenous information to modulate plant growth, and their signaling activity has become the main focus of this thesis.

## **I.2. Gibberellins**

GAs form a big family of diterpenoid compounds found in plants, fungi and bacteria of which only a few regulate plant growth. The first GA was originally isolated in 1938 as a metabolite from the rice fungal pathogen *Gibberella fujikuroi* (Sawada) Wollen. Infection of plants by the fungus resulted in exaggerated stem elongation, ultimately causing the plant to fall over (Yamaguchi, 2008). Further research showed

that GAs synthesized by the fungus were the compounds responsible for the exaggerated growth and lodging. In the middle 50s, evidence was found that plants could also synthesize GAs, and their role as endogenous growth regulators began to be established (Radley, 1956; Phinney BO, 1957).

To understand the impact that GAs have on plant biology and agriculture, it is worth mentioning that the “Green Revolution” in the 1960s and 1970s, was associated with the use of new dwarf varieties of rice and other cereals, which now are well-characterized mutants in GA metabolism and GA signaling (Peng et al., 1999). Interestingly, these plants were not only smaller in size (with the associated benefits for agriculture, such as increased resources for grain production, higher yield per cultured surface, etc), but also displayed higher tolerance to severe weather conditions as wind or drought (Peng et al., 1999).

Molecular-genetic analysis of these varieties, and of equivalent mutants in model plants such as *Arabidopsis*, has shown that GAs not only regulate plant size, but also other developmental processes, such as germination and flowering, as well as the resistance to different stress factors (Table I.1).


**Table 1: Biological processes regulated by GAs**

Biological process	Phenotype of GA-deficiency	Reference
Germination	Impaired germination	(Lee et al., 2002), (Ogawa et al., 2003)
Photomorphogenesis	Derepressed photomorphogenesis in darkness	(Alabadí et al., 2004), (Achard et al., 2007)
Vegetative growth (roots, aerial part)	Impaired growth	(Veen, 1980), (Peng et al., 1997), (King et al., 2001), (Fu and Harberd, 2003)
Floral induction	Late flowering	(Wilson et al., 1992), (Blázquez et al., 1998)
Flower development	Retarded growth of floral organs	(Yu et al., 2004)
Pollen development	Male sterility	(Wilson et al., 1992), (Goto N, 1999)
Fruit induction	Impaired fruit development	(Garcia-Martinez et al., 1997), (Singh et al., 2002)
Abiotic stress	Increased tolerance	(Achard et al., 2008), (Achard et al., 2006)
Biotic stress	Increased resistance	(Navarro et al., 2008)

### I.3. Gibberellin metabolism

Gibberellins are synthesized from geranylgeranyl diphosphate (GGDP) through a long and complex metabolic pathway which involves several different enzymes and cell compartments (Figure I.1). The first step occurs within the plastid, where the geranyl-geranyl diphosphate (GGDP) is converted to *ent*-kaurene through the activity of *ent*-copalyl diphosphate synthase (CPS) and *ent*-kaurene synthase (KS) ((Sun and Kamiya, 1994; Aach H, 1997; Sun, 1997; Helliwell et al., 2001). *Ent*-kaurene is then converted to GA<sub>12</sub> by *ent*-Kaurene oxidase (KO), located in the membrane of the plastid, and *ent*-kaurenoic acid oxidase (KAO), located in the endoplasmic reticulum (Helliwell et al., 2001; Nelson et al., 2004; Appleford et al., 2006). Successive steps from GA<sub>12</sub> to GA<sub>4</sub> –the main bioactive GA in *Arabidopsis thaliana*– occur in the cytoplasm through the activity of GA 20-oxidases (GA20ox) and GA 3-oxidases (GA3ox) (Spray et al., 1996; Itoh et al., 2001; Appleford et al., 2006). GA<sub>12</sub> is also a substrate for GA 13-oxidases (GA13ox) that will produce GA<sub>54</sub> which is a precursor of GA<sub>1</sub> – the main bioactive gibberellin in rice (Yamaguchi, 2008).

GAs are enzymatically inactivated by different means. The best characterized process is the 2-oxidation of the bioactive GAs, catalyzed by GA 2-oxidases (GA2ox). In fact, these enzymes can also oxidize the C2 of the precursors of bioactive GAs, so they also regulate substrate availability for GA3ox during GA synthesis (Thomas et al., 1999; Schomburg et al., 2003; Lee and Zeevaart, 2005). Another deactivation process found in rice is the 16 $\alpha$ ,17-epoxydation of GA<sub>4</sub> and its precursors (Zhu et al., 2006). The rice mutant *eui*, which lacks the corresponding gene –*eui*–, accumulates huge amounts of bioactive GAs (Zhu et al., 2006). Another deactivation process more recently found in *Arabidopsis* involves the methylation of C6 carboxyl groups of bioactive GAs and their precursors by GA methyltransferases (GAMT). Accordingly, ectopic expression of *GAMT* genes in different plant species causes a GA-deficient dwarf phenotype (Varbanova et al., 2007). In addition, GAs can be converted into conjugates in plants (Schneider et al., 1992; Schliemann, 1994). Theoretically, conjugation of GAs to glucose might render inactive molecules, but there are no experimental data available to date regarding the effect of this process on the concentration of bioactive GAs. The discovery of GA-glycosyl transferases and reverse genetic studies will help to understand their role on GAs metabolism.



**Fig I.1: Representation of predicted subcellular localization of GA metabolism enzymes and sequence or reactions.**

**GGDP**: geranylgeranyldiphosphate **CPS**: ent-copalyl diphosphate synthase; **KS**: ent-kaurene synthase; **ER**: endoplasmic reticulum, **KO**: ent-kaurene oxidase; **KAO**: ent-kaurenoic acid oxidase; **GAMT**: gibberellin methyltransferase; **GA<sub>2ox</sub>**, **GA<sub>3ox</sub>**, **GA<sub>16,17ox</sub>**, **GA<sub>20ox</sub>**: GA oxidases. Adapted from Yamaguchi, S. 2008

The levels of bioactive GAs level are maintained in plants through feedback and feedforward mechanisms (Hedden and Phillips, 2000; Olszewski et al., 2002). Transcript analysis shows that GA signaling targets *GA<sub>20ox</sub>*, *GA<sub>3ox</sub>* and *GA<sub>2ox</sub>* to establish homeostasis. For example expression levels of *GA<sub>20ox</sub>* and *GA<sub>3ox</sub>* is enhanced under GA deficiency conditions whereas expression level decreases after exogenous GA treatment (Chiang et al., 1995; Phillips et al., 1995; Yamaguchi et al., 1998; Xu et al., 1999; O'Neill and Ross, 2002). In contrast, *GA<sub>2ox</sub>* are upregulated after GA treatment (Thomas et al., 1999).

Apart from this mechanism where the GA pathway itself regulates bioactive GA levels, there are additional internal or external cues that can influence GA homeostasis. For instance, several hormones have been shown to affect the expression of GA metabolism genes such as *GA<sub>20ox</sub>*, *GA<sub>3ox</sub>* and *GA<sub>2ox</sub>*. Among them, auxins have been shown to induce the expression of GA metabolism genes in *Arabidopsis* (Frigerio et al., 2006) and other plants resulting in actual changes in GA concentration (Ross et al., 2000; Wolbang and Ross, 2001; Wolbang et al., 2004). Similarly, brassinosteroids have been proposed to induce the expression of *AtGA<sub>20ox1</sub>* (Bouquin et al., 2001). However it is not known whether this induction leads to increased levels of bioactive GA *AtGA<sub>20ox1</sub>* (Jager et al., 2005). On the other hand, a negative effect has been observed of ABA on these GA biosynthetic genes (Schomburg et al., 2003). In the case of ethylene both a positive and a negative effect on GA content have been observed




depending on the developmental context (Hoffmann-Benning and Kende, 1992; Achard et al., 2007).

Likewise, environmental cues have been revealed as important modulators of the levels of bioactive GAs. Light is among the leading cues affecting developmental traits, and several reports have linked light signaling to the regulation of GA biosynthesis in seeds, through an increase in *GA20ox* and *GA3ox* expression, and repression of *GA2ox* (Yamaguchi et al., 1998; Oh et al., 2006; Seo et al., 2006; Yamauchi et al., 2007). Interestingly, the effect of light on GA biosynthesis is the opposite after germination, *i.e.* during seedling development. Illumination of etiolated seedlings causes a very rapid decrease in the expression of GA biosynthesis genes, accompanied by a comparable increase in the expression of *GA2ox* genes (Achard et al., 2007; Alabadí et al., 2008). Temperature is another important external cue that affects plant development in part through the modulation of bioactive GA levels. For instance, it has been shown that in dark-imbibed after-ripened *Arabidopsis* seeds, cold temperatures promote GA synthesis through up-regulation of *GA20ox2* and *GA3ox1* and down-regulation of *GA2ox2* (Yamauchi et al., 2004). Similarly to what happens with light, the effect of temperature in seedlings is opposite to the one in seeds: a shift to higher temperatures induces, in the hypocotyl, a fast up-regulation of *GA20ox1* and *GA3ox1* and down-regulation of *GA2ox1* to promote seedling growth (Stavang et al., 2009). Finally, stress is known to decelerate plant growth, and this is, in part, due to a decrease in GA content. It has been shown that *Arabidopsis* plants grown on higher salt concentrations have lower bioactive GA content (Achard et al., 2006), and upregulation of *GA2ox7* expression has been proposed to be cause (Yamaguchi, 2008). All these observations point out that GA homeostasis is tightly regulated by different internal and external cues that will help to create a final output where optimal GA content is synthesized.

#### **I.4. Gibberellin signaling**

Most of the current knowledge of the molecular mechanism of GA signaling comes from the availability of mutants affected in their response to GAs. According to their phenotypes, three classes of mutants have been isolated in different plant species (Figure I.2): (1) Recessive GA-sensitive dwarves mutants such as *gal* in *Arabidopsis* (Sun and Kamiya, 1994) (2) Dominant GA-insensitive dwarves such as *gai* in *Arabidopsis*, *D8* in maize, and *Rht-B1b/Rht-D1b* in wheat (Koornneef, 1985; Harberd

and Freeling, 1989; Peng and Harberd, 1993, 1997; Peng et al., 1999). Their dominant or semi-dominant behavior was taken as representative of gain-of-function alleles of GA signaling elements with a negative role. Also recessive GA-insensitive dwarves such as *gid1* and *gid2* in rice or *Atgid1* and *sly1* in Arabidopsis (McGinnis et al., 2003; Sasaki et al., 2003; Ueguchi-Tanaka et al., 2005; Griffiths et al., 2006), initially thought to be loss-of-function alleles of GA signaling elements necessary for GA action. And (3) recessive slender mutants such as *slr1-1* of rice or the quintuple *della* mutant of Arabidopsis (Ikeda et al., 2001; Feng et al., 2008) that mimic the exaggerated growth caused by continuous GA application (Potts, 1985).




**Figure I.2 Schematic representation of the three classes of GA mutants**  
Adapted from Harberd, N et al, 2009

Studies of the above categories of mutants enabled a formal genetic definition of the mechanism by which GAs promote growth, long before the molecular basis of this mechanism was apparent. In fact the early hypothesis that GAs would act as “inhibitors of a repressor”, formulated upon the study of slender pea mutants (Brian, 1957), has been later substantiated by molecular genetic approaches in Arabidopsis and rice (Harberd et al., 2009), and has resulted in a model known as the GA-GID1-DELLA mechanism of GA response regulation.

## I.5. DELLA proteins

GA signaling is fairly simple judging from the limited number of components that integrate the pathway. The key components are the DELLA proteins, which act as the repressors whose activity needs to be counteracted by GAs. The first DELLA gene isolated was *GAI* from Arabidopsis (Koornneef, 1985), and it was soon realized that there were five *DELLA* paralogs in Arabidopsis (*GAI*, *RGA*, *RGL1*, *RGL2* and *RGL3*) and closer relatives, while other plant species would have only one ortholog like *SLR1*

in rice (Ikeda et al., 2001). All of them share three common features: (1) a DELLA domain within the N-terminus; (2) a GRAS region within the C-terminus; and (3) they are nuclear localized (Peng and Harberd, 1997; Ikeda et al., 2001; Silverstone et al., 2001; Lee et al., 2002; Wen and Chang, 2002).




**Figure I.3: Schematic representation of a DELLA protein**

Studies with RGA:GFP fusions in *Arabidopsis* showed that DELLA proteins accumulate in the nuclei of cells with low GA concentration, but they suffer rapid destabilization in the presence of GAs (Silverstone et al., 2001). Moreover, GA-induced degradation is absolutely dependent on the DELLA motif, so that *DELLA* mutant alleles lacking this motif (such as *gai-1* and *rga-Δ17*) are stable even in the presence of GAs (Peng et al., 1997; Dill et al., 2001).

Contrary to rice, where a knockout mutant in the single *DELLA* gene displays a phenotype that resembles constitutive GA activity (Ikeda et al., 2001), redundancy of *DELLA* genes in *Arabidopsis* requires the combination of multiple knockouts to obtain a slender phenotype. Moreover, detailed analysis of single and multiple *DELLA* mutants in this species has revealed both distinct and overlapping functions for individual DELLAs in the regulation of plant development.

For instance, the **growth** defect caused by GA deficiency (such as in the *gal-3* mutant) can be partially overcome by a knockout mutation in *RGA* but not *GAI* alone, although concurrent elimination of the two *DELLA* genes causes almost complete recovery of the wild-type size (Dill and Sun, 2001; King et al., 2001). This indicates that both RGA and GAI participate in the control of plant size, with RGA having a more prominent role. Interestingly, a different situation is found when analyzing **flower development**. The *gal-3* mutant presents defective flowers with very short immature stamens and almost absent petals, which cannot be restored by simultaneous loss of RGA and GAI function. Full recovery of flower development can only be attained when *RGA*, *RGL1* and *RGL2* are inactivated, irrespective of the presence of GAI (Cheng et al., 2004). Regarding **germination**, RGL2 has been proposed as the main DELLA

protein that needs to be inactivated during GA-induced breaking of dormancy (Lee et al., 2002). This idea was based on the observation that a single knockout of *RGL2*, but not of any other *DELLA* gene, was able to rescue the germination defect of the *gal-3* mutant. However, the other *DELLA* genes have also been found to regulate germination under different contexts. For instance, RGA and GAI also participate in far-red light mediated repression of germination through the stimulation of ABA biosynthesis (Piskurewicz et al., 2008).



**Figure I.4 Arabidopsis DELLA proteins subfunctionalization.**

Each DELLA is represented with a different color and the contribution of each one to a determinate process is related to the size of typography; the bigger the more important.


These results raise the question of what evolutionary mechanisms have prompted the subfunctionalization of *DELLA* genes in those plant species (such as the Brassicaceae) where recent duplications have occurred.

## **I.6. Gibberellin-induced DELLA degradation**

According to the “inhibitor of an inhibitor” model, DELLA proteins restrain plant growth and other GA responses, while GAs exert their activity by releasing DELLA repression. As previously stated, this is achieved through the degradation of

DELLA proteins, and the GA receptor is an essential component of the degradation machinery.

The soluble GA receptor is encoded by a single gene in rice (*GA-INSENSITIVE DWARF1*, *GID1*) (Ueguchi-Tanaka et al., 2005), and three paralogs in Arabidopsis (*GID1a-c*) with almost overlapping functions (Griffiths et al., 2006; Nakajima et al., 2006; Iuchi et al., 2007). The GID1 protein possesses a central pocket that accommodates bioactive GAs. Upon binding, GA causes an allosteric change in GID1 that results in the N-terminus forming a lid to the pocket (Murase et al., 2008; Shimada et al., 2008). Once in place, the outer surface of the lid interacts with the DELLA protein, specifically with the N-terminal region defined by the DELLA and VHYNP domains (Murase et al., 2008) (Fig I.3). The formation of the GA-GID1-DELLA complex is thought to induce a conformational change in the GRAS domain of the DELLA protein, which then interacts with GID1 and stabilizes the complex (Fig I.5). The stabilized complex is then recognized by a specific SCF E3 ubiquitin-ligase complex involving the F-box proteins *AtSLY1* and *AtSNE* in Arabidopsis or *OsGID2* in rice, respectively (McGinnis et al., 2003; Sasaki et al., 2003; Griffiths et al., 2006; Willige et al., 2007; Hirano et al., 2010; Ariizumi et al., 2011). In turn, SCF<sup>SLY1/GID2</sup> promotes the polyubiquitinylation and subsequent destruction of DELLAs by the 26S proteasome (Fig I. 6a) (McGinnis et al., 2003; Sasaki et al., 2003; Dill et al., 2004; Fu et al., 2004). Hence, *OsGID1* single knockout mutants or *AtGID1* triple knockout mutants display a severe GA deficiency phenotype which cannot be reverted by GA application, suggesting that these are very likely the only GA receptors in plants, at least relevant for growth regulation. Interestingly, one of the Arabidopsis GID proteins -*AtGID1b*- is able to interact with DELLA proteins in the absence of GAs because its aminoacid sequence forms a partially closed lid independent of the presence of GAs and hence it is able to interact and promote the DELLA proteolytic degradation (Yamamoto et al., 2010). Soybean (*Glycine max*) and *Brassica napus* also have GID1s similar to *AtGID1b*, indicating that these unique GID1s occur in various dicots and may have important functions in these plants (Yamamoto et al., 2010).



**Figure I.5 GA-GID-DELLA complex**

a) Schematic representation of the GA-GID-DELLA complex formation.

b) 3D model of the GA-GID-DELLA complex from two points of view.


Adapted from Murase, K. et al. 2008

There are many indications that GAs regulate growth and most of the other processes through changes in gene expression. Indeed, one of the earliest molecular events associated with GA signaling was the regulation of alpha-amylase gene expression in barley aleurone cells (Varner JE, 1965 ). More recently, transcriptomic analysis provided a fairly comprehensive view of the different changes and patterns of gene expression in response to GAs in different contexts, such as germination, vegetative growth and flower development (Ogawa et al., 2003; Cao et al., 2006; Nemhauser et al., 2006; Zentella et al., 2007; Hou et al., 2008). Two interesting conclusions can be drawn from these analyses. Firstly, GAs regulate distinct sets of genes in the different tissues examined and in the different developmental stages (Cao et al., 2006). And secondly, all the changes in gene expression in response to GAs are mediated by DELLA proteins (Cao et al., 2006). Of course, given that all these approaches provide a snapshot of gene expression in a specific moment of a plant's life, they do not distinguish between direct targets for GAs/DELLAs and secondary effects.

It seemed reasonable that DELLA proteins act as transcription factors, based on two additional observations: their C-terminus is similar to that of other GRAS proteins like SHORTROOT (SHR) and SCARECROW (SCR), which have been shown to regulate transcription (Levesque et al., 2006; Cui et al., 2007); and DELLA proteins accumulate in the nucleus (Silverstone et al., 1998; Ogawa et al., 2000). Although there is no evidence for direct interaction between DELLA proteins and DNA *in vitro*, chromatin immunoprecipitation (ChIP) experiments with a stable version of RGA have shown an association between DELLA proteins and the promoters of a few genes whose expression is regulated by GAs (Zentella et al., 2007). Therefore, the most likely possibility is that DELLA proteins interact with other DNA-binding transcription factors to modify their activity, and the identification of such partners of DELLA proteins would be pivotal to understand and manipulate GA signaling.

An important clue about the identity of such transcription factors was found in the study of the repression of photomorphogenesis that GAs exert in etiolated seedlings (Alabadí et al., 2004). The systematic analysis of multiple light signaling mutants under GA-deficient conditions revealed that only two types of transcription factors were required for the regulation of photomorphogenesis by DELLA proteins (Alabadí et al., 2008): the bZIP protein HY5 (ELONGATED HYPOCOTYL5), and two members of the PIF (PHYTOCHROME-INTERACTING FACTOR) family of bHLH transcription factors, PIF3 and PIF4. HY5 and PIFs have opposite roles in light signal transduction: while HY5 promotes photomorphogenesis (Ang and Deng, 1994; Lee et al., 2007), PIF proteins are required for etiolated growth and the repression of light-induced gene expression (Ni et al., 1998; Leivar et al., 2008; Leivar et al., 2009). The observation that *hy5* mutants were more resistant to the accumulation of DELLA proteins, and *pif* mutants were hypersensitive in the same conditions was a strong indication that DELLAs might act through some of these transcription factors.

Indeed, PIF proteins have been found to interact physically with the LHR1 domain of DELLA proteins (Fig I.3) and through the bHLH domain, and this interaction prevents binding of the transcription factors to their target promoters and their subsequent activation (de Lucas et al., 2008; Feng et al., 2008). This important finding not only identifies PIFs as a transcription factors through which GAs regulate gene expression (Fig I.6), but it also provides a molecular framework for the interaction between GA and other signaling pathways, given that the availability of PIF proteins is strictly regulated by light and the circadian clock.



**Figure I.6 Transcriptional regulation by DELLA proteins.**

Schematic representation of how GAs regulate transcription of target genes.

The physical interaction of DELLA proteins with other members of the bHLH family has been recently proved. DELLA can interact and sequester SPATULA (SPT) to control cotyledon expansion (Josse et al., 2011). Similarly, DELLAs can interact with ALCATRAZ (ALC) and avoid its function to control fruit patterning (Arnaud et al., 2010). Thus, inactivation of bHLH proteins through interaction with DELLA proteins seems to be a general mechanism of action of GA signaling pathway.

It is thought that DELLA proteins act as dimmers in planta through the interaction with their LHR1 domain (Fig I.3) (Itoh et al., 2002). As mentioned before, DELLA proteins belong to the GRAS family where many transcription factors are found. Interestingly, it was recently shown that DELLA can interact with one member of this family, SCARECROW-LIKE3 –SCL3- to control several aspects of plant development such as germination, hypocotyls length and root growth (Heo et al., 2011; Zhang et al., 2011).

But DELLA proteins seem to be able to interact with other protein different than transcription factors. This is the case for the interaction of DELLA with the JASMONATE ZIM-domain (JAZ) proteins, which are negative regulators of the jasmonate signaling pathway (Hou et al., 2010). In the absence of jasmonate, JAZ proteins accumulate in the nucleus sequestering the jasmonate-response transcription factor MYC2 (Chini et al., 2007). Interaction of DELLA proteins through the N-terminal region (DELLA domains and LHR1 (Fig I.3)) with JAZ proteins release MYC2 protein that can trigger the jasmonate response (Hou et al., 2010).

Thus, different strategies seem to have evolved for the DELLA control of gene expression. Remarkably, the LHR1 domain seems to be most important regarding



interaction with transcription regulators. Hence, it will be important to identify additional interactors of DELLA proteins –possibly, but not only, transcription factors–, and also establish the list of direct target genes for DELLA proteins in specific tissues/processes.

## I.8. Bibliography

- Aach H BH, Robinson DG, Graebe JE** (1997) ent -Kaurene synthase is located in proplastids of meristematic shoot tissues *Planta* **202**: 211-219
- Achard P, Baghour M, Chapple A, Hedden P, Van Der Straeten D, Genschik P, Moritz T, Harberd NP** (2007) The plant stress hormone ethylene controls floral transition via DELLA-dependent regulation of floral meristem-identity genes. *Proc Natl Acad Sci U S A* **104**: 6484-6489
- Achard P, Cheng H, De Grauwe L, Decat J, Schoutteten H, Moritz T, Van Der Straeten D, Peng J, Harberd NP** (2006) Integration of plant responses to environmentally activated phytohormonal signals. *Science* **311**: 91-94
- Achard P, Liao L, Jiang C, Desnos T, Bartlett J, Fu X, Harberd NP** (2007) DELLAs Contribute to Plant Photomorphogenesis. *Plant Physiol* **143**: 1163-1172
- Achard P, Renou JP, Berthome R, Harberd NP, Genschik P** (2008) Plant DELLAs restrain growth and promote survival of adversity by reducing the levels of reactive oxygen species. *Curr Biol* **18**: 656-660
- Alabadí D, Gallego-Bartolomé J, García-Cárcel L, Orlando L, Rubio V, Martínez C, Frigerio M, Iglesias-Pedraz JM, Espinosa A, Deng XW, Blázquez MA** (2008) Gibberellins modulate light signaling pathways to prevent Arabidopsis seedling de-etiolation in darkness. *Plant J* **53**: 324-335
- Alabadí D, Gil J, Blázquez MA, García-Martínez JL** (2004) Gibberellins repress photomorphogenesis in darkness. *Plant Physiol* **134**: 1050-1057
- Ang LH, Deng XW** (1994) Regulatory hierarchy of photomorphogenic loci: allele-specific and light-dependent interaction between the HY5 and COP1 loci. *Plant Cell* **6**: 613-628
- Appleford NE, Evans DJ, Lenton JR, Gaskin P, Croker SJ, Devos KM, Phillips AL, Hedden P** (2006) Function and transcript analysis of gibberellin-biosynthetic enzymes in wheat. *Planta* **223**: 568-582
- Ariizumi T, Lawrence PK, Steber CM** (2011) The role of two f-box proteins, SLEEPY1 and SNEEZY, in arabidopsis gibberellin signaling. *Plant Physiol* **155**: 765-775
- Arnaud N, Girin T, Sorefan K, Fuentes S, Wood TA, Lawrenson T, Sablowski R, Ostergaard L** (2010) Gibberellins control fruit patterning in Arabidopsis thaliana. *Genes Dev* **24**: 2127-2132
- Bari R, Jones JD** (2009) Role of plant hormones in plant defence responses. *Plant Mol Biol* **69**: 473-488
- Blázquez MA, Green R, Nilsson O, Sussman MR, Weigel D** (1998) Gibberellins promote flowering of arabidopsis by activating the LEAFY promoter. *Plant Cell* **10**: 791-800
- Bouquin T, Meier C, Foster R, Nielsen ME, Mundy J** (2001) Control of specific gene expression by gibberellin and brassinosteroid. *Plant Physiol* **127**: 450-458
- Brian PW** (1957) The effects of some microbial metabolic products on plant growth. *Symp. Soc. Exp. Biol.* **11**: 166-182
- Cao D, Cheng H, Wu W, Soo HM, Peng J** (2006) Gibberellin mobilizes distinct DELLA-dependent transcriptomes to regulate seed germination and floral development in Arabidopsis. *Plant Physiol* **142**: 509-525

- Cheng H, Qin L, Lee S, Fu X, Richards DE, Cao D, Luo D, Harberd NP, Peng J** (2004) Gibberellin regulates Arabidopsis floral development via suppression of DELLA protein function. *Development* **131**: 1055-1064
- Chiang HH, Hwang I, Goodman HM** (1995) Isolation of the Arabidopsis GA4 locus. *Plant Cell* **7**: 195-201
- Chini A, Fonseca S, Fernandez G, Adie B, Chico JM, Lorenzo O, Garcia-Casado G, Lopez-Vidriero I, Lozano FM, Ponce MR, Micol JL, Solano R** (2007) The JAZ family of repressors is the missing link in jasmonate signalling. *Nature* **448**: 666-671
- Cui H, Levesque MP, Vernoux T, Jung JW, Paquette AJ, Gallagher KL, Wang JY, Blilou I, Scheres B, Benfey PN** (2007) An evolutionarily conserved mechanism delimiting SHR movement defines a single layer of endodermis in plants. *Science* **316**: 421-425
- de Lucas M, Davière JM, Rodríguez-Falcón M, Pontin M, Iglesias-Pedraz JM, Lorrain S, Fankhauser C, Blázquez MA, Titarenko E, Prat S** (2008) A molecular framework for light and gibberellin control of cell elongation. *Nature* **451**: 480-484
- Dill A, Jung HS, Sun TP** (2001) The DELLA motif is essential for gibberellin-induced degradation of RGA. *Proc Natl Acad Sci U S A* **98**: 14162-14167
- Dill A, Sun T** (2001) Synergistic derepression of gibberellin signaling by removing RGA and GAI function in Arabidopsis thaliana. *Genetics* **159**: 777-785
- Dill A, Thomas SG, Hu J, Steber CM, Sun TP** (2004) The Arabidopsis F-box protein SLEEPY1 targets gibberellin signaling repressors for gibberellin-induced degradation. *Plant Cell* **16**: 1392-1405
- Feng S, Martinez C, Gusmaroli G, Wang Y, Zhou J, Wang F, Chen L, Yu L, Iglesias-Pedraz JM, Kircher S, Schafer E, Fu X, Fan LM, Deng XW** (2008) Coordinated regulation of Arabidopsis thaliana development by light and gibberellins. *Nature* **451**: 475-479
- Frigerio M, Alabadí D, Pérez-Gómez J, García-Cárcel L, Phillips AL, Hedden P, Blázquez MA** (2006) Transcriptional regulation of gibberellin metabolism genes by auxin signaling in Arabidopsis. *Plant Physiol* **142**: 553-563
- Fu X, Harberd NP** (2003) Auxin promotes Arabidopsis root growth by modulating gibberellin response. *Nature* **421**: 740-743
- Fu X, Richards DE, Fleck B, Xie D, Burton N, Harberd NP** (2004) The Arabidopsis mutant sleepy1gar2-1 protein promotes plant growth by increasing the affinity of the SCFSLY1 E3 ubiquitin ligase for DELLA protein substrates. *Plant Cell* **16**: 1406-1418
- Garcia-Martinez JL, Lopez-Diaz I, Sanchez-Beltran MJ, Phillips AL, Ward DA, Gaskin P, Hedden P** (1997) Isolation and transcript analysis of gibberellin 20-oxidase genes in pea and bean in relation to fruit development. *Plant Mol Biol* **33**: 1073-1084
- Goto N PR** (1999) Role of gibberellin in the development of floral organs of the gibberellin-deficient mutant, ga1-1, of Arabidopsis thaliana. *Can J Bot* **77**: 944-954
- Griffiths J, Murase K, Rieu I, Zentella R, Zhang ZL, Powers SJ, Gong F, Phillips AL, Hedden P, Sun TP, Thomas SG** (2006) Genetic characterization and functional analysis of the GID1 gibberellin receptors in Arabidopsis. *Plant Cell* **18**: 3399-3414
- Harberd NP, Belfield E, Yasumura Y** (2009) The angiosperm gibberellin-GID1-DELLA growth regulatory mechanism: how an "inhibitor of an inhibitor" enables flexible response to fluctuating environments. *Plant Cell* **21**: 1328-1339
- Harberd NP, Freeling M** (1989) Genetics of dominant gibberellin-insensitive dwarfism in maize. *Genetics* **121**: 827-838
- Hedden P, Phillips AL** (2000) Gibberellin metabolism: new insights revealed by the genes. *Trends Plant Sci* **5**: 523-530
- Helliwell CA, Sullivan JA, Mould RM, Gray JC, Peacock WJ, Dennis ES** (2001) A plastid envelope location of Arabidopsis ent-kaurene oxidase links the plastid and endoplasmic reticulum steps of the gibberellin biosynthesis pathway. *Plant J* **28**: 201-208
- Heo JO, Chang KS, Kim IA, Lee MH, Lee SA, Song SK, Lee MM, Lim J** (2011) Funneling of gibberellin signaling by the GRAS transcription regulator scarecrow-like 3 in the Arabidopsis root. *Proc Natl Acad Sci U S A* **108**: 2166-2171

- Hirano K, Asano K, Tsuji H, Kawamura M, Mori H, Kitano H, Ueguchi-Tanaka M, Matsuoka M** (2010) Characterization of the Molecular Mechanism Underlying Gibberellin Perception Complex Formation in Rice. *Plant Cell*
- Hoffmann-Benning S, Kende H** (1992) On the role of abscisic Acid and gibberellin in the regulation of growth in rice. *Plant Physiol* **99**: 1156-1161
- Hou X, Hu WW, Shen L, Lee LY, Tao Z, Han JH, Yu H** (2008) Global Identification of DELLA Target Genes during Arabidopsis Flower Development. *Plant Physiol*
- Hou X, Lee LY, Xia K, Yan Y, Yu H** (2010) DELLAs modulate jasmonate signaling via competitive binding to JAZs. *Dev Cell* **19**: 884-894
- Ikeda A, Ueguchi-Tanaka M, Sonoda Y, Kitano H, Koshioka M, Futsuhara Y, Matsuoka M, Yamaguchi J** (2001) slender rice, a constitutive gibberellin response mutant, is caused by a null mutation of the SLR1 gene, an ortholog of the height-regulating gene GAI/RGA/RHT/D8. *Plant Cell* **13**: 999-1010
- Itoh H, Ueguchi-Tanaka M, Sato Y, Ashikari M, Matsuoka M** (2002) The gibberellin signaling pathway is regulated by the appearance and disappearance of SLENDER RICE1 in nuclei. *Plant Cell* **14**: 57-70
- Itoh H, Ueguchi-Tanaka M, Sentoku N, Kitano H, Matsuoka M, Kobayashi M** (2001) Cloning and functional analysis of two gibberellin 3 beta -hydroxylase genes that are differently expressed during the growth of rice. *Proc Natl Acad Sci U S A* **98**: 8909-8914
- Iuchi S, Suzuki H, Kim YC, Iuchi A, Kuromori T, Ueguchi-Tanaka M, Asami T, Yamaguchi I, Matsuoka M, Kobayashi M, Nakajima M** (2007) Multiple loss-of-function of Arabidopsis gibberellin receptor AtGID1s completely shuts down a gibberellin signal. *Plant J* **50**: 958-966
- Jager CE, Symons GM, Ross JJ, Smith JJ, Reid JB** (2005) The brassinosteroid growth response in pea is not mediated by changes in gibberellin content. *Planta* **221**: 141-148
- Jaillais Y, Chory J** (2010) Unraveling the paradoxes of plant hormone signaling integration. *Nat Struct Mol Biol* **17**: 642-645
- Josse EM, Gan Y, Bou-Torrent J, Stewart KL, Gilday AD, Jeffree CE, Vaistij FE, Martinez-Garcia JF, Nagy F, Graham IA, Halliday KJ** (2011) A DELLA in Disguise: SPATULA Restrains the Growth of the Developing Arabidopsis Seedling. *Plant Cell*
- King KE, Moritz T, Harberd NP** (2001) Gibberellins are not required for normal stem growth in Arabidopsis thaliana in the absence of GAI and RGA. *Genetics* **159**: 767-776
- Koornneef M, Elgersma, A., Hanhart, C.J., van Loenen-Martinet, E.P., van Rign, L., and Zeevaart, J.A.D.** (1985) A gibberellin insensitive mutant of Arabidopsis thaliana. *Physiol. Plant* **65**: 33-39
- Lee DJ, Zeevaart JA** (2005) Molecular cloning of GA 2-oxidase3 from spinach and its ectopic expression in Nicotiana glauca. *Plant Physiol* **138**: 243-254
- Lee J, He K, Stolz V, Lee H, Figueroa P, Gao Y, Tongprasit W, Zhao H, Lee I, Deng XW** (2007) Analysis of transcription factor HY5 genomic binding sites revealed its hierarchical role in light regulation of development. *Plant Cell* **19**: 731-749
- Lee S, Cheng H, King KE, Wang W, He Y, Hussain A, Lo J, Harberd NP, Peng J** (2002) Gibberellin regulates Arabidopsis seed germination via RGL2, a GAI/RGA-like gene whose expression is up-regulated following imbibition. *Genes Dev* **16**: 646-658
- Leivar P, Monte E, Oka Y, Liu T, Carle C, Castillon A, Huq E, Quail PH** (2008) Multiple phytochrome-interacting bHLH transcription factors repress premature seedling photomorphogenesis in darkness. *Curr Biol* **18**: 1815-1823
- Leivar P, Tepperman JM, Monte E, Calderon RH, Liu TL, Quail PH** (2009) Definition of Early Transcriptional Circuitry Involved in Light-Induced Reversal of PIF-Imposed Repression of Photomorphogenesis in Young Arabidopsis Seedlings. *Plant Cell* **21**: 3535-3553
- Levesque MP, Vernoux T, Busch W, Cui H, Wang JY, Blilou I, Hassan H, Nakajima K, Matsumoto N, Lohmann JU, Scheres B, Benfey PN** (2006) Whole-genome analysis of the SHORT-ROOT developmental pathway in Arabidopsis. *PLoS Biol* **4**: e143

- McGinnis KM, Thomas SG, Soule JD, Strader LC, Zale JM, Sun TP, Steber CM** (2003) The Arabidopsis SLEEPY1 gene encodes a putative F-box subunit of an SCF E3 ubiquitin ligase. *Plant Cell* **15**: 1120-1130
- Murase K, Hirano Y, Sun TP, Hakoshima T** (2008) Gibberellin-induced DELLA recognition by the gibberellin receptor GID1. *Nature* **456**: 459-463
- Nakajima M, Shimada A, Takashi Y, Kim YC, Park SH, Ueguchi-Tanaka M, Suzuki H, Katoh E, Iuchi S, Kobayashi M, Maeda T, Matsuoka M, Yamaguchi I** (2006) Identification and characterization of Arabidopsis gibberellin receptors. *Plant J* **46**: 880-889
- Navarro L, Bari R, Achard P, Lison P, Nemri A, Harberd NP, Jones JD** (2008) DELLAs control plant immune responses by modulating the balance of jasmonic acid and salicylic acid signaling. *Curr Biol* **18**: 650-655
- Nelson DR, Schuler MA, Paquette SM, Werck-Reichhart D, Bak S** (2004) Comparative genomics of rice and Arabidopsis. Analysis of 727 cytochrome P450 genes and pseudogenes from a monocot and a dicot. *Plant Physiol* **135**: 756-772
- Nemhauser JL, Hong F, Chory J** (2006) Different plant hormones regulate similar processes through largely nonoverlapping transcriptional responses. *Cell* **126**: 467-475
- Ni M, Tepperman JM, Quail PH** (1998) PIF3, a phytochrome-interacting factor necessary for normal photoinduced signal transduction, is a novel basic helix-loop-helix protein. *Cell* **95**: 657-667
- O'Neill DP, Ross JJ** (2002) Auxin regulation of the gibberellin pathway in pea. *Plant Physiol* **130**: 1974-1982
- Ogawa M, Hanada A, Yamauchi Y, Kuwahara A, Kamiya Y, Yamaguchi S** (2003) Gibberellin biosynthesis and response during Arabidopsis seed germination. *Plant Cell* **15**: 1591-1604
- Ogawa M, Kusano T, Katsumi M, Sano H** (2000) Rice gibberellin-insensitive gene homolog, OsGAI, encodes a nuclear-localized protein capable of gene activation at transcriptional level. *Gene* **245**: 21-29
- Oh E, Yamaguchi S, Kamiya Y, Bae G, Chung WI, Choi G** (2006) Light activates the degradation of PIL5 protein to promote seed germination through gibberellin in Arabidopsis. *Plant J* **47**: 124-139
- Olszewski N, Sun TP, Gubler F** (2002) Gibberellin signaling: biosynthesis, catabolism, and response pathways. *Plant Cell* **14 Suppl**: S61-80
- Peng J, Carol P, Richards DE, King KE, Cowling RJ, Murphy GP, Harberd NP** (1997) The Arabidopsis GAI gene defines a signaling pathway that negatively regulates gibberellin responses. *Genes Dev* **11**: 3194-3205
- Peng J, Harberd NP** (1993) Derivative Alleles of the Arabidopsis Gibberellin-Insensitive (gai) Mutation Confer a Wild-Type Phenotype. *Plant Cell* **5**: 351-360
- Peng J, Harberd NP** (1997) Gibberellin deficiency and response mutations suppress the stem elongation phenotype of phytochrome-deficient mutants of Arabidopsis. *Plant Physiol* **113**: 1051-1058
- Peng J, Richards DE, Hartley NM, Murphy GP, Devos KM, Flintham JE, Beales J, Fish LJ, Worland AJ, Pelica F, Sudhakar D, Christou P, Snape JW, Gale MD, Harberd NP** (1999) 'Green revolution' genes encode mutant gibberellin response modulators. *Nature* **400**: 256-261
- Phillips AL, Ward DA, Uknes S, Appleford NE, Lange T, Huttly AK, Gaskin P, Graebe JE, Hedden P** (1995) Isolation and expression of three gibberellin 20-oxidase cDNA clones from Arabidopsis. *Plant Physiol* **108**: 1049-1057
- Phinney BO WC, Ritzel MB, Neely PM** (1957) Evidence for gibberellin-like substances from flowering plants. *Proc. Nat. Acad. Sci. (U.S.A.)* **43**: 398-404
- Piskurewicz U, Jikumaru Y, Kinoshita N, Nambara E, Kamiya Y, Lopez-Molina L** (2008) The Gibberellic Acid Signaling Repressor RGL2 Inhibits Arabidopsis Seed Germination by Stimulating Abscissic Acid Synthesis and ABI5 Activity. *Plant Cell* **20**: 2729-2745

- Potts WC, Reid, J.B., and Murfet, I.C.** (1985) Internode length in *Pisum*. Gibberellins and the slender phenotype. *Physiol. Plant* **63**: 357-364
- Radley M** (1956) Occurrence of substances similar to gibberellic acid in higher plants. *Nature* **178**: 1070-1071
- Ross JJ, O'Neill DP, Smith JJ, Kerckhoffs LH, Elliott RC** (2000) Evidence that auxin promotes gibberellin A1 biosynthesis in pea. *Plant J* **21**: 547-552
- Sasaki A, Itoh H, Gomi K, Ueguchi-Tanaka M, Ishiyama K, Kobayashi M, Jeong DH, An G, Kitano H, Ashikari M, Matsuoka M** (2003) Accumulation of phosphorylated repressor for gibberellin signaling in an F-box mutant. *Science* **299**: 1896-1898
- Schliemann GSaW** (1994) Gibberellin conjugates: an overview *Plant Growth Reg* **15**: 247-260
- Schneider G, Jensen E, Spray CR, Phinney BO** (1992) Hydrolysis and re-conjugation of gibberellin A20 glucosyl ester by seedlings of *Zea mays* L. *Proc Natl Acad Sci U S A* **89**: 8045-8048
- Schomburg FM, Bizzell CM, Lee DJ, Zeevaart JA, Amasino RM** (2003) Overexpression of a novel class of gibberellin 2-oxidases decreases gibberellin levels and creates dwarf plants. *Plant Cell* **15**: 151-163
- Seo M, Hanada A, Kuwahara A, Endo A, Okamoto M, Yamauchi Y, North H, Marion-Poll A, Sun TP, Koshida T, Kamiya Y, Yamaguchi S, Nambara E** (2006) Regulation of hormone metabolism in *Arabidopsis* seeds: phytochrome regulation of abscisic acid metabolism and abscisic acid regulation of gibberellin metabolism. *Plant J* **48**: 354-366
- Shimada A, Ueguchi-Tanaka M, Nakatsu T, Nakajima M, Naoe Y, Ohmiya H, Kato H, Matsuoka M** (2008) Structural basis for gibberellin recognition by its receptor GID1. *Nature* **456**: 520-523
- Silverstone AL, Ciampaglio CN, Sun T** (1998) The *Arabidopsis* RGA gene encodes a transcriptional regulator repressing the gibberellin signal transduction pathway. *Plant Cell* **10**: 155-169
- Silverstone AL, Jung HS, Dill A, Kawaide H, Kamiya Y, Sun TP** (2001) Repressing a repressor: gibberellin-induced rapid reduction of the RGA protein in *Arabidopsis*. *Plant Cell* **13**: 1555-1566
- Singh DP, Jermakow AM, Swain SM** (2002) Gibberellins are required for seed development and pollen tube growth in *Arabidopsis*. *Plant Cell* **14**: 3133-3147
- Spray CR, Kobayashi M, Suzuki Y, Phinney BO, Gaskin P, MacMillan J** (1996) The dwarf-1 (dt) Mutant of *Zea mays* blocks three steps in the gibberellin-biosynthetic pathway. *Proc Natl Acad Sci U S A* **93**: 10515-10518
- Stavang JA, Gallego-Bartolome J, Gomez MD, Yoshida S, Asami T, Olsen JE, Garcia-Martinez JL, Alabadi D, Blazquez MA** (2009) Hormonal regulation of temperature-induced growth in *Arabidopsis*. *Plant J* **60**: 589-601
- Sun T-pK, Yuji** (1997) Regulation and cellular localization of ent-kaurene synthesis *Physiologia Plantarum* **101**: 701-708
- Sun TP, Kamiya Y** (1994) The *Arabidopsis* GA1 locus encodes the cyclase ent-kaurene synthetase A of gibberellin biosynthesis. *Plant Cell* **6**: 1509-1518
- Thomas SG, Phillips AL, Hedden P** (1999) Molecular cloning and functional expression of gibberellin 2- oxidases, multifunctional enzymes involved in gibberellin deactivation. *Proc Natl Acad Sci U S A* **96**: 4698-4703
- Ueguchi-Tanaka M, Ashikari M, Nakajima M, Itoh H, Katoh E, Kobayashi M, Chow TY, Hsing YI, Kitano H, Yamaguchi I, Matsuoka M** (2005) GIBBERELLIN INSENSITIVE DWARF1 encodes a soluble receptor for gibberellin. *Nature* **437**: 693-698
- Varbanova M, Yamaguchi S, Yang Y, McKelvey K, Hanada A, Borochoy R, Yu F, Jikumaru Y, Ross J, Cortes D, Ma CJ, Noel JP, Mander L, Shulaev V, Kamiya Y, Rodermeier S, Weiss D, Pichersky E** (2007) Methylation of gibberellins by *Arabidopsis* GAMT1 and GAMT2. *Plant Cell* **19**: 32-45
- Varner JE CG, Chrispeels MJ.** (1965 ) Gibberellic acid-controlled synthesis of alpha-amylase in barley endosperm. *J Cell Physiol.* **66**: 55-67

- Veen MKaJH** (1980) Induction and analysis of gibberellin sensitive mutants in *Arabidopsis thaliana* (L.) heynh. . Theoretical and Applied Genetics **58**: 257-263
- Wen CK, Chang C** (2002) *Arabidopsis* RGL1 encodes a negative regulator of gibberellin responses. Plant Cell **14**: 87-100
- Willige BC, Ghosh S, Nill C, Zourelidou M, Dohmann EM, Maier A, Schwechheimer C** (2007) The DELLA Domain of GA INSENSITIVE Mediates the Interaction with the GA INSENSITIVE DWARF1A Gibberellin Receptor of *Arabidopsis*. Plant Cell
- Wilson RN, Heckman JW, Somerville CR** (1992) Gibberellin Is Required for Flowering in *Arabidopsis thaliana* under Short Days. Plant Physiol **100**: 403-408
- Wolbang CM, Chandler PM, Smith JJ, Ross JJ** (2004) Auxin from the developing inflorescence is required for the biosynthesis of active gibberellins in barley stems. Plant Physiol **134**: 769-776
- Wolbang CM, Ross JJ** (2001) Auxin promotes gibberellin biosynthesis in decapitated tobacco plants. Planta **214**: 153-157
- Xu YL, Li L, Gage DA, Zeevaart JA** (1999) Feedback regulation of GA5 expression and metabolic engineering of gibberellin levels in *Arabidopsis*. Plant Cell **11**: 927-936
- Yamaguchi S** (2008) Gibberellin metabolism and its regulation. Annu Rev Plant Biol **59**: 225-251
- Yamaguchi S, Smith MW, Brown RG, Kamiya Y, Sun T** (1998) Phytochrome regulation and differential expression of gibberellin 3beta-hydroxylase genes in germinating *Arabidopsis* seeds. Plant Cell **10**: 2115-2126
- Yamamoto Y, Hirai T, Yamamoto E, Kawamura M, Sato T, Kitano H, Matsuoka M, Ueguchi-Tanaka M** (2010) A rice *gid1* suppressor mutant reveals that gibberellin is not always required for interaction between its receptor, GID1, and DELLA proteins. Plant Cell **22**: 3589-3602
- Yamauchi Y, Ogawa M, Kuwahara A, Hanada A, Kamiya Y, Yamaguchi S** (2004) Activation of gibberellin biosynthesis and response pathways by low temperature during imbibition of *Arabidopsis thaliana* seeds. Plant Cell **16**: 367-378
- Yamauchi Y, Takeda-Kamiya N, Hanada A, Ogawa M, Kuwahara A, Seo M, Kamiya Y, Yamaguchi S** (2007) Contribution of gibberellin deactivation by AtGA2ox2 to the suppression of germination of dark-imbibed *Arabidopsis thaliana* seeds. Plant Cell Physiol **48**: 555-561
- Yu H, Ito T, Zhao Y, Peng J, Kumar P, Meyerowitz EM** (2004) Floral homeotic genes are targets of gibberellin signaling in flower development. Proc Natl Acad Sci U S A **101**: 7827-7832
- Zentella R, Zhang ZL, Park M, Thomas SG, Endo A, Murase K, Fleet CM, Jikumaru Y, Nambara E, Kamiya Y, Sun TP** (2007) Global analysis of *della* direct targets in early gibberellin signaling in *Arabidopsis*. Plant Cell **19**: 3037-3057
- Zhang ZL, Ogawa M, Fleet CM, Zentella R, Hu J, Heo JO, Lim J, Kamiya Y, Yamaguchi S, Sun TP** (2011) Scarecrow-like 3 promotes gibberellin signaling by antagonizing master growth repressor DELLA in *Arabidopsis*. Proc Natl Acad Sci U S A **108**: 2160-2165
- Zhu Y, Nomura T, Xu Y, Zhang Y, Peng Y, Mao B, Hanada A, Zhou H, Wang R, Li P, Zhu X, Mander LN, Kamiya Y, Yamaguchi S, He Z** (2006) ELONGATED UPPERMOST INTERNODE encodes a cytochrome P450 monooxygenase that epoxidizes gibberellins in a novel deactivation reaction in rice. Plant Cell **18**: 442-456

## **Objectives**





Current knowledge about hormone signaling seems to establish a clear trend for future work, whose leitmotiv could be the establishment of the molecular models that explain: (1) crosstalk between signaling pathways; and (2) the fact that a single signal triggers different specific responses depending on the spatial and developmental context. Given the characteristics of GA action described in the previous sections, we have chosen GA signaling as a model to address the molecular mechanism for both the **specificity** of GA activity, and also the **crosstalk** between GA and other input signals.


For the first question, redundancy of *DELLA* genes in *Arabidopsis* offers an excellent playground to address the relative importance of promoter vs. coding region divergence in the subfunctionalization of duplicated genes, and its impact in GA signaling. For the second question, our strategy has been to focus in a particular process, known to be concurrently regulated by multiple signals, and dissect the participation of GAs at the molecular level. Such is the case of cell expansion whose regulation involves several players acting in a single spatial domain. Besides GAs, it is known to be regulated by light and by other hormones, including auxin, BRs, and ethylene. But an even more interesting situation is that of differential growth, by which the cells in one side of the organ expand more, compared to the cells in the opposite side. This results in the formation of a curvature, found in tropic responses (phototropism, gravitropism...) and in the apical hook of etiolated seedlings. Therefore, our purpose was to identify direct targets of DELLA proteins in differential growth and to investigate the participation of GAs in this context.

Consequently, the objectives of this work were:

1. To assess the ability of different DELLA proteins to perform each other's role. This would be addressed by expressing two *DELLA* genes under the control of the reciprocal promoters, and analyzing the ability of the chimeras to complement *della* mutant phenotypes (Chapter 1).
2. To identify direct target genes regulated by DELLA proteins in etiolated seedlings, focusing on those known to be related to differential growth. This would be approached through transcriptomic analysis of an inducible version of *gai-1* and subsequent molecular studies (Chapters 2, 3 and 4).



## Chapter 1



# Transcriptional diversification and functional conservation between DELLA proteins in Arabidopsis

Gallego-Bartolomé J, Minguet EG, Marín JA, Prat S, Blázquez MA, Alabadí D.

*Mol Biol Evol* 27(6): 1247-1256 (2010)



## **1.1. Abstract**

Plasticity and robustness of signaling pathways partly rely on genetic redundancy, although the precise mechanism that provides functional specificity to the different redundant elements in a given process is often unknown. In *Arabidopsis*, functional redundancy in gibberellin signaling has been largely attributed to the presence of five members of the DELLA family of transcriptional regulators. Here we demonstrate that two evolutionarily and functionally divergent DELLA proteins, RGL2 and RGA, can perform exchangeable functions when they are expressed under control of the reciprocal promoter. Furthermore, both DELLA proteins display equivalent abilities to interact with PIF4 and with other bHLH transcription factors with a reported role in the control of cell growth and seed germination. Therefore, we propose that functional diversification of *Arabidopsis* DELLA proteins has largely relied on changes in their gene expression patterns rather than in their ability to interact with different regulatory partners, model also supported by a clustering analysis of *DELLA* transcript profiles over a range of organs and growth conditions which revealed specific patterns of expression for each of these genes.

## **1.2. Introduction**

Gene duplications are considered as the major source for variation and the generation of evolutionary novelties (Ohno, 1970). Although the most common fate for duplicated genes is gene loss (Lynch and Conery, 2000), duplicated copies are released from mutational constraints, enabling the evolution of new functions (neofunctionalization) (Ohno, 1970; Taylor and Raes, 2004). Alternatively, both gene duplicates can undergo simultaneous reduction of their activity thereby maintaining the total capacity of the ancestral gene (sub-functionalization) (Force et al., 1999).

In addition to providing genetic robustness against deleterious mutations through functional redundancy, gene duplications seem also to be at the core of the mechanisms that provide an unusually high degree of plasticity and robustness to plant signaling pathways (Smith, 1990; Pickett and Meeks-Wagner, 1995; Casal et al., 2004). For instance, it has been proposed that the multiplicity of responses triggered by auxin is governed by the optimized interaction of more than 20 Aux/IAA-ARF pairs in each cell type (Weijers and Jurgens, 2004; Weijers et al., 2005). However, this phenomenon is less understood in other hormonal pathways, in which signaling is transduced by a set of structurally and phylogenetically related proteins that represent branching points in the action of a single given hormone. Such is the case for protein phosphatases type-2C in abscisic acid signaling (Rodriguez, 1998), the ARR response regulators in cytokinin signaling (To et al., 2004), and the JAZ family of proteins involved in jasmonic acid signaling (Chini et al., 2007) among others.

Much work has been devoted to understand the molecular mechanisms that allow the maintenance of gene duplicates in model organisms. Nonetheless, of particular relevance is establishing the extent of the relative contribution to the functional divergence of paralogous genes of variation in the regulatory sequences versus to those in the coding region. The accumulation of polymorphisms in the coding regions of amylase (Goto et al., 2005) and fatty-acid desaturase genes (Fang et al., 2009) in *Drosophila*, and  $\beta$ -defensins in mice and humans (Maxwell et al., 2003), underscore the importance of rapid variation in the coding sequence of recently duplicated genes to generate functional divergence. However, large-scale analyses of expression divergence among duplicated genes in yeast, plants and humans, provide a less clear-cut view. In particular, it has been shown that half of the recently duplicated

genes in Arabidopsis (Blanc and Wolfe, 2004) or rice (Li et al., 2009) have divergent expression patterns, although no correlation is found between expression divergence and time since duplication (Haberer et al., 2004). Actually, it is likely that functional divergence between duplicated genes occurs through both mechanisms, as suggested by the analysis of paralogs of human transcription factors: if the DNA binding site motifs of the transcription factor paralogs are similar, their expression has diverged, while two paralogs that are highly expressed in a tissue tend to have dissimilar DNA binding site motifs (Singh and Hannenhalli, 2008).

Despite all suggestive evidence based on correlations obtained through genomic analyses of duplicated genes, direct experimental evidence for the relative importance of promoter vs coding sequence divergence is scarce. A significant exception is the demonstration that diversification of cis elements in the promoters have been essential to solve “adaptive conflicts” in the ancestor of the duplicated genes (Hittinger and Carroll, 2007). For this reason we chose to analyze the degree of conservation of the actual molecular activities of members belonging to a small family of Arabidopsis transcriptional regulators.

In Arabidopsis, functional redundancy in GA signaling has been largely attributed to the presence of five members of the DELLA family of nuclear-localized transcriptional regulators: GAI, RGA, RGL1, RGL2 and RGL3. These proteins accumulate under low GA concentrations and act as repressors of GA-activated processes, while a local increase in hormone concentration triggers proteasome-dependent degradation of the DELLA proteins, by the concurrent action of the GA receptor GID1 and the E3 ubiquitin ligase SCF<sup>SLY1</sup> complex (Dill et al., 2004; Griffiths et al., 2006; Nakajima et al., 2006). Molecular genetic analyses have shown that each member of the DELLA family performs specific but also overlapping roles in plant development. For example, GAI and RGA are the main regulators of cell expansion in vegetative tissues, since simultaneous loss of GAI and RGA function suppresses the dwarf phenotype of GA-deficient plants to a large extent (Dill and Sun, 2001; King et al., 2001). However, male fertility is primarily regulated by RGA (but not GAI), together with RGL1 and RGL2 (Cheng et al., 2004; Tyler et al., 2004). On the other hand, RGL2 has been proposed to be the main regulator of germination, since knock-out mutations in *RGL2*, but not the other DELLA genes, allow germination of seeds also when GA synthesis is impaired (Lee et al., 2002; Tyler et al., 2004).

Sequence comparisons show a high degree of conservation between the five DELLA proteins, but also enough differences are observed in their N-terminal third that might justify their different activities (Hussain et al., 2005). On the other hand, it cannot be ruled out that functional specificity of the DELLA genes relies on their different expression profiles, as evidenced by RT-qPCR expression analyses of different organs (Tyler et al., 2004). To distinguish between these two possibilities, we have examined the ability of RGL2 and RGA to perform exchangeable functions, by expressing *RGL2* under the control of the *RGA* promoter and vice versa.

### 1.3. Materials and Methods

#### Plant lines and growth conditions

*Arabidopsis thaliana* mutant plants (*gal-3*, *gal-3 gai-t6*, *gal-3 gai-t6 rga-24*, and *rgl2-1*) were all in the *Ler* ecotype. For germination tests, *Ler* and *rgl2-1* seeds were surface-sterilized and sown on sterile Whatman filter papers placed in plates of half-strength MS medium (Duchefa) with 0.8% w/v agar and 1% w/v sucrose, and stratified at 4°C for 7 days in darkness. *gal*, *gal gai-t6* and *gal gai-t6 rga-24* seeds were sterilized and imbibed at 4°C for 7 days in water containing 20  $\mu$ M GA<sub>3</sub>. Before transfer to MS plates, they were extensively rinsed with sterile water to remove any remaining GA<sub>3</sub>.

Germination took place under continuous white fluorescent light (90–100  $\mu$ mol m<sup>-2</sup> sec<sup>-1</sup>) at 23°C in a Percival growth chamber E-30B. Soil-grown plants were kept at 23°C in cabinets with 16-h photoperiod.

#### Construction of vectors and generation of transgenic lines

For promoter-swapping experiments, 2-kb promoter regions of *RGA* and *RGL2*, which included the 5' untranslated region, were PCR amplified from genomic DNA of wild-type *Ler* plants using the pairs of oligonucleotide primers:

pRGA-F-XbaI (TCTAGATATAACCTCATCCATCTATAG) and pRGA-R-XbaI (TCTAGATTACAAGATCTGATGGAG) for pRGA; and pRGL2-F-XbaI (TCTAGATCAGGATGCGAGGTTAAGAATGG) and pRGL2-R-HindIII (AAGCTTTTACTT TACTTCATGGGT) for pRGL2.



The PCR products were subcloned into pCR2.1 (Invitrogen) and transferred into the MCS of the Gateway<sup>TM</sup> binary vector pSBright (Bensmihen et al., 2004) by *Xba*I digestion in case of pRGA to generate pSBright-pRGA and *Xba*I/*Hind*III for pRGL2 to generate pSBright-pRGL2. The *RGA* and *RGL2* cDNAs were obtained from the REGIA consortium as pDONR201 (Invitrogen) clones. Both cDNAs were subcloned into pSBright-pRGA or -pRGL2 using LR clonase (Invitrogen).

The constructs were introduced into the *Agrobacterium tumefaciens* strain C58 by electroporation, and these were then used to transform the *Arabidopsis gal gai-t6 rga-24* and *rgl2-1* mutants by the floral dip method (Clough and Bent, 1998). Transgenic seedlings in the T<sub>1</sub> and T<sub>2</sub> generations were selected based on their resistance to glufosinate. Transgenic lines with a 3:1 (resistant:sensitive) segregation ratio were selected and at least 19 homozygous lines were identified in the T<sub>3</sub> generation for each construct. Data from two representative lines per construct are shown in this work.

### **Phylogenetic analysis**

Iterative search by tblastn was done on the NCBI public database (<http://www.ncbi.nlm.nih.gov>) with the *Arabidopsis thaliana* DELLA amino acid sequences as baits, and representative full-length sequences were selected. Species-specific databases were also consulted to determine the number of DELLA genes present in each genome. Alignments of protein sequences were done with CLUSTALX (Thompson et al., 1997) and only the informative part of the alignments were used for subsequent analyses. A phylogenetic tree was obtained with the PhyML software v2.4.4 (Guindon and Gascuel, 2003) using the JTT amino acid substitution model as indicated upon comparison of the different models with Prottest (Abascal et al., 2005). The consensus tree was supported by Bootstrap analysis (n=1000). Visualization and manipulation of trees were made with TreeView (Page, 1996).

### **Real-time quantitative RT-PCR**

Total RNA from imbibed seeds was extracted using a modified RNAeasy Mini kit (Qiagen) protocol, where the grinded tissue was previously incubated with 600 µl of RLT-PVP buffer (540RLT buffer from RNAeasy Mini Kit + 60 µl PVP40 10% + 6 µl β-mercaptoethanol) followed by a 30 s centrifugation, recovery of the supernatant, and application to the lilac column. For adult plant tissues, total RNA extraction was carried

out as described previously (Frigerio et al., 2006). cDNA synthesis and quantitative PCR, as well as primer sequences for amplification of GA metabolism and EF1- $\alpha$  genes, have been described previously (Frigerio et al., 2006).

The primers used for the quantitative PCR analysis of mRNA levels of *GFP*, *RGA*, *RGL2*, *GAI*, *ATHB16* and *AtMYB34* were, respectively: GFPqRT-F (TCATATGAAGCGGCACGACTT) and GFPqRT-R (GATGGTCCTCTCCTGCAC GTA); RGAqRT-F (ACTTCGACGGGTACGCAGAT) and RGAqRT-R (TGTCGT CACCGTCGTTCC); RGL2qRT-F (GACGGCGCGTAGAGTTCAC) and RGL2qRT-R (TGCATCCCTTGATTAAGCCC); GAIqRT-F (GCTTATGCAGGCTCTTGCG) and GAIqRT-R (AACCGGAAAACAGGAGGACC); ATHB-16qRT-F (GCGCCGTTCTT AACGACGAAACAA) and ATHB-16qRT-R (TAAGAACTCCCGCCAGTAACCGT); MYB34qRT-F (TTAACCGCGTCGCAAGCAAATACG) and MYB34qRT-R (TTGAGC AATGTGGAGGTCGGAGAA).

#### **Protein extraction and western blot**

Total proteins were extracted by homogenizing seedlings in one volume of cold extraction buffer [50 mM Tris-HCl pH 7.5, 150 mM NaCl, 1 mM EDTA, 1 mM DTT, 10% glycerol, 1 mM PMSF, and 1×complete protease inhibitor cocktail (Roche)]. Extracts were centrifuged at 13000 g for 10 min at 4°C. Protein concentration in the supernatants was quantified by the Bradford assay (Bradford, 1976). Aliquots (40 µg) of denatured total proteins were separated in Precise™ 8% Tris-HEPES-SDS gels (Pierce) and transferred onto PVDF membrane (Bio-Rad). RGA-GFP fusion was detected using the monoclonal anti-GFP antibody (clone JL-8) from Clontech.

#### **Yeast two hybrid assay**

For two-hybrid experiments, truncated forms of RGA and RGL2, lacking the DELLA domain, were PCR amplified from a first strand cDNA of *Ler* seedlings using the following combinations of oligonucleotide primers:

RGA1-F (CACCTGGTTGACTCGCAAGAGAACG) and RGA1-R (GTCAAACCTCA GTACGCCGCCG) for RGA; and RGL2-F (CACCTCGTTGACTCTCAGGAGAC CG) and RGL2-R (GCCGCGACTCAGGCGAGTTTCC) for RGL2.

For PCR amplification of the complete coding regions for the PIF4, PIL2, PIL5 and SPT bHLH factors, the combinations of primers used were:

PIF4-F (CACCATGGAACACCAAGGTTGGAG) and PIF4-R (GGCTCACCAACCTAGTGGTCC) for PIF4; PIL2-F (CACCATGATGTTCTTACCAACCG) and PIL2-R (CAGGGAGAATTCCTTCATCTG) for PIL2; PIL5-F (CACCATGCATCATTTTGTCCTG) and PIL5-R (GTTAACCTGTTGTGTGGTTTC) for PIL5; and SPT-F (CACCATGATATCACAGAGAGAAGAAAG) and SPT-R (GGACACTGTTCAAGTAATTCG) for SPT.

The PCR products were subcloned into the pENTR<sup>TM</sup>/D vector using the pENTR<sup>TM</sup>/D-TOPO<sup>®</sup> cloning kit (Invitrogen) and mobilized by LR clonase (Invitrogen) into the pGBKT7 and pGADT7 Gateway vectors (kindly provided by Marta Boter) generated by inserting the Gateway *ccdB* cassette into the MCS *NdeI*–*XhoI*/*Sall* sites of these vectors.

Constructs were transformed into the AH109 yeast strain (*MATa ura3-52 his3-200 trp1-901 leu2-3,112 gal4Δ gal80Δ LYS2::GAL1<sub>UAS</sub>-GAL1<sub>TATA</sub>-HIS3 GAL2<sub>UAS</sub>-GAL2<sub>TATA</sub>-ADE2 URA3::MEL1<sub>UAS</sub>-MEL1<sub>TATA</sub>-lacZ*), using the lithium acetate/PEG method, and yeast cells containing the different DELLA-BD and bHLH-AD fusion combinations were selected on SD-Leu-Trp and SD-Leu-Trp-His-Ade plates. Each construct was also transformed with the pGBKT7 or pGADT7 empty vectors to test for auto-activation activity. Protein extracts were obtained from the transformed yeast cells and western probed with anti-HA (Roche) and anti-GAL4BD (Santa Cruz) antibodies to ensure proper expression of the protein fusions.

### **Confocal Microscopy**

Seedlings were rinsed for 2 min with 10 µg/ml propidium iodide (PI), then 5 min with water. Fresh stained seedlings were mounted on slides only with water. Images were taken using a Leica TCS SL confocal laser microscope (Leica, <http://www.leica.com>) with excitation at 488 nm. For GFP detection, channel 1 was configured between 500-540 nm; and for PI detection, channel 2 was configured between 590-660 nm.

## 1.4. Results and discussion


### 1.4.1. Phylogenetic relationships of DELLA proteins in Angiosperms

Examination of public gene sequence databases and the available full genome sequences of several plant species indicates that, unlike in monocots, genes encoding DELLA proteins in dicots are frequently duplicated. In some cases, such as the *Brassicaceae*, the genomes contain up to five DELLA genes. To investigate the origin and the possible evolutionary history of these duplications, we assembled full-length sequences of DELLA proteins from different dicots and monocots, and analyzed their phylogenetic relationships. As shown in Figure 1.1A, two large clades (I and II) of DELLA proteins can be found in dicots. However, while species in the Rosids genera such as *Populus*, *Pisum*, *Medicago* and the *Brassicaceae* possess members in both subfamilies, species in the Asterids like *Solanum lycopersicum* and *Latuca sativa* possess either only one DELLA protein or two paralogs, respectively, that always group in clade I. In these cases, the absence of DELLA proteins associated with clade II could be a consequence of the lack of sequence information for these species, although a loss-of-function mutant identified in *S. lycopersicum* in the single reported *DELLA* gene displays a phenotype that covers all the functions attributed to GAs in this organism (Marti et al., 2007; Bassel et al., 2008; Jasinski et al., 2008). Hence, it is likely that it indeed represents the only functional *DELLA* gene in tomato.

The presence of two clades of DELLA proteins suggests that the diversification of this family was initiated by a duplication of a single ancestor within the Rosids. After this event, subsequent independent duplications would have given rise to the variety of DELLA proteins present in different species. In fact, the observation of syntenic regions in the *Arabidopsis* genome that include the five DELLA genes (Figure 1.1B) reveals a possible mechanism for the multiplication of these genes in the *Brassicaceae*, involving the rearrangement of large chromosomal fragments.

### 1.4.2. Expression of chimeric versions of DELLA genes


According to genetic analysis, repeated duplication of DELLA proteins in *Arabidopsis* has been accompanied by certain degree of functional diversification, given that mutants in the different DELLA genes are affected only in a subset of responses regulated by GAs.



(A) Phylogenetic maximum likelihood tree of DELLA proteins displaying two large clades (blue and orange). See Material and Methods for details on sequence analysis. Arabidopsis proteins are in bold. Numbers represent percentage bootstrap value (n=1000). Monocot sequences are depicted in green. (B) Syntenic regions in Arabidopsis chromosomes that include DELLA genes. *At*, *Arabidopsis thaliana*; *Bn*, *Brassica napus*; *Gb*, *Gossypium barbadense*; *Ls*, *Lactuca sativa*; *Mh*, *Malus hupehensis*; *Mt*, *Medicago truncatula*; *Os*, *Oryza sativa*; *Ps*, *Pisum sativum*; *Pt*, *Populus trichocarpa*; *Pv*, *Phaseolus vulgaris*; *Sl*, *Solanum lycopersicum*; *Zm*, *Zea mays*. GAH stands for GAI Homolog. *gp* is the number of gene pairs syntenically conserved between the indicated regions.

Therefore, a 2-kb fragment of the *RGL2* promoter was used to drive the expression of the *RGL2* (R2R2) or *RGA* (R2RA) coding sequences fused to *GFP* as a visualization marker, and the constructs were introduced into the *rgl2-1* mutant.

Nineteen and twenty-one independent transformants were isolated respectively. Given that *RGL2* is expressed predominantly in seeds (Lee et al., 2002; Tyler et al., 2004), the expression of the corresponding transgenes was measured by RT-qPCR in the seeds of the transformants and, based on their expression level, two homozygous lines from each class were selected for further analyses: one representative of the lower-expressing lines, and one for the higher-expressing lines (Figure 1.2B). None of the transgenes displayed significant expression in tissues in which *RGL2* is not expressed (data not shown). Similarly, a 2-kb fragment of the *RGA* promoter was fused to the *RGA* (RARA) or *RGL2* (RAR2) coding sequences and *GFP*, and the resulting constructs were introduced into the triple knockout line *gal-3 gai-t6 rga-24*. Twenty-two and twenty-five independent transformants were isolated. The functionality of the constructs was tested through the detection of the GFP-DELLA proteins in extracts of 7-day-old seedlings and, as expected, their stability was severely reduced after incubation of the seedlings with 50  $\mu$ M GA<sub>3</sub> for 3 h (Fig. 1.2C). Furthermore, the transgenic DELLA fusion proteins displayed nuclear localization (Fig. 1.2D), and two homozygous lines of each class were selected for further analyses.




**Figure 1.2.** Transgenic lines expressing *RGA* and *RGL2* under the control of their own and reciprocal promoters.

(A) Scheme of the constructs used in this study. (B) Expression of *RGL2pro::GFP:DELLA* transgenes, determined by RT-qPCR analysis of *GFP* in *rgl2* mutants. Error bars represent SD of three replicates. (C) Production of functional DELLA protein in transgenic lines expressing *RGApro::GFP:DELLA* in *gal-3 gai-t6 rga-24* mutants, determined by western blot in seedlings with mock and 50  $\mu$ M GA<sub>3</sub> treatments. (D) Nuclear localization of DELLA proteins in the roots of transgenic plants described in (C). Line pRGA-GFP is used as control (Dill et al., 2001). Size bar is 40  $\mu$ m.

#### **1.4.3. Functional substitution of RGA by RGL2**

Loss of *GAI*, encoding *ent*-copalylidiphosphate synthase (Sun and Kamiya, 1994), blocks the early steps in GA biosynthesis and causes severe dwarfism due to the accumulation of DELLA proteins, which affect the size of the shoot, the leaves, the hypocotyls and other organs (Silverstone et al., 2001). This defect cannot be rescued by a mutation of *GAI*, due to redundancy with *RGA*, but simultaneous knockout mutations of both DELLA genes restores growth almost to the size of a wild type (Dill and Sun, 2001; King et al., 2001). On the other hand, the endogenous *RGL2* gene does not have a role in cell expansion, since mutations in this gene do not restore growth (Lee et al., 2002; Tyler et al., 2004), in agreement with the very low expression of *RGL2* in these tissues. However, our analysis of the *RAR2* lines indicates that the RGL2 protein is capable of exerting RGA function when expressed under control of the *RGA* promoter. As shown in Figure 1.3A, expression of RGA under the 2-kb *RGA* promoter could complement the loss of RGA function, to different extent depending on the line examined. Even more interestingly, expression of *RGL2* could also complement the lack of *RGA* function to an equivalent extent, when expressed under the *RGA* promoter. Such complementation ability was observed not only in leaves and shoots (Figures 1.3A and 1.3B), but also in etiolated hypocotyls (results not shown). Actually, the degree of stem dwarfism rescue was dependent on the age of the plant but not on the transgenic DELLA used (Figure 1.3B), an effect that might be caused by the lack of additional regulatory sequences lying beyond the 2-kb promoter fragment used in this study.

RGA is also involved in feed-back regulation of GA metabolism genes in tissues in which *RGL2* is not normally expressed, such as hypocotyls, leaves and shoots (Silverstone et al., 1998; Dill and Sun, 2001; Frigerio et al., 2006). *RGL2*, when expressed under the control of the *RGA* promoter, was able to restore feed-back control of the *GA20ox1*, *GA20ox2* and *GA20ox8* genes, which was lost in the *gai-3 gai-t6 rga-24* mutant (Figure 1.3C). Therefore, the RGL2 protein seems to display the biochemical activity characteristic of RGA function, functional divergence of these genes being caused, at least in this case, by changes in their spatial pattern of expression. To confirm this hypothesis, we performed the reciprocal analysis by testing the ability of RGA to substitute RGL2.



**Figure 1.3. Complementation by *RGL2* of *rga* loss-of-function mutants.**


(A) Photograph of four-week-old plants grown under long days. (B) Plant height at different ages of control plants and plants transformed with *RGAPro::GFP:DELLA*, showing that *RGA* and *RGL2* are capable of inhibiting shoot elongation to a similar extent when expressed under the *RGA* promoter. Error bars represent SD ( $n > 15$ ). (C) Expression of GA metabolism genes subject to regulation by RGA in shoots, measured by RT-qPCR, showing that RGA and RGL2 complement the loss of RGA function. Error bars represent SD of three replicates. Only two representative homozygous transgenic lines are shown for each construct, although equivalent results were observed for over 14 lines.

#### 1.4.4. Functional substitution of RGL2 by RGA

A major function of RGL2 in seed physiology is revealed by the observation that GA-deficient seeds are able to germinate only if RGL2 activity is suppressed, but not when any of the other four *DELLA* genes are mutated (Lee et al., 2002; Cao et al., 2005). This is particularly evident for instance in the *rgl2-1* allele, whose seeds germinate in the presence of 20 or even 120  $\mu$ M PAC, while these inhibitor concentrations reduce germination efficiency of wild-type seeds to values below 5% (Figure 1.4A). As expected, expression of *RGL2* under the control of its 2-kb promoter region caused a decrease in germination of *rgl2-1* in the presence of PAC, although it never reached the values of the wild type (Figure 1.4A). More importantly, expression of *RGA* under the control of the *RGL2* promoter (R2RA lines) also complemented the



*rgl2-1* to an equivalent extent when compared to RGL2, indicating that RGA and RGL2 proteins can perform equivalent functions during seed germination, provided that they are both expressed under the *RGL2* promoter. This conclusion was confirmed at the molecular level by measuring the expression level of *ATHB-16* and *MYB34* in germinating seeds, by RT-qPCR. These genes are normally up-regulated during germination and have been proposed to be repressed by RGL2 in imbibed seeds (Ogawa et al., 2003; Cao et al., 2006). As shown on Figure 1.4B, both marker genes were induced in *rgl2-1* mutant seeds during germination in GA-limiting conditions, compared to seeds of the wild type and the transgenic lines expressing RGL2 and RGA under the control of the *RGL2* promoter.




**Figure 1.4. Complementation by RGA of *rgl2* loss-of-function mutants.**

(A) Germination of wild type and *rgl2* mutant seeds transformed with *RGL2pro::GFP:DELLA* under increasing concentrations of paclobutrazol (PAC). All lines tested germinated with almost 100% efficiency in the absence of PAC.  $n > 200$  seeds. (B) Expression of germination marker genes in seeds imbibed for 11 days in the presence of 20 μM PAC, 6 days after exposure to light. Error bars represent SD of three replicates. Only two representative homozygous transgenic lines are shown for each construct, although equivalent results were observed for over 12 lines.

The observation that the RGA protein can perform the function of RGL2 in seeds seems to be in conflict with the fact that RGA is also expressed in seeds, but suppression of this gene does not allow germination of GA-deficient seeds (Tyler et al., 2004; Cao et al., 2005). Where does the specificity of RGL2 function reside in wild-type seeds? One possibility is that expression of *RGA* is restricted to cell types in the seed which are not relevant for germination. This is unlikely because RGA and GAI have indeed been found to contribute to control seed germination under normal conditions (Cao et al., 2005). Another possibility is that the *RGA* and *RGL2* promoters are differentially regulated under GA deficiency, as suggested by the observation that the RGL2 protein is more abundant than RGA in seeds incubated with PAC (Piskurewicz et al., 2008). To explore this option, we analyzed the expression of *GAI*, *RGA* and *RGL2* during seed imbibition and germination, in the GA-deficient *gal* mutant, and in its corresponding parental wild type. Interestingly, while *RGA* and *RGL2* expression levels were equivalent in the wild type during germination (when seeds are exposed to light), *RGL2* expression was highly predominant in GA-deficient seeds (Figure 1.5A). Besides, elements mediating GA-regulation of the *RGL2* promoter would lie- within the 2-kb fragment used in the R2R2 and R2RA lines, as confirmed by RT-qPCR analysis of the GFP fusion transcript in germinating seeds of the transgenic lines in mock and GA-deficient (20  $\mu$ M PAC) conditions (Figure 1.5B).

#### 1.4.5. Conservation of DELLA protein interactions

The results presented here indicate that a major driving force in the sub-functionalization of DELLA proteins may in fact rely on different expression patterns arisen after duplication, hence pointing to variations in the promoters of the *DELLA* genes as the main cause for the differential roles of DELLAs in plant development. Common tools for promoter analysis did not allow the identification of specific sequences that explain the divergent behaviour of the *DELLA* promoters. However, cluster analysis of the expression of the five Arabidopsis *DELLA* genes using the over 100 conditions microarray data available through Genevestigator (Zimmermann et al., 2004) revealed a topology that faithfully reproduces the phylogenetic structure of the DELLA group (Figure 1.6; see also Figure 1.1A), suggesting an intimate link between the evolution of DELLA function and *DELLA* gene expression patterns.




**Figure 1.5. Expression of *DELLA* genes during seed imbibition and germination.**

(A) Time course of the expression of *GAI*, *RGA* and *RGL2* during imbibition of wild-type and *gal* mutant seeds in darkness (shaded area in the graphs) and after exposure to light, determined by RT-qPCR. (B) Expression of *RGL2pro::GFP:DELLA* transgenes, determined by RT-qPCR analysis of *GFP* in *rgl2* mutants 48 h after exposure of imbibed seeds to light. In this experiment, seeds were incubated with mock or 20μM PAC solutions. Error bars represent SD of three biological replicates.

This view is coherent with the observation that at least one fifth of the alterations responsible for phenotypic evolution in multicellular organisms reside in regulatory regions (Stern and Orgogozo, 2008).

On the other hand, the study of the expression patterns of duplicate genes using microarrays has established a link between expression divergence and coding-sequence divergence in animals (Makova and Li, 2003; Conant and Wagner, 2004; Li et al.,


2005). Since the correlation between these two processes is less clear in *Arabidopsis* (Blanc and Wolfe, 2004), and there is experimental evidence that recently duplicated proteins in *Arabidopsis* undergo rapid changes in protein activity (Tominaga et al., 2007), we decided to investigate the extent of divergence in DELLA protein activity that could have arisen as a result of expression changes.



**Figure 1.6. Cluster analysis of *DELLA* gene expression.**

(A) Expression patterns of *DELLA* genes during plant development and across tissues. (B) Expression of *DELLA* genes in response to biotic and abiotic stress treatments. Multiple microarray data were analyzed with the tools provided by Genevestigator (Zimmermann et al., 2004) (<https://www.genevestigator.ethz.ch>), and the dendrogram displaying the relationships between *DELLA* gene expression patterns was constructed using all the experiments in the database. On the left panel, the intensity of the blue colour is proportional to the level of expression, while on the right panel, green and red colours indicate lower and higher levels, respectively, compared to the control situation in each treatment.

Although biochemical activity of DELLA proteins is not yet fully established, at least two features have been defined that are intimately linked to the protein sequences: (i) they have been proposed to regulate gene expression through protein-protein interactions with transcription factors (TFs) of the bHLH family (de Lucas et al., 2008; Feng et al., 2008); and (ii) they interact physically with the GID1 GA receptors (Nakajima et al., 2006; Ueguchi-Tanaka et al., 2007; Murase et al., 2008). To evaluate the degree of conservation of the biochemical properties of these proteins, we examined the ability of RGA and RGL2 to interact with PIF4 and also with other bHLH TFs of subfamily 15, with a reported role in seed germination control (Penfield et al., 2005; Oh et al., 2006; Kim et al., 2008; Oh et al., 2009). As seen previously, RGA displayed strong interaction with PIF4 and a similar interaction ability was also observed for RGL2 (Figure 1.7). PIF4-DELLA interaction has been reported to involve the bHLH DNA recognition domain, which is highly conserved among members of subfamily 15. Thus, it is possible that this interaction is not restricted to PIF3/PIF4 but involves other members of this gene family. In fact, we observed a strong interaction between RGA and RGL2 and the bHLH proteins PIL5, PIL2 and SPATULA (SPT), suggesting that DELLAs may interact with all members of this TF subfamily.



**Figure 1.7. Conserved interaction between DELLA proteins and bHLH transcription factors.**

(A) Growth of the yeast cells transformed with the DELLA-GAL4BD and bHLH-GAL4AD constructs on SD-Leu-Trp (SD-LT) and SD-Leu-Trp-His-Ade (SD-4) plates. (B) Western blot detection of the GAL4BD and GAL4AD fusion proteins using anti-GAL4BD and anti-HA antibodies.

DELLA proteins have been also described to display differential abilities to interact with the three GID1 receptors in Arabidopsis, based on yeast two-hybrid analyses (Nakajima et al., 2006). Taking into account the reported differential expression pattern of the three Arabidopsis GID1 receptors and the five DELLA genes, it seems likely that GA signaling is governed by combinatorial tissue- and stage-specific expression patterns of DELLA proteins and most likely also by subsequent diversification of their interactors.

Therefore, the experimental evidence presented here demonstrates the existence of at least two levels of regulation that have contributed with different impact to the evolutionary fixation of diversity in GA signaling: first, and more critically, the sub-functionalization of *DELLA* gene expression patterns (either spatially or in terms of regulatory responses), and then a more recent optimization of protein interactions between DELLAs and different sets of TFs (and possibly other proteins) within the cellular context defined by the new expression domain for each DELLA protein. It remains to be assessed if the multiplication of *DELLA* genes in Rosids, and especially in the *Brassicaceae*, represents an improvement in the plasticity of the responses mediated by GAs.

### Acknowledgements

We deeply appreciate the help of Marta Trénor and Laura García-Cárcel in the initial stages of this work. We also thank Tai-ping Sun (Duke University) and the Arabidopsis Biological Resource Center for seeds, Marta Boter for the pGBKT7 and pGADT7 Gateway vectors, Santiago Elena (IBMCP, CSIC-UPV) for useful comments on the manuscript, and François Percy (IRTSV, CNRS-CEA) for fruitful discussions and hosting MAB. Work in the authors' laboratories is funded by grants BIO2007-60923 and BIO2005-07284 from the Spanish Ministry of Science and Innovation. JGB is the recipient of a CSIC I3P Fellowship and JAM is the recipient of a Fellowship from the Fundación "la Caixa".

## 1.5. Bibliography

Abascal F, Zardoya R, Posada D (2005) ProtTest: selection of best-fit models of protein evolution. *Bioinformatics* **21**: 2104-2105

- Bassel GW, Mullen RT, Bewley JD** (2008) Procera is a putative DELLA mutant in tomato (*Solanum lycopersicum*): effects on the seed and vegetative plant. *J Exp Bot* **59**: 585-593
- Bensmihen S, To A, Lambert G, Kroj T, Giraudat J, Parcy F** (2004) Analysis of an activated ABI5 allele using a new selection method for transgenic *Arabidopsis* seeds. *FEBS Lett* **561**: 127-131
- Blanc G, Wolfe KH** (2004) Functional divergence of duplicated genes formed by polyploidy during *Arabidopsis* evolution. *Plant Cell* **16**: 1679-1691
- Bradford MM** (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* **72**: 248-254
- Cao D, Cheng H, Wu W, Soo HM, Peng J** (2006) Gibberellin mobilizes distinct DELLA-dependent transcriptomes to regulate seed germination and floral development in *Arabidopsis*. *Plant Physiol* **142**: 509-525
- Cao D, Hussain A, Cheng H, Peng J** (2005) Loss of function of four DELLA genes leads to light- and gibberellin-independent seed germination in *Arabidopsis*. *Planta* **223**: 105-113
- Casal JJ, Fankhauser C, Coupland G, Blázquez MA** (2004) Signalling for developmental plasticity. *Trends Plant Sci* **9**: 309-314
- Cheng H, Qin L, Lee S, Fu X, Richards DE, Cao D, Luo D, Harberd NP, Peng J** (2004) Gibberellin regulates *Arabidopsis* floral development via suppression of DELLA protein function. *Development* **131**: 1055-1064
- Chini A, Fonseca S, Fernandez G, Adie B, Chico JM, Lorenzo O, Garcia-Casado G, Lopez-Vidriero I, Lozano FM, Ponce MR, Micol JL, Solano R** (2007) The JAZ family of repressors is the missing link in jasmonate signalling. *Nature* **448**: 666-671
- Clough SJ, Bent AF** (1998) Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J* **16**: 735-743
- Conant GC, Wagner A** (2004) Duplicate genes and robustness to transient gene knock-downs in *Caenorhabditis elegans*. *Proc Biol Sci* **271**: 89-96
- de Lucas M, Davière JM, Rodríguez-Falcón M, Pontin M, Iglesias-Pedraz JM, Lorrain S, Fankhauser C, Blázquez MA, Titarenko E, Prat S** (2008) A molecular framework for light and gibberellin control of cell elongation. *Nature* **451**: 480-484
- Dill A, Jung HS, Sun TP** (2001) The DELLA motif is essential for gibberellin-induced degradation of RGA. *Proc Natl Acad Sci U S A* **98**: 14162-14167
- Dill A, Sun T** (2001) Synergistic derepression of gibberellin signaling by removing RGA and GAI function in *Arabidopsis thaliana*. *Genetics* **159**: 777-785
- Dill A, Thomas SG, Hu J, Steber CM, Sun TP** (2004) The *Arabidopsis* F-box protein SLEEPY1 targets gibberellin signaling repressors for gibberellin-induced degradation. *Plant Cell* **16**: 1392-1405
- Fang S, Ting CT, Lee CR, Chu KH, Wang CC, Tsaor SC** (2009) Molecular evolution and functional diversification of fatty acid desaturases after recurrent gene duplication in *Drosophila*. *Mol Biol Evol* **26**: 1447-1456
- Feng S, Martinez C, Gusmaroli G, Wang Y, Zhou J, Wang F, Chen L, Yu L, Iglesias-Pedraz JM, Kircher S, Schafer E, Fu X, Fan LM, Deng XW** (2008) Coordinated regulation of *Arabidopsis thaliana* development by light and gibberellins. *Nature* **451**: 475-479
- Force A, Lynch M, Pickett FB, Amores A, Yan YL, Postlethwait J** (1999) Preservation of duplicate genes by complementary, degenerative mutations. *Genetics* **151**: 1531-1545
- Frigerio M, Alabadi D, Pérez-Gómez J, García-Cárcel L, Phillips AL, Hedden P, Blázquez MA** (2006) Transcriptional regulation of gibberellin metabolism genes by auxin signaling in *Arabidopsis*. *Plant Physiol* **142**: 553-563
- Goto H, Szmids AE, Yamazaki T, Inomata N** (2005) Effect of nucleotide polymorphism in cis-regulatory and coding regions on amylase activity and fitness in *Drosophila melanogaster*. *Heredity* **95**: 369-376

- Griffiths J, Murase K, Rieu I, Zentella R, Zhang ZL, Powers SJ, Gong F, Phillips AL, Hedden P, Sun TP, Thomas SG** (2006) Genetic characterization and functional analysis of the GID1 gibberellin receptors in Arabidopsis. *Plant Cell* **18**: 3399-3414
- Guindon S, Gascuel O** (2003) A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Syst Biol* **52**: 696-704
- Haberer G, Hindemitt T, Meyers BC, Mayer KF** (2004) Transcriptional similarities, dissimilarities, and conservation of cis-elements in duplicated genes of Arabidopsis. *Plant Physiol* **136**: 3009-3022
- Hittinger CT, Carroll SB** (2007) Gene duplication and the adaptive evolution of a classic genetic switch. *Nature* **449**: 677-681
- Hussain A, Cao D, Cheng H, Wen Z, Peng J** (2005) Identification of the conserved serine/threonine residues important for gibberellin-sensitivity of Arabidopsis RGL2 protein. *Plant J* **44**: 88-99
- Jasinski S, Tattersall A, Piazza P, Hay A, Martinez-Garcia JF, Schmitz G, Theres K, McCormick S, Tsiantis M** (2008) PROCERA encodes a DELLA protein that mediates control of dissected leaf form in tomato. *Plant J* **56**: 603-612
- Kim DH, Yamaguchi S, Lim S, Oh E, Park J, Hanada A, Kamiya Y, Choi G** (2008) SOMNUS, a CCCH-type zinc finger protein in Arabidopsis, negatively regulates light-dependent seed germination downstream of PIL5. *Plant Cell* **20**: 1260-1277
- King KE, Moritz T, Harberd NP** (2001) Gibberellins are not required for normal stem growth in Arabidopsis thaliana in the absence of GAI and RGA. *Genetics* **159**: 767-776
- Lee S, Cheng H, King KE, Wang W, He Y, Hussain A, Lo J, Harberd NP, Peng J** (2002) Gibberellin regulates Arabidopsis seed germination via RGL2, a GAI/RGA-like gene whose expression is up-regulated following imbibition. *Genes Dev* **16**: 646-658
- Li WH, Yang J, Gu X** (2005) Expression divergence between duplicate genes. *Trends Genet* **21**: 602-607
- Li Z, Zhang H, Ge S, Gu X, Gao G, Luo J** (2009) Expression pattern divergence of duplicated genes in rice. *BMC Bioinformatics* **10 Suppl 6**: S8
- Lynch M, Conery JS** (2000) The evolutionary fate and consequences of duplicate genes. *Science* **290**: 1151-1155
- Makova KD, Li WH** (2003) Divergence in the spatial pattern of gene expression between human duplicate genes. *Genome Res* **13**: 1638-1645
- Marti C, Orzaez D, Ellul P, Moreno V, Carbonell J, Granell A** (2007) Silencing of DELLA induces facultative parthenocarpy in tomato fruits. *Plant J* **52**: 865-876
- Maxwell AI, Morrison GM, Dorin JR** (2003) Rapid sequence divergence in mammalian beta-defensins by adaptive evolution. *Mol Immunol* **40**: 413-421
- Murase K, Hirano Y, Sun TP, Hakoshima T** (2008) Gibberellin-induced DELLA recognition by the gibberellin receptor GID1. *Nature* **456**: 459-463
- Nakajima M, Shimada A, Takashi Y, Kim YC, Park SH, Ueguchi-Tanaka M, Suzuki H, Katoh E, Iuchi S, Kobayashi M, Maeda T, Matsuoka M, Yamaguchi I** (2006) Identification and characterization of Arabidopsis gibberellin receptors. *Plant J* **46**: 880-889
- Ogawa M, Hanada A, Yamauchi Y, Kuwahara A, Kamiya Y, Yamaguchi S** (2003) Gibberellin biosynthesis and response during Arabidopsis seed germination. *Plant Cell* **15**: 1591-1604
- Oh E, Kang H, Yamaguchi S, Park J, Lee D, Kamiya Y, Choi G** (2009) Genome-Wide Analysis of Genes Targeted by PHYTOCHROME INTERACTING FACTOR 3-LIKE5 during Seed Germination in Arabidopsis. *Plant Cell* **21**: 403-419
- Oh E, Yamaguchi S, Kamiya Y, Bae G, Chung WI, Choi G** (2006) Light activates the degradation of PIL5 protein to promote seed germination through gibberellin in Arabidopsis. *Plant J* **47**: 124-139
- Ohno S** (1970) Evolution by gene duplication. Springer, New York
- Page RD** (1996) TreeView: an application to display phylogenetic trees on personal computers. *Comput Appl Biosci* **12**: 357-358



- Penfield S, Josse EM, Kannangara R, Gilday AD, Halliday KJ, Graham IA** (2005) Cold and light control seed germination through the bHLH transcription factor SPATULA. *Curr Biol* **15**: 1998-2006
- Pickett FB, Meeks-Wagner D** (1995) Seeing double: appreciating genetic redundancy. *Plant Cell* **7**: 1347-1356
- Piskurewicz U, Jikumaru Y, Kinoshita N, Nambara E, Kamiya Y, Lopez-Molina L** (2008) The Gibberellic Acid Signaling Repressor RGL2 Inhibits Arabidopsis Seed Germination by Stimulating Abscissic Acid Synthesis and ABI5 Activity. *Plant Cell* **20**: 2729-2745
- Rodriguez PL** (1998) Protein phosphatase 2C (PP2C) function in higher plants. *Plant Mol Biol* **38**: 919-927
- Silverstone AL, Ciampaglio CN, Sun T** (1998) The Arabidopsis RGA gene encodes a transcriptional regulator repressing the gibberellin signal transduction pathway. *Plant Cell* **10**: 155-169
- Silverstone AL, Jung HS, Dill A, Kawaide H, Kamiya Y, Sun TP** (2001) Repressing a repressor: gibberellin-induced rapid reduction of the RGA protein in Arabidopsis. *Plant Cell* **13**: 1555-1566
- Singh LN, Hannenhalli S** (2008) Functional diversification of paralogous transcription factors via divergence in DNA binding site motif and in expression. *PLoS One* **3**: e2345
- Smith H** (1990) Signal perception, differential expression within multigene families and the molecular basis for phenotypic plasticity. *Plant Cell Environ* **13**: 585-594
- Stern DL, Orgogozo V** (2008) The loci of evolution: how predictable is genetic evolution? *Evolution* **62**: 2155-2177
- Sun TP, Kamiya Y** (1994) The Arabidopsis GA1 locus encodes the cyclase ent-kaurene synthetase A of gibberellin biosynthesis. *Plant Cell* **6**: 1509-1518
- Taylor JS, Raes J** (2004) Duplication and divergence: the evolution of new genes and old ideas. *Annu Rev Genet* **38**: 615-643
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG** (1997) The CLUSTAL\_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* **25**: 4876-4882
- To JP, Haberer G, Ferreira FJ, Deruere J, Mason MG, Schaller GE, Alonso JM, Ecker JR, Kieber JJ** (2004) Type-A Arabidopsis response regulators are partially redundant negative regulators of cytokinin signaling. *Plant Cell* **16**: 658-671
- Tominaga R, Iwata M, Okada K, Wada T** (2007) Functional analysis of the epidermal-specific MYB genes CAPRICE and WEREWOLF in Arabidopsis. *Plant Cell* **19**: 2264-2277
- Tyler L, Thomas SG, Hu J, Dill A, Alonso JM, Ecker JR, Sun TP** (2004) DELLA proteins and gibberellin-regulated seed germination and floral development in Arabidopsis. *Plant Physiol* **135**: 1008-1019
- Ueguchi-Tanaka M, Nakajima M, Katoh E, Ohmiya H, Asano K, Saji S, Hongyu X, Ashikari M, Kitano H, Yamaguchi I, Matsuoka M** (2007) Molecular interactions of a soluble gibberellin receptor, GID1, with a rice DELLA protein, SLR1, and gibberellin. *Plant Cell* **19**: 2140-2155
- Weijers D, Benkova E, Jager KE, Schlereth A, Hamann T, Kientz M, Wilmoth JC, Reed JW, Jurgens G** (2005) Developmental specificity of auxin response by pairs of ARF and Aux/IAA transcriptional regulators. *Embo J* **24**: 1874-1885
- Weijers D, Jurgens G** (2004) Funneling auxin action: specificity in signal transduction. *Curr Opin Plant Biol* **7**: 687-693
- Zimmermann P, Hirsch-Hoffmann M, Hennig L, Gruissem W** (2004) GENEVESTIGATOR. Arabidopsis microarray database and analysis toolbox. *Plant Physiol* **136**: 2621-2632



## Chapter 2



# **Transcriptional regulation by DELLA proteins during etiolated development in *Arabidopsis thaliana***

Gallego-Bartolomé J, Alabadí D, Blázquez MA



## **2.1. Introduction**

Plants are sessile organisms that cannot change their location as a strategy to optimize their access to energy sources or in response to the environment. Thus, adjusting their growth and choosing the correct developmental program has to be precise and robust otherwise chances of survival could be reduced. This need has forced the development of very sophisticated sensing mechanisms and signal transduction pathways to respond properly to fluctuating environmental conditions. Plant hormones play an instructive role on this as they control many, if not all, developmental responses in plants (Alabadi et al., 2009; Jaillais and Chory, 2010).

Gibberellins (GAs) are one of the classical plant hormones. They regulate several processes during the plant life cycle such as germination, vegetative growth or flowering (Yamaguchi, 2008) through gene transcriptional regulation (Ogawa et al., 2003; Cao et al., 2006; Zentella et al., 2007; Hou et al., 2008). This transcriptional regulation relies on the activity of the nuclear, GA-regulated DELLA proteins (Harberd et al., 2009). In brief, DELLAs accumulate in the absence of GAs blocking the transcriptional response to the hormone. When GA levels increase, the binding of the hormone to its receptor, GID1, promotes the formation of a GA-GID1-DELLA complex (Ueguchi-Tanaka et al., 2005; Shimada et al., 2008) that favors the recognition of the DELLA protein by the SCF<sup>SLY</sup> ubiquitin ligase (Hirano et al., 2010) and the subsequent ubiquitination. This modification leads to DELLA degradation by the 26S proteasome (Fu et al., 2002; Itoh et al., 2003) and transcriptional changes to the hormone take place.

Two observations support the idea that DELLAs are transcriptional regulators: first, chromatin immunoprecipitation (ChIP) experiments reveal that DELLAs sit at the vicinity of promoters of certain GA-regulated genes (Zentella et al., 2007; Zhang et al., 2011). Second, DELLAs interact physically with transcription factors and other transcriptional regulators. For example, they interact with bHLH transcription factors of the PIF clade and inhibit their ability to bind DNA (de Lucas et al., 2008; Feng et al., 2008), as well as with other members of the bHLH family (Arnaud et al., 2010; Gallego-Bartolome et al., 2010). Also, they interact with JAZ proteins, which are transcriptional regulators that negatively regulate jasmonate signaling (Hou et al., 2010), and with SCL3, a transcriptional regulator that belongs to the GRAS family (Heo et al., 2011; Zhang et al., 2011). In addition, genetic evidence indicates that the bZIP

transcription factor HY5 mediates the promotion of photomorphogenesis by DELLA (Alabadi et al., 2008).

Despite these recent advances, we still lack a broader view of the mechanisms by which DELLA proteins regulate the large variety of GA responses. A bottom-up strategy to dissect further this fundamental aspect of GA signaling is to identify and classify GA target genes according to their expression domain or the process in which they participate. In this regard, global analyses of DELLA-regulated transcription in two different developmental contexts –vegetative growth and floral development– have shown that only 3.6% of the target genes are shared between the two sets (Zentella et al., 2007; Hou et al., 2008). This observation underscores the importance of the developmental context in which GA signaling is investigated.


GAs are important regulators of the skotomorphogenic developmental program (Alabadi et al., 2004; Alabadi et al., 2008). In order to dissect how GAs regulate this process, we have searched for early target genes of DELLAs in etiolated seedlings. For that purpose, we have examined global, rapid changes in gene expression after compromising the GA signaling pathway in dark-grown seedlings. This approach allowed us 1) to identify which cellular pathways are directly regulated by GAs to promote skotomorphogenesis; and 2) to identify gene targets that will serve as markers to further dissect the mechanisms by which DELLAs regulate gene expression.

## 2.2. Results & Discussion

### 2.2.1. Identification of genes rapidly regulated by GAI in etiolated seedlings

We sought to identify in a global and unbiased way genes whose expression was modulated rapidly in response to a change in GA activity in etiolated seedlings by using a transgenic line that expresses a gain-of-function version of the DELLA protein GAI under the control of a temperature-inducible promoter, *HS::gai-1* (Alabadi et al., 2008). To determine the optimal duration of the heat treatment needed to strongly induce *gai-1* transcript accumulation, we placed 2-day-old etiolated *HS::gai-1* seedlings at 37°C for 30, 60, and 120 minutes, and then analyzed expression of the transgene by qRT-PCR over a time-course (Figure 2.1A). The 30-min treatment was sufficient to strongly and transiently induce *gai-1* transcript accumulation. To confirm that the inductive treatment resulted in an increase of GAI activity, we checked the expression of the *GA20ox2* and *GA3ox1* genes, that encode key enzymes in the GA biosynthetic pathway subject to

feedback regulation by DELLA proteins (Peng et al., 1997; Dill et al., 2001; Zentella et al., 2007; Arana et al., 2011). As expected, transcripts of both genes accumulated strongly in seedlings following the heat shock, but only in the 30-min treatment was this accumulation transitory (Figures 2.1B and C); moreover, expression of these genes did not change significantly in response to the temperature treatment in wild-type seedlings (data not shown).



**Figure 2.1** Effect of transient *gai-1* induction on known DELLA target genes.

Two-day-old, etiolated *HS::gai-1* and wild type Col-0 plants grown at 22°C received a 37°C heat-shock treatment for different periods (30, 60, 120 min) and then returned to 22°C. Samples were collected at the indicated times. Expression of the transgene (A), as well as of *AtGA20ox2* (B) and *AtGA3ox1* (C) genes was monitored by RT-qPCR.

Given that the induction protocol was appropriate to modulate the expression of GAI target genes, we interrogated the transcriptome of two-day-old etiolated *HS::gai-1* seedlings at 0, 1, 2, and 4 hours after starting a 30-min heat shock at 37°C. Expression was compared using triplicate RNA samples from whole transgenic seedlings at each time point and the corresponding wild-type seedlings by hybridization of 70-mer oligonucleotide arrays representing the majority of the *Arabidopsis* genes (<http://www.ag.arizona.edu/microarray>). The application of a Significance Analysis of

Microarrays criterium (Tusher et al., 2001) with a false discovery rate of 8.74% and a 1.5-fold cutoff value allowed us to identify 148 genes differentially expressed during the first four hours after the induction of *gai-1* activity. This list represented the genes putatively regulated by GAI in etiolated seedlings (Supplemental Table 1); among them, 58 were downregulated and 90 induced (Figure 2.2A).


### **2.2.2. Comparison of DELLA-regulated genes in different developmental situations**

Recent studies have identified by a similar approach early target genes of the Arabidopsis DELLA protein RGA in aerial tissue of light-grown seedlings (Zentella et al., 2007) and in flowers of Arabidopsis (Hou et al., 2008), as well as genes responding rapidly to GA application (Zentella et al., 2007). Despite the functional similarities between these two DELLA proteins (Gallego-Bartolome et al., 2010), comparison of the sets of genes regulated by GAI and RGA showed little overlap. Out of the 148 GAI targets in etiolated seedlings, 19 genes overlapped with RGA targets in seedlings (Zentella et al., 2007) and 11 in flowers (Hou et al., 2008), which corresponds to 13% and 7% of the GAI-regulated genes respectively (Figures 2.2B and C). Only five genes overlapped in all conditions (Figure 2.2B) and, remarkably, four of them encode members of the GA pathway (*GA20ox1*, *GA20ox2*, *GA3ox1*, and *GID1b*) supporting the notion that the strong regulation of GA activity by DELLA proteins extends to several tissues and growth conditions. However, beyond this regulatory process, a limited overlap in targets displayed by DELLA proteins is evident. It is unlikely that this effect is the consequence of the different expression patterns of the *DELLA* genes used in these studies, given that ubiquitous promoters were used to drive their expression (Zentella et al., 2007; Alabadí et al., 2008). Rather, the low degree of overlap probably reflects the presence of very different sets of transcription factors available for DELLA interaction in etiolated seedlings (our current study) compared with light-grown seedlings and flowers.

### **2.2.3. GAI regulates target genes in part through PIFs and HY5 transcription factors**

The proper control of the developmental switch between skotomorphogenesis and photomorphogenesis after germination is triggered by light through the activation






**Figure 2.2 Transcriptomic analysis of early targets of DELLA proteins.**

(A) Heatmap representation of the 148 best scored genes (q-value ≤ 8). (B) Illustration of the overlap with the datasets of DELLA target genes in two other developmental situations (Zentella et al., 2007; Hou et al., 2009) (C) Heatmap representation of the differential expression of genes overlapping in the three datasets. Red and blue colors in the heatmaps represent induced and repressed genes, respectively.

of transcription factors that promote photomorphogenesis, like ELONGATED HYPOCOTYL5 (HY5), and the inactivation of other transcription factors that promote etiolated growth, such as the PHYTOCHROME-INTERACTING FACTORS, (PIFs) (Alabadí and Blázquez, 2009). Remarkably, GAs counterbalance the effect of light by regulating negatively HY5 protein levels (Alabadi et al 2008), and also through a physical interaction between DELLAs and PIFs that prevents the binding of these transcription factors to their target promoters (de Lucas et al., 2008; Feng et al., 2008). To investigate at the molecular level the extent of these functional interactions, we compared the list of GAI targets with the available lists of genes regulated by HY5 and the PIFs. We reasoned that this comparison would allow us to identify which GAI-

regulated genes depend on the activity of these transcription factors, and delineate the transcriptional network that mediates the GA-control on this developmental switch. While a faithful dataset of *in vivo* target genes for HY5 in light-grown seedlings has been generated by ChIP-to-chip experiments (Lee et al., 2007), the only available list of putative PIF targets can be extracted from transcriptomic analyses of dark- and light-grown wild-type and *pifQ* mutants (Leivar et al., 2009). As shown in Figure 2.3, almost half of the GAI regulated targets are either regulated by HY5, the PIFs, or both, supporting the relevance of these transcription factors in transcriptional regulation by DELLAs.



**Figure 2.3** Meta-analysis comparing microarray data from *HS::gai-1*, HY5 and PIF targets.

Venn diagram of microarray data from *HS::gai-1*, HY5 targets and quadruple *pif* mutant (*pifQ*) show common genes regulated by GAI, HY5 and PIF proteins. Heatmaps show the behaviour of common GAI-HY5, GAI-HY5-PIF and GAI-PIF targets in different light conditions.

**Wt R/D**, data comes from differential expressed genes under red light compared to dark in a WT. **pifQ/wt D**, data comes from differential expressed genes among quadruple *pif* mutant compared to wt in darkness. **pifQ/wt R**, data comes from differential expressed genes among quadruple *pif* mutant compared to wt under red light.


The comparisons are consistent with current models of light and GA regulation. For instance, many of the genes whose promoters are bound by HY5 are coherently regulated by light treatments, and also by DELLA accumulation (Figure 2.3). Only a few of them displayed conflictive regulation by light and by DELLAs (induced by light, bound by HY5, repressed by DELLAs), probably indicating that these targets common to HY5 and DELLAs are not regulated jointly, but in parallel. In the case of PIFs, it is well established that DELLAs have a negative effect on PIFs activity (de Lucas et al., 2008; Feng et al., 2008). In agreement with this, many genes that are targets for both PIFs and DELLAs show the same behavior for DELLA accumulation and for PIF deficiency (Figure 2.3). An indication that this regulation is biologically relevant is that endodermis-specific expression of *PIF1* in *pifQ* mutants restores the formation of the apical hook (Kim et al., 2011), and this tissue specificity is also observed for the regulation of the apical hook by GAs (Gallego-Bartolome et al., 2011). But there are also some cases where the opposite behavior is observed, suggesting either that DELLA regulation of those targets does not proceed through PIFs, or that not all individual PIFs have equivalent activities and abilities to interact with DELLAs *in vivo*.

#### **2.2.4. Promoter analysis of GAI regulated targets suggests new transcription factors mediating DELLA activity.**

Although half of the GAI targets are likely regulated by HY5 and PIFs, there is no obvious connection between these two transcription factors and the rest of the genes regulated by GAI. To get hints regarding the identity of the additional transcription factors mediating DELLA regulation, we investigated the enrichment of particular *cis* elements among the promoters of genes up- and downregulated in *HS::gai-1* using ELEMENT (<http://element.cgrb.oregonstate.edu/>) (Nemhauser et al., 2004). This tool returns those 3-8 bp sequences that are over-represented in the 1000 bp upstream region that precedes the transcription start site of target genes, compared to those same regions through the whole Arabidopsis genome. According to this analysis, apart from a small number of putative *cis* elements with unknown identity (Figure 2.4A), the promoters of genes induced by GAI are enriched in the Dof (AAAG) (Yanagisawa, 2004) and ARR1 (NGATT) (Sakai et al., 2000) binding sites. Interestingly, both types of transcription factors have been related to GAs. For example, Dof proteins have been implicated in the regulation of GA signaling and biosynthesis in Arabidopsis and barley, possibly in the DELLA-mediated feedback regulation of the GA pathway (Gabriele et al., 2010; (Mena

et al., 2002; Zou et al., 2008). And ARR1 has been shown to mediate the control of root meristem size in response to GAs through the up-regulation of *ARR1* expression by DELLA proteins (Moubayidin et al., 2010).

A



B




Figure 2.4 Over-represented *cis* elements among DELLA-regulated promoters.

(A) Logos of over-represented *cis* elements in the promoters of induced and repressed targets in the *HS::gai-1* microarray experiment. (B) Logos of an over-represented *cis* elements in the promoters of induced and repressed genes coming from the joint dataset of *HS::gai-1*, *rga-Δ17* (Zentella et al, 2007) and GA/floral (Hou et al (2009) microarray targets. The logo representation was obtained at <http://weblogo.berkeley.edu/> (Crooks et al., 2004).

To investigate if this analysis allows the identification of DELLA-related regulatory sequences common to different developmental contexts, we examined the enrichment of *cis* elements in the dataset containing all DELLA target promoters found in all available experiments (Zentella et al., 2007; Hou et al., 2008). Surprisingly, the analysis of the promoters GAI induced targets showed an enrichment in two known regulatory sequences: the G-box (CACGTG) (Giuliano et al., 1988; Riechmann et al., 1996) and a sequence similar to the CArG box (CC(A/T)<sub>6</sub>GG) (Riechmann et al., 1996), which also includes a Dof binding site (AAAG) (Figure 4B). The presence of G-boxes is reasonable, taking into account that they are bound both by bHLH and bZIP transcription factors (Foster et al., 1994; Lee et al., 2007), like the PIFs and HY5, for which strong molecular interactions exist with respect to GA signaling (de Lucas et al., 2008; Feng et al., 2008; Alabadi et al. 2008). However, no link between MADS-box transcription factors and GAs has been established yet.

On the other hand, the E-box CATGTG appeared as an over-represented sequence both in the etiolated and in the joint dataset of DELLA repressed targets (Figure 2.4). E-boxes (CAnnTG) are usually bound by bHLH proteins. Unlike the G-box, which is a particular case of an E-box bound by PIFs (Giuliano et al., 1988; Huq and Quail, 2002; de Lucas et al., 2008; Feng et al., 2008), the CATGTG (or CACATG, in the opposite orientation) is bound by the brassinosteroid signaling elements BZR1 and BES1 (Moubayidin et al., 2010; Sun et al., 2010; Yu et al., 2011). Moreover, this element is enriched in promoters of dawn-phased genes that oscillate under short-day photocycles, and it is important for gating their expression by the circadian clock (Michael et al., 2008). Thus, the enrichment of this E-box element could subtend the connection between DELLA proteins and circadian regulation of transcription (Arana et al., 2011) or point to new interactions between the GA and brassinosteroid pathways.

### **2.2.5. Gene ontology analysis of GAI-regulated genes**

Next, to identify the basic biological processes that are regulated by GAs in etiolated seedlings at the molecular level, we tried to extract any significantly over-represented Gene Ontology term (GO) (Ashburner et al., 2000) from our gene list by using the FatiGO algorithm (Al-Shahrour et al., 2005). This analysis suggests that GAI is closely involved in the control of GA homeostasis, growth, other hormone pathways, light, and stress responses and in controlling transcriptional networks (Table 2.1; see below).

Table 2.1 GO term of non redundant categories of *HS::gai-1* targets

BIOLOGICAL PROCESS				MOLECULAR FUNCTION			
name	p-value	genes in this group		GO	p-value	genes in this group	
Response to gibberellin stimulus	2,4E-09	AT2G01570	RGA1	oxidoreductase activity	6E-05	AT4G25420	GA20OX1
Gibberellic acid mediated signaling	5,1E-08	AT3G05120	GID1A			AT1G60980	ATGA20OX4
Gibberellin biosynthetic process	1,2E-06	AT2G37640	EXP3			AT4G21200	GA2OX8
		AT1G67100	LBD40			AT1G15550	GA3OX1
		AT1G66350	RGL1			AT5G51810	GA20OX2
		AT4G25420	GA20OX1	transcription factor activity	2E-05	AT5G56860	GNC
		AT5G25900	GA3			AT3G60390	HAT3
		AT3G63010	GID1B			AT1G49560	MYB TF
		AT2G04240	XERICO			AT4G00050	UNE10
		AT1G15550	GA3OX1			AT5G28300	trihelix DNA-bin
		AT5G51810	GA20OX2			AT1G56650	PAP1
		AT5G67480	BT4			AT3G50890	AtHB28
						AT3G18010	WOX1
						AT4G32280	IAA29
						AT1G53910	RAP2.12
Regulation of transcription	0,00485	AT3G28857	PRE5			AT2G02450	ANAC035
		AT4G39070	STH7			AT2G42380	AtBZIP34
		AT1G66380	MYB114			AT1G66380	MYB114
		AT3G60390	HAT3			AT4G39070	STH7
		AT4G30180	bHLH146			AT1G69690	TCP TF
		AT1G49560	MYB TF			AT3G06590	AIF2
		AT4G00050	UNE10			AT5G39860	PRE1
		AT5G28300	trihelix DNA-bind			AT1G21910	AtERF012
		AT1G53910	RAP2.12			AT2G01570	RGA1
		AT5G14750	ATMYB66			AT4G30180	bHLH146
		AT1G14600	Myb-like TF			AT1G66350	RGL1
		AT1G69690	TCP TF			AT3G15540	IAA19
		AT1G56650	PAP1			AT3G28730	ATHMG
		AT3G06590	AIF2			AT5G14750	ATMYB66
		AT5G15150	ATHB-3			AT1G14600	Myb-like TF
		AT2G01570	RGA1			AT5G41920	SCL25
		AT5G41920	SCL25			AT5G15150	ATHB-3
		AT4G32890	GATA9			AT4G32890	GATA9
response to red or far red light	0,00085	AT2G01570	RGA1	monooxygenase activity	0,0025	AT5G25900	GA3
		AT5G04190	PKS4			AT4G28720	YUCCA8
		AT2G37640	EXP3			AT2G26710	BAS1
		AT4G32280	IAA29			AT1G58440	XF1
		AT4G25260	invertase inhibitor			AT5G38970	BR6OX1
		AT1G15550	GA3OX1				
		AT5G51810	GA20OX2				
response to jasmonic acid stimulus	0,0193	AT1G66350	RGL1	lyase activity	0,0244	AT3G51430	YLS2
		AT2G01570	RGA1			AT3G07010	pectate lyase
		AT1G66380	MYB114			AT1G27980	DPL1
		AT5G13220	JAZ10			AT1G67750	pectate lyase
		AT1G56650	PAP1			AT5G28020	CYSD2
response to salt stress	0,0366	AT1G13930	response to stress			AT4G37770	ACS8
		AT2G01570	RGA1			AT5G36160	putative C-S lya
		AT1G66350	RGL1				
		AT1G56650	PAP1				
		AT2G33380	RD20				
		AT2G04240	XERICO				
unidimensional cell growth	0,0115	AT5G51810	GA20OX2				
		AT4G25420	GA20OX1				
		AT2G37640	EXP3				
		AT2G20750	ATEXPB1				
		AT2G40610	ATEXPA8				

## 2.2.6. Direct regulation of the GA pathway by DELLA proteins

The control of the homeostasis of GA levels in the plant is finely achieved through feedback and feedforward mechanisms that require the activity of the different elements of the GA signaling pathways (Hedden and Phillips, 2000; Yamaguchi, 2008).

Besides, this mechanism also operates to control the level of GA receptors' transcripts (Griffiths et al., 2006). Recently Zentella et al. (2007) demonstrated the involvement of the DELLA protein RGA in this mechanism, as they showed RGA directly up-regulates the expression of *AtGA20ox2*, *AtGA3ox1*, *GA INSENSITIVE DWARF1a* (*GID1a*), and *GID1b* genes. In addition to these genes, we found *AtGA20ox1*, and *AtGA20ox4* among the GAI up-regulated genes, and *AtGA20ox8*, *GAI*, and *RGL1* among the GAI down-regulated genes (Figure 2.5). The regulation of these genes by GAI was confirmed by analyzing their transcript levels in several GA-related mutants and transgenic lines (Figure 2.5A). Control on the expression of the majority of genes was shared with RGA and with other DELLA proteins –see for example regulation of *AtGA20ox8* gene expression that was repressed by PAC treatment to similar levels in the wild type and in the double null mutant *gai-t6 rga-24*.


The rapid change in the expression of these genes in response to *gai-1* accumulation suggested to us that they might be direct targets. Thus, we tested this possibility by using transgenic lines that express a translational fusion between *gai-1* and the glucocorticoid receptor domain from rats, under the control of the *GAI* promoter (Gallego-Bartolome et al., 2011). Dexamethasone treatment mimicked the effect on target gene expression that a heat-shock treatment provokes in the *HS::gai-1* line, and this effect was not abolished by cycloheximide indicating these genes are directly regulated by GAI, i.e. independently of protein synthesis (Figure 2.5B). These results broaden our current understanding of DELLA control on the homeostasis of the GA pathway and indicate that in etiolated seedlings this control includes: 1) genes coding for GA 2-oxidases, which indicates that DELLA proteins directly participate in the feedforward mechanism. And 2) genes coding for DELLA proteins, indicating that DELLA activity negatively regulates the expression of their own genes.

#### **2.2.7. DELLA proteins mediate direct cross-regulation with auxin and ethylene pathways**


Our analysis indicates that the crosstalk between GAs and other hormones could be exerted at the transcriptional level. Among the relevant targets for GAI, we identified several genes related to auxin synthesis and signaling, such as the negative auxin signaling intermediates *AUXIN/INDOLE-3-ACETIC ACID19* (*Aux/IAA19*) (Tatematsu, Kumagai et al. 2004) and *Aux/IAA29*, two auxin-inducible *SMALL AUXIN UPREGULATED* genes, and also *INDOLE-3-ACETIC ACID*

*METHYLTRANSFERASE1* (*IAMT1*) (Qin, Gu et al. 2005) and *YUCCA3* (*YUC3*) involved in IAA inactivation (Li, Hou et al. 2007) and biosynthesis (Zhao, Christensen et al. 2001), respectively (Table 2.1 and Supplemental Table 1).

A



B



**Figure 2.5. DELLA transcriptional regulation of GA metabolism and signaling genes.**

A) Validation of *gai-1* target genes in independent experiment using PAC-grown seedlings and GA mutants. B) Direct regulation by *gai-1* of target genes. Experiments with *pGAI::gai-1-GR* transgenic line demonstrated that “*de novo*” protein synthesis is not necessary for *gai-1* modulation of target expression.




The ethylene biosynthesis genes *ACC SYNTHASE8 (ACS8)* and *ACS5/ETO2* (Vogel, Woeste et al. 1998; Yamagami, Tsuchisaka et al. 2003) were also among the genes downregulated by GAI, extending the control by GAs to hormones other than auxin. Transcriptional control of these genes by GAI was direct, and was exerted redundantly with other DELLAs (Figure 2.6A and B) (Gallego-Bartolome, Arana et al. 2011; Gallego-Bartolome, Kami et al. 2011).


#### **2.2.8. DELLAs impinge on transcriptional networks**

Several GO terms referring to transcription factors were also overrepresented among the GAI-regulated genes (Table 1 and Supplemental Table 1). The regulation by GAs of several genes of diverse families, including homeobox-leucine zipper (*HOMEBOX-LEUCINE ZIPPER PROTEIN3* and *7*; *HAT3* and *HAT7*) (Ciarbelli et al., 2008), MYB (*PRODUCTION OF ANTHOCYANIN PIGMENT1*, *PAP1*) (Borevitz et al., 2000), TCP (*TCP15*), and HLH (*PACLOBUTRAZOL RESISTANT1* and *5*; *PRE1* and *PRE5*) (Lee et al., 2006), was confirmed in various GA-related mutants and transgenic lines (Figure 2.6A). Importantly, the expression of these genes was directly regulated by GAI (Figure 2.6B), which indicates that GA activity directly impinges cellular transcriptional networks to amplify its signal and thus to control the skotomorphogenic development. Remarkably, our results provide a molecular explanation for the observation that *PRE1* expression, and likely that of *PRE5*, are induced by GAs and act as positive elements promoting several GA responses by acting downstream of GAI (Lee et al., 2006), and suggests that these HLH proteins also exert a role promoting skotomorphogenesis. Furthermore, this regulation might amplify the effect of the GA pathway on transcriptional networks since both proteins might act as dominant negative regulators of others bHLH by forming non DNA-binding heterodimers, as demonstrated, for instance, with the HLH protein HFR1 in the control of the shade avoidance response (Hornitschek et al., 2009).

A



B



**Figure 2.6. DELLA transcriptional regulation of transcription factor and hormone-related genes.**

A) Validation of *gai-1* target genes in independent experiment using PAC-grown seedlings and GA mutants. B) Direct regulation by *gai-1* of target genes. Experiments with *pGAI::gai-1-GR* transgenic line demonstrated that “*de novo*” protein synthesis is not necessary for *gai-1* modulation of target expression.

### 2.3. Conclusions

The enormous plasticity in plant development depends on highly wired, interconnected signaling networks that properly integrate endogenous and environmental cues (Casal et al., 2004). In many cases, the cross-regulation between pathways occurs at the level of transcriptional regulation (Kuppusamy et al., 2008). The

output from the GA pathway largely relies on the activity of the transcriptional regulators DELLA proteins. Our transcriptomic analysis of DELLA responsive genes in etiolated seedlings reveals that the activity of the GA pathway directly influences on other hormone pathways –ethylene and auxin- and on pre-existing transcriptional networks. Further, the *in silico* promoter analysis of DELLA-regulated genes suggests that transcription factors of the Dof and type-B ARRs classes, as well as the brassinosteroid signaling elements, E-box binding proteins BES1 and BZR1, could mediate the activity of GAs. Thus, the GA pathway contributes, at least at the transcriptional level, to the wiring of the signaling network that underlies plant plasticity.

## **2.4. Materials and methods**

### **Plant material and growth conditions**

*Arabidopsis thaliana* GA signaling dominant mutant *rga-Δ17* (Dill et al., 2001), the double loss-of-function *rga-24 gai-t6* (King et al., 2001) and *pGAI::gai-1-GR* (Gallego-Bartolomé, Kami et al, 2011) are in the *Ler* background, while *HS::gai-1* and the *35S::gai-1 (gai-1ox)* (Alabadí et al., 2008) are derived from Col-0 accession. Seeds were sterilized and stratified for 6 days in water at 4°C. Germination took place under white fluorescent light (90–100  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ) at 22°C for 6 h in a Percival growth chamber E-30B (<http://www.percival-scientific.com>). Seeds were plated in plates of half-strength MS medium with 0.8% (w/v) agar and 1% (w/v) sucrose supplemented with either 1  $\mu\text{M}$  PAC or mock treatment and grown in darkness at 22°C for 3 days. For short-term treatments, seedlings were incubated in the dark in water supplemented with 10  $\mu\text{M}$  CHX and/or 10  $\mu\text{M}$  DEX. MS and PAC were from Duchefa (<http://www.duchefa.com>). DEX and CHX were from Sigma (<http://www.sigmaaldrich.com>).

### **Real-time quantitative RT-PCR**

RNA extraction, cDNA synthesis, quantitative RT-PCR (qRT-PCR), analysis, and primer sequences for amplification of *AtGA20ox2* and *EF1- $\alpha$*  genes, used to normalize all expression data, have been described (Frigerio et al., 2006). qRT-PCR oligonucleotides sequences for the other target genes are listed in the following table.

Gene	AGI number	forward primer (5' to 3')	reverse primer (5' to 3')
20ox1	At4g25420	CTTCCATCAACGTTCTCGAGC	GGTTTTGAAGGTCGATGAGAGG
20ox2	At5g51810	AGAAACCTTCCATTGACATTCCA	AGAGATCGATGAACGGGACG
20ox4	At1g60980	CTATCCAAAATGCAAGCAACCA	CAGTGAGGCCCCGTACCTAGT
2ox8	At4g21200	CATGGAGCAATGGCATGTACA	GGTTCGTCATCACACGGTGTT
3ox1	At1g15550	GATCTCCTCTTCTCCGCTGCT	GAGGGATGTTTTACCCGGTG
RGL1	At1g66350	TCAGTGGCGGTAACTCGGT	GGGATGAGCTAAGAGGCGATG
GID1a	At3g05120	GTGACGGTTAGAGACCGCA	TCCCTCGGGTAAAAACGCTT
GID1b	At3g63010	TCGCCCTGACGGTTCTTTC	TTACGGTCAAGGAACTCGGC
GAI 3'UTR	At1g14920	AATGAATTGATCTGTTGAACCGG	GGCTTCGGTCGGAAATCTATC
PRE1	At5g39860	CAAAATTTTGCCGGAGATTGG	GGCTGATGCCTTATCAGAACG
PRE5	At3g28857	CTCCGATGACCAGATGATCGA	GGCAAAAAGTACGAGGACTT
HAT7	At5g15150	TCCCTTCTTCGATCCGATCC	CAATATGGGTAGAGGTCGTGGTC
PAP1	At1g56650	TTGGTTCCTGAAGCGACGAC	GTCAAAAAGCCAAGGTGTCCC
IAMT1	At5g55250	CTGCCTTCTCCTTGCAATTGG	TATCCGTCACACTTTCGGGC
YUC3	At1g04610	AAGATCAAAATCGTCCCCGG	CTCAACTTTGCCTTTGCCGA

To analyze expression of transgenic *gai-1* in the *pHsp::gai-1* seedlings, we used an oligonucleotide annealing to the 5' UTR of the HSP18.2 gene, which is included in the construct, as the forward primer (5' CCCGAAAAGCAACGAACAAT-3'), and an oligonucleotide annealing to the *gai-1* coding region as the reverse primer (5'-TCATTCATCATCATAGTCTTCTTATCTTGA-3').

### Gene expression analysis by long oligonucleotide microarrays

Seeds of *Arabidopsis* Col-0 and *HS::gai-1* transgenic line were sterilized, sown, stratified, and germinated as described above. Seedlings were grown for 3 days in darkness at 22°C. Then both wild type and transgenic seedlings were moved to 37°C for 30 minutes. After the heat-shock treatment plates were moved back to 22°C. Samples were collected at time points 0, 1, 2, and 4 hours after the beginning of the heat treatment. Three independent biological replicates were used for the analysis. Total RNA from whole seedlings was extracted as described above. RNA amplification, labeling, and hybridization of microarray slides were carried out as described (Bueso et al., 2007). Scanning of the slides, quantification of spots, and normalization were performed as previously described (Stavang et al., 2009).

## Promoter analysis

Promoter analysis (<http://element.cgrb.oregonstate.edu/>) was done as previously described (14) using the ELEMENT webtool (<http://element.cgrb.oregonstate.edu/>). Logos were built using the Weblogo webtool (<http://weblogo.berkeley.edu/>). The cluster lists are formulated by using the highest-count promoter core elements. All longer elements containing the core element are clustered together. PLACE database (<http://www.dna.affrc.go.jp/PLACE/>) was used to identify any known CIS-acting element.

## 2.5. Bibliography

- Al-Shahrour F, Diaz-Uriarte R, Dopazo J** (2005) Discovering molecular functions significantly related to phenotypes by combining gene expression data and biological information. *Bioinformatics* **21**: 2988-2993
- Alabadí D, Blázquez MA** (2009) Molecular interactions between light and hormone signaling to control plant growth. *Plant Mol Biol* **69**: 409-417
- Alabadi D, Blázquez MA, Carbonell J, Ferrandiz C, Perez-Amador MA** (2009) Instructive roles for hormones in plant development. *Int J Dev Biol* **53**: 1597-1608
- Alabadí D, Gallego-Bartolomé J, García-Cárcel L, Orlando L, Rubio V, Martínez C, Frigerio M, Iglesias-Pedraz JM, Espinosa A, Deng XW, Blázquez MA** (2008) Gibberellins modulate light signaling pathways to prevent Arabidopsis seedling de-etiolation in darkness. *Plant J* **53**: 324-335
- Alabadi D, Gallego-Bartolomé J, Orlando L, Garcia-Carcel L, Rubio V, Martinez C, Frigerio M, Iglesias-Pedraz JM, Espinosa A, Deng XW, Blázquez MA** (2008) Gibberellins modulate light signaling pathways to prevent Arabidopsis seedling de-etiolation in darkness. *Plant J* **53**: 324-335
- Alabadí D, Gil J, Blázquez MA, García-Martínez JL** (2004) Gibberellins repress photomorphogenesis in darkness. *Plant Physiol* **134**: 1050-1057
- Arana MV, Marin-de la Rosa N, Maloof JN, Blázquez MA, Alabadi D** (2011) Circadian oscillation of gibberellin signaling in Arabidopsis. *Proc Natl Acad Sci U S A*
- Arnaud N, Girin T, Sorefan K, Fuentes S, Wood TA, Lawrenson T, Sablowski R, Ostergaard L** (2010) Gibberellins control fruit patterning in Arabidopsis thaliana. *Genes Dev* **24**: 2127-2132
- Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, Harris MA, Hill DP, Issel-Tarver L, Kasarskis A, Lewis S, Matese JC, Richardson JE, Ringwald M, Rubin GM, Sherlock G** (2000) Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet* **25**: 25-29
- Borevitz JO, Xia Y, Blount J, Dixon RA, Lamb C** (2000) Activation tagging identifies a conserved MYB regulator of phenylpropanoid biosynthesis. *Plant Cell* **12**: 2383-2394
- Bueso E, Alejandro S, Carbonell P, Perez-Amador MA, Fayos J, Belles JM, Rodriguez PL, Serrano R** (2007) The lithium tolerance of the Arabidopsis cat2 mutant reveals a cross-talk between oxidative stress and ethylene. *Plant J* **52**: 1052-1065
- Cao D, Cheng H, Wu W, Soo HM, Peng J** (2006) Gibberellin mobilizes distinct DELLA-dependent transcriptomes to regulate seed germination and floral development in Arabidopsis. *Plant Physiol* **142**: 509-525
- Casal JJ, Fankhauser C, Coupland G, Blázquez MA** (2004) Signalling for developmental plasticity. *Trends Plant Sci* **9**: 309-314


- Ciarbelli AR, Ciolfi A, Salvucci S, Ruzza V, Possenti M, Carabelli M, Fruscalzo A, Sessa G, Morelli G, Ruberti I** (2008) The Arabidopsis homeodomain-leucine zipper II gene family: diversity and redundancy. *Plant Mol Biol* **68**: 465-478
- Crooks GE, Hon G, Chandonia JM, Brenner SE** (2004) WebLogo: a sequence logo generator. *Genome Res* **14**: 1188-1190
- de Lucas M, Daviere JM, Rodriguez-Falcon M, Pontin M, Iglesias-Pedraz JM, Lorrain S, Fankhauser C, Blazquez MA, Titarenko E, Prat S** (2008) A molecular framework for light and gibberellin control of cell elongation. *Nature* **451**: 480-484
- de Lucas M, Davière JM, Rodríguez-Falcón M, Pontin M, Iglesias-Pedraz JM, Lorrain S, Fankhauser C, Blázquez MA, Titarenko E, Prat S** (2008) A molecular framework for light and gibberellin control of cell elongation. *Nature* **451**: 480-484
- Dill A, Jung HS, Sun TP** (2001) The DELLA motif is essential for gibberellin-induced degradation of RGA. *Proc Natl Acad Sci U S A* **98**: 14162-14167
- Feng S, Martinez C, Gusmaroli G, Wang Y, Zhou J, Wang F, Chen L, Yu L, Iglesias-Pedraz JM, Kircher S, Schafer E, Fu X, Fan LM, Deng XW** (2008) Coordinated regulation of Arabidopsis thaliana development by light and gibberellins. *Nature* **451**: 475-479
- Foster R, Izawa T, Chua NH** (1994) Plant bZIP proteins gather at ACGT elements. *FASEB J* **8**: 192-200
- Frigerio M, Alabadi D, Pérez-Gómez J, García-Cárcel L, Phillips AL, Hedden P, Blázquez MA** (2006) Transcriptional regulation of gibberellin metabolism genes by auxin signaling in Arabidopsis. *Plant Physiol* **142**: 553-563
- Fu X, Richards DE, Ait-Ali T, Hynes LW, Ougham H, Peng J, Harberd NP** (2002) Gibberellin-mediated proteasome-dependent degradation of the barley DELLA protein SLN1 repressor. *Plant Cell* **14**: 3191-3200
- Gallego-Bartolome J, Arana MV, Vandenbussche F, Zadnikova P, Minguet EG, Guardiola V, Van Der Straeten D, Benkova E, Alabadi D, Blazquez MA** (2011) Hierarchy of hormone action controlling apical hook development in Arabidopsis. *Plant J*
- Gallego-Bartolome J, Kami C, Fankhauser C, Alabadi D, Blazquez MA** (2011) A hormonal regulatory module that provides flexibility to tropic responses. *Plant Physiol*
- Gallego-Bartolome J, Minguet EG, Marin JA, Prat S, Blazquez MA, Alabadi D** (2010) Transcriptional diversification and functional conservation between DELLA proteins in Arabidopsis. *Mol Biol Evol* **27**: 1247-1256
- Giuliano G, Pichersky E, Malik VS, Timko MP, Scolnik PA, Cashmore AR** (1988) An evolutionarily conserved protein binding sequence upstream of a plant light-regulated gene. *Proc Natl Acad Sci U S A* **85**: 7089-7093
- Griffiths J, Murase K, Rieu I, Zentella R, Zhang ZL, Powers SJ, Gong F, Phillips AL, Hedden P, Sun TP, Thomas SG** (2006) Genetic characterization and functional analysis of the GID1 gibberellin receptors in Arabidopsis. *Plant Cell* **18**: 3399-3414
- Harberd NP, Belfield E, Yasumura Y** (2009) The angiosperm gibberellin-GID1-DELLA growth regulatory mechanism: how an "inhibitor of an inhibitor" enables flexible response to fluctuating environments. *Plant Cell* **21**: 1328-1339
- Hedden P, Phillips AL** (2000) Gibberellin metabolism: new insights revealed by the genes. *Trends Plant Sci* **5**: 523-530
- Heo JO, Chang KS, Kim IA, Lee MH, Lee SA, Song SK, Lee MM, Lim J** (2011) Funneling of gibberellin signaling by the GRAS transcription regulator scarecrow-like 3 in the Arabidopsis root. *Proc Natl Acad Sci U S A* **108**: 2166-2171
- Hirano K, Asano K, Tsuji H, Kawamura M, Mori H, Kitano H, Ueguchi-Tanaka M, Matsuoka M** (2010) Characterization of the Molecular Mechanism Underlying Gibberellin Perception Complex Formation in Rice. *Plant Cell*
- Hornitschek P, Lorrain S, Zoete V, Michielin O, Fankhauser C** (2009) Inhibition of the shade avoidance response by formation of non-DNA binding bHLH heterodimers. *Embo J* **28**: 3893-3902

- Hou X, Hu WW, Shen L, Lee LY, Tao Z, Han JH, Yu H** (2008) Global Identification of DELLA Target Genes during Arabidopsis Flower Development. *Plant Physiol*
- Hou X, Hu WW, Shen L, Lee LY, Tao Z, Han JH, Yu H** (2008) Global identification of DELLA target genes during Arabidopsis flower development. *Plant Physiol* **147**: 1126-1142
- Hou X, Lee LY, Xia K, Yan Y, Yu H** (2010) DELLAs modulate jasmonate signaling via competitive binding to JAZs. *Dev Cell* **19**: 884-894
- Huq E, Quail PH** (2002) PIF4, a phytochrome-interacting bHLH factor, functions as a negative regulator of phytochrome B signaling in Arabidopsis. *EMBO J* **21**: 2441-2450
- Itoh H, Matsuoka M, Steber CM** (2003) A role for the ubiquitin-26S-proteasome pathway in gibberellin signaling. *Trends Plant Sci* **8**: 492-497
- Jaillais Y, Chory J** (2010) Unraveling the paradoxes of plant hormone signaling integration. *Nat Struct Mol Biol* **17**: 642-645
- King KE, Moritz T, Harberd NP** (2001) Gibberellins are not required for normal stem growth in Arabidopsis thaliana in the absence of GAI and RGA. *Genetics* **159**: 767-776
- Kuppusamy KT, Walcher CL, Nemhauser JL** (2008) Cross-regulatory mechanisms in hormone signaling. *Plant Mol Biol*
- Lee J, He K, Stolz V, Lee H, Figueroa P, Gao Y, Tongprasit W, Zhao H, Lee I, Deng XW** (2007) Analysis of transcription factor HY5 genomic binding sites revealed its hierarchical role in light regulation of development. *Plant Cell* **19**: 731-749
- Lee S, Yang KY, Kim YM, Park SY, Kim SY, Soh MS** (2006) Overexpression of PRE1 and its homologous genes activates Gibberellin-dependent responses in Arabidopsis thaliana. *Plant Cell Physiol* **47**: 591-600
- Leivar P, Tepperman JM, Monte E, Calderon RH, Liu TL, Quail PH** (2009) Definition of early transcriptional circuitry involved in light-induced reversal of PIF-imposed repression of photomorphogenesis in young Arabidopsis seedlings. *Plant Cell* **21**: 3535-3553
- Li L, Hou X, Tsuge T, Ding M, Aoyama T, Oka A, Gu H, Zhao Y, Qu LJ** (2007) The possible action mechanisms of indole-3-acetic acid methyl ester in Arabidopsis. *Plant Cell Rep*
- Mena M, Cejudo FJ, Isabel-Lamonedá I, Carbonero P** (2002) A role for the DOF transcription factor BPBF in the regulation of gibberellin-responsive genes in barley aleurone. *Plant Physiol* **130**: 111-119
- Michael TP, Breton G, Hazen SP, Priest H, Mockler TC, Kay SA, Chory J** (2008) A morning-specific phytohormone gene expression program underlying rhythmic plant growth. *PLoS Biol* **6**: e225
- Moubayidin L, Perilli S, Dello Ioio R, Di Mambro R, Costantino P, Sabatini S** (2010) The rate of cell differentiation controls the Arabidopsis root meristem growth phase. *Curr Biol* **20**: 1138-1143
- Nemhauser JL, Hong F, Chory J** (2006) Different plant hormones regulate similar processes through largely nonoverlapping transcriptional responses. *Cell* **126**: 467-475
- Nemhauser JL, Mockler TC, Chory J** (2004) Interdependency of brassinosteroid and auxin signaling in Arabidopsis. *PLoS Biol* **2**: E258
- Ogawa M, Hanada A, Yamauchi Y, Kuwahara A, Kamiya Y, Yamaguchi S** (2003) Gibberellin biosynthesis and response during Arabidopsis seed germination. *Plant Cell* **15**: 1591-1604
- Peng J, Carol P, Richards DE, King KE, Cowling RJ, Murphy GP, Harberd NP** (1997) The Arabidopsis GAI gene defines a signaling pathway that negatively regulates gibberellin responses. *Genes Dev* **11**: 3194-3205
- Qin G, Gu H, Zhao Y, Ma Z, Shi G, Yang Y, Pichersky E, Chen H, Liu M, Chen Z, Qu LJ** (2005) An indole-3-acetic acid carboxyl methyltransferase regulates Arabidopsis leaf development. *Plant Cell* **17**: 2693-2704
- Riechmann JL, Krizek BA, Meyerowitz EM** (1996) Dimerization specificity of Arabidopsis MADS domain homeotic proteins APETALA1, APETALA3, PISTILLATA, and AGAMOUS. *Proc Natl Acad Sci U S A* **93**: 4793-4798

- Sakai H, Aoyama T, Oka A** (2000) Arabidopsis ARR1 and ARR2 response regulators operate as transcriptional activators. *Plant J* **24**: 703-711
- Shimada A, Ueguchi-Tanaka M, Nakatsu T, Nakajima M, Naoe Y, Ohmiya H, Kato H, Matsuoka M** (2008) Structural basis for gibberellin recognition by its receptor GID1. *Nature* **456**: 520-523
- Stavang JA, Gallego-Bartolome J, Gomez MD, Yoshida S, Asami T, Olsen JE, Garcia-Martinez JL, Alabadi D, Blazquez MA** (2009) Hormonal regulation of temperature-induced growth in Arabidopsis. *Plant J* **60**: 589-601
- Sun Y, Fan XY, Cao DM, Tang W, He K, Zhu JY, He JX, Bai MY, Zhu S, Oh E, Patil S, Kim TW, Ji H, Wong WH, Rhee SY, Wang ZY** (2010) Integration of brassinosteroid signal transduction with the transcription network for plant growth regulation in Arabidopsis. *Dev Cell* **19**: 765-777
- Tatematsu K, Kumagai S, Muto H, Sato A, Watahiki MK, Harper RM, Liscum E, Yamamoto KT** (2004) MASSUGU2 encodes Aux/IAA19, an auxin-regulated protein that functions together with the transcriptional activator NPH4/ARF7 to regulate differential growth responses of hypocotyl and formation of lateral roots in Arabidopsis thaliana. *Plant Cell* **16**: 379-393
- Tusher VG, Tibshirani R, Chu G** (2001) Significance analysis of microarrays applied to the ionizing radiation response. *Proc Natl Acad Sci U S A* **98**: 5116-5121
- Ueguchi-Tanaka M, Ashikari M, Nakajima M, Itoh H, Katoh E, Kobayashi M, Chow TY, Hsing YI, Kitano H, Yamaguchi I, Matsuoka M** (2005) GIBBERELLIN INSENSITIVE DWARF1 encodes a soluble receptor for gibberellin. *Nature* **437**: 693-698
- Vogel JP, Woeste KE, Theologis A, Kieber JJ** (1998) Recessive and dominant mutations in the ethylene biosynthetic gene ACS5 of Arabidopsis confer cytokinin insensitivity and ethylene overproduction, respectively. *Proc Natl Acad Sci U S A* **95**: 4766-4771
- Yamagami T, Tsuchisaka A, Yamada K, Haddon WF, Harden LA, Theologis A** (2003) Biochemical diversity among the 1-amino-cyclopropane-1-carboxylate synthase isozymes encoded by the Arabidopsis gene family. *J Biol Chem* **278**: 49102-49112
- Yamaguchi S** (2008) Gibberellin metabolism and its regulation. *Annu Rev Plant Biol* **59**: 225-251
- Yanagisawa S** (2004) Dof domain proteins: plant-specific transcription factors associated with diverse phenomena unique to plants. *Plant Cell Physiol* **45**: 386-391
- Yu X, Li L, Zola J, Aluru M, Ye H, Foudree A, Guo H, Anderson S, Aluru S, Liu P, Rodermeier S, Yin Y** (2011) A brassinosteroid transcriptional network revealed by genome-wide identification of BES1 target genes in Arabidopsis thaliana. *Plant J* **65**: 634-646
- Zentella R, Zhang ZL, Park M, Thomas SG, Endo A, Murase K, Fleet CM, Jikumaru Y, Nambara E, Kamiya Y, Sun TP** (2007) Global analysis of della direct targets in early gibberellin signaling in Arabidopsis. *Plant Cell* **19**: 3037-3057
- Zhang ZL, Ogawa M, Fleet CM, Zentella R, Hu J, Heo JO, Lim J, Kamiya Y, Yamaguchi S, Sun TP** (2011) Scarecrow-like 3 promotes gibberellin signaling by antagonizing master growth repressor DELLA in Arabidopsis. *Proc Natl Acad Sci U S A* **108**: 2160-2165
- Zhao Y, Christensen SK, Fankhauser C, Cashman JR, Cohen JD, Weigel D, Chory J** (2001) A role for flavin monooxygenase-like enzymes in auxin biosynthesis. *Science* **291**: 306-309
- Zou X, Neuman D, Shen QJ** (2008) Interactions of two transcriptional repressors and two transcriptional activators in modulating gibberellin signaling in aleurone cells. *Plant Physiol* **148**: 176-186



## Chapter 3



**A hormonal regulatory module that  
provides flexibility to tropic responses**

Gallego-Bartolome J, Kami C, Fankhauser C, Alabadi D, Blazquez MA.

**Plant Physiol** DOI:10.1104/pp.111.173971 (2011)



### **3.1. Abstract**

Plants orient their growth depending on directional stimuli such as light and gravity, in a process known as tropic response. Tropisms result from asymmetrical accumulation of auxin across the responding organ relative to the direction of the stimulus, which causes differential growth rates on both sides of the organ. Here we show that gibberellins (GAs) attenuate the gravitropic reorientation of stimulated hypocotyls of dark-grown *Arabidopsis* seedlings. We show that the modulation occurs through induction of the expression of the negative regulator of auxin signaling *IAA19/MSG2*. The biological significance of this regulatory mechanism involving GAs and auxin seems to be the maintenance of a high degree of flexibility in tropic responses. This notion is further supported by observations that GA-deficient seedlings showed a much lower variance in the response to gravity compared to wild-type seedlings and that the attenuation of gravitropism by GAs resulted in an increased phototropic response. This suggests that the interplay between auxin and GAs may be particularly important for plant orientation under competing tropic stimuli.

### **3.2. Introduction**

One hundred and thirty years ago, Darwin described that plants can sense their environment and orient themselves for optimal growth and development (Darwin, 1880). Among the signals that promote a tropic response in plants, gravity is unique in that it is constant and unidirectional. Besides, it generally induces the underground tissues to bend towards the signal, and the aerial parts against the stimulating vector. Like in other tropisms, when plants perceive a change in their position relative to the gravity vector, they respond by differential growth on either side of the affected organ (Esmon et al., 2005), and several hormones have been involved in the control of these responses. Among them, auxin is instrumental because it forms a lateral gradient in response to the stimulus and thus establishes the framework for differential growth (Rashotte et al., 2000; Esmon et al., 2006). The differential response to auxin on either side of an organ has been shown to depend on the correct functioning of polar auxin transport and activity of auxin efflux carriers (Friml et al., 2002), and also on the activity of specific Aux/IAA and AUXIN RESPONSE FACTOR (ARF) transcriptional regulators (Harper et al., 2000; Tatematsu et al., 2004). Moreover, brassinosteroids have been proposed to enhance tropic reorientation by facilitating polar auxin transport (Meudt, 1987; Li et al., 2005; Kim et al., 2007).

Gibberellins (GAs) are also known to promote cell expansion (Cowling and Harberd, 1999). The molecular mechanism of GA signaling proceeds through GA-induced degradation of repressor proteins of the DELLA family by the proteasome, thereby activating transcription of growth-promoting genes (Schwechheimer, 2008). Given that GAs regulate growth, sometimes as a subsidiary signal of auxin action (Frigerio et al., 2006), the obvious hypothesis is that GAs would mediate the promotion of differential growth during gravitropic reorientation. However, here we show molecular evidence for a different role of GAs on gravitropism through the attenuation of auxin responsiveness, that results in an increased ability to modulate growth under competing tropic signals.

### **3.3. Results**


#### **3.3.1. Gibberellin deficiency enhances gravitropic reorientation**

To test if GAs are necessary for the promotion of the differential cell expansion that underlies a tropic response, we examined the response of Arabidopsis etiolated hypocotyls to a gravitropic stimulus, under GA-limiting conditions. Surprisingly, paclobutrazol (PAC)-induced deficiency in GA biosynthesis not only did not impair gravitropic reorientation but, on the contrary, the hypocotyls of GA-deficient seedlings displayed an enhanced response to the gravitropic stimulus and a faster reorientation (Fig. 3.1A). This effect was observed at a low PAC concentration that did not inhibit seed germination, and was fully reverted by GA<sub>3</sub> application, demonstrating the specificity of the inhibitor. Moreover, an enhanced rate of response was also evident in the *gai-1D* and *rga-Δ17* mutants, which express dominant versions of the DELLA proteins GIBBERELLIN INSENSITIVE (GAI) and REPRESSOR OF GA1 (RGA) respectively, that constitutively block GA-induced growth (Peng et al., 1997; Dill et al., 2001), (Fig. 3.1B, 3.1C). Since GA-deficiency also causes dwarfism, it is possible that the enhanced gravitropic response were due to an intrinsic capacity of smaller seedlings to display differential growth and bending. However, this is not the case, because transient induction of the dominant allele *gai-1D* increased the response to a gravitropic stimulus without affecting the size of the seedlings (Alabadí et al., 2008) (Fig. 3.1D). Therefore, we conclude that GAs attenuate the gravitropic response in aerial tissues, and that this regulation is likely a direct consequence of DELLA activity.

#### **3.3.2. Expression of *IAA19/MSG2* is repressed by DELLA proteins**

DELLA proteins regulate gene expression in response to GAs (Zentella et al., 2007; de Lucas et al., 2008; Feng et al., 2008). Thus, to elucidate the molecular mechanism that underlies the regulation of gravitropism by GAs, we investigated by microarray analysis the transcriptional changes associated with transient expression of *gai-1D* in two-day-old dark-grown seedlings, i.e. under conditions where it promotes gravitropism (Fig. 3.1D).

To achieve transient expression of *gai-1D*, we used a transgenic line harbouring the *gai-1D* gene under the control of a heat-shock inducible promoter (Alabadí et al., 2008). Incubation of 3-day-old etiolated seedlings at 37°C for 30 min led to altered




**Fig. 3.1** Gravitropic reorientation of hypocotyls of GA deficient seedlings.

(A), (B), (C) Seedlings derived from *Ler* were grown for 3 days in darkness, and the plates were turned 90° at time 0. (D) *Col* and *HS::gai-1D* seedlings were grown for 3 days in darkness and subject to a 30-min heat shock at 37°C before turning the plates 90°. Reorientation of the hypocotyls was monitored in darkness as explained in Materials and Methods. When used, paclobutrazol (PAC) concentration was 0.4 μM. Error bars represent SD (n>16 individual seedlings). The effect of PAC, *gai-1D* and *rgaΔ17* and *HS::gai-1D* with respect to the wild type was statistically significant (p<0.05) as indicated by a two-way ANOVA test.

expression of around 150 genes 4 hours after the inductive stimulus when compared with wild-type seedlings subject to the same treatment (Chapter II).

Remarkably, expression of *IAA19/MASSUGU2* (*IAA19/MSG2*), which encodes a member of the Aux/IAA family of proteins that negatively regulate auxin signaling (Tatematsu et al., 2004; Overvoorde et al., 2005), was steadily downregulated (Fig. 3.2A). The two closest paralogs of *IAA19/MSG2*, *IAA5* and *IAA6/SHORT HYPOCOTYL1* (*IAA6/SHY1*), were also repressed following *gai-1D* induction (Fig. 3.2A), and the expression of all these genes was consistently lower in dark-grown seedlings of the *gai-1D* and *rga-Δ17* mutants, or in the presence of 1 μM PAC (Fig. 3.2B).

To determine if the regulation of these genes by GAI was direct, we constructed a glucocorticoid-inducible version of *gai-1D* by fusing it to the rat glucocorticoid receptor domain (*GR*) (Lloyd et al., 1994; Aoyama and Chua, 1997), under the control of the *GAI* promoter. As expected, induction of *gai-1D* translocation into the nucleus by dexamethasone application caused repression of *IAA19/MSG2*, *IAA5* and *IAA6/SHY1*, and this repression was not blocked by cycloheximide (Fig. 3.2C).




**Fig. 3.2 Regulation of Aux/IAA gene expression by DELLA proteins.**

(A) Expression of *IAA5*, *IAA6* and *IAA19* determined by RT-qPCR in 3-day-old dark-grown *HS::gai-1D* seedlings after a 30-min heat shock at 37°C. Values at each time-point are relative to the expression of each gene in seedlings not subject to heat shock (A). Error bars represent s.e.m. (n=3 biological replicates).

(B) The expression of *IAA5*, *IAA6*, and *IAA19* was determined by RT-qPCR in 3-day-old wild-type and mutant seedlings grown in darkness. The values indicate the expression level of each gene relative to the value of the wild type grown in MS. Error bars represent SD (n=3).

(C) Expression of *IAA5*, *IAA6* and *IAA19* determined by RT-qPCR in 3-day-old dark-grown seedlings expressing a fusion between *gai-1D* and the glucocorticoid receptor (*GR*) under the control of the *GAI* promoter. 3-day-old dark-grown seedlings were transferred into flasks and incubated with soft shaking for 6 h with a mock solution or with 10 μM dexamethasone (DEX) to allow *gai-1D:GR* moving to the nucleus and regulate target genes. Alternatively, seedlings were also incubated with 10 μM cycloheximide (CHX) to prevent de novo protein synthesis during mock or DEX treatments. Values are relative to the expression of each gene in mock-treated seedlings. Error bars represent s.e.m. (n=3 biological replicates).

The repression of *IAA19/MSG2* expression upon GA deficiency was also evident in etiolated seedlings harbouring a transcriptional fusion between the *IAA19/MSG2* promoter and the *GUS* reporter gene (Fig.3.3A). Higher expression levels were detected, as previously reported (Tatematsu et al., 2004), in the apical part of dark-grown seedlings, and the level of expression decreased upon PAC application in a dose-dependent manner (Fig 3.3B).




**Fig. 3.3 Transcriptional regulation of *MSG2/IAA19* by GA in etiolated seedlings**

**(A)** Transcriptional regulation of *MSG2/IAA19* by GA in etiolated seedlings during gravitropic reorientation. PAC concentration was 1  $\mu\text{M}$ .  $\text{GA}_3$  concentration was 10  $\mu\text{M}$  **(B)** Localization of *MSG2/IAA19* expression in etiolated seedlings. Photographs reproduce the reported expression of *MSG2* in the apex of hypocotyls and in the upper part of the roots. Scale bars represent 0.5 mm.



The observation that *IAA19/MSG2* is a direct target for GAI transcriptional regulation provides a likely mechanism for the attenuation of gravitropism by GAs. The dominant *msg2-1* mutation that prevents *IAA19/MSG2* destabilization by auxin has been shown to impair gravitropic responses presumably by blocking ARF activity (Tatematsu et al., 2004). Thus, to investigate the degree of involvement of *IAA19/MSG2* in the repression of the gravitropic response by GAs, we asked whether DELLA accumulation would alleviate the agravitropic phenotype of the seedlings that carry the hyperstable allele *msg2-1*. In agreement with our observation that PAC decreases the activity of the *IAA19/MSG2* promoter (Fig. 3.4), growth of etiolated seedlings in the presence of 0.4  $\mu$ M PAC prevented the accumulation in the nuclei of *msg2-1::GFP* protein expressed from the *IAA19/MSG2* promoter (Muto et al., 2007) (Fig. 3.4A). And, concurrently, this treatment restored an almost normal reorientation capacity to *msg2-1* mutant seedlings (Fig. 3.4B). We thus conclude that GAs modulate gravitropism at least partly through the transcriptional regulation of *IAA19/MSG2* expression.




**Fig. 3.4 Involvement of *IAA19/MSG2* in the regulation of gravitropism by GA.**

(A) Reduction of *IAA19/Msg2-1* protein levels caused by impairment of GA biosynthesis. A *pMSG2::msg2-1::GFP* transgenic line was grown for 3 days in darkness with and without 0.4  $\mu$ M PAC, and GFP fluorescence in hypocotyls was visualized under a confocal microscope. (B) Gravitropic reorientation of hypocotyls of *msg2-1* seedlings. Seedlings were grown for 3 days in darkness on control media or in media supplemented with 0.4  $\mu$ M PAC, and the plates were turned 90° at time 0. Reorientation of the hypocotyls was monitored in darkness as explained in Materials and Methods. Error bars represent SD (n>16 individual seedlings).

### 3.3.3. Physiological relevance of the regulation of gravitropism by gibberellins

The results shown here indicate that GAs –and hence the level of DELLA proteins– influence the gravitropic response of aerial tissues, but what is the physiological relevance of this regulation? Can changes in DELLA abundance within the physiological range still affect gravitropism? Indeed, seedlings of the quadruple DELLA KO mutant did suffer a delay in the gravitropic response when grown in the light (Fig. 3.5A), a situation that causes DELLA accumulation in the wild type (Achard et al., 2007), confirming that the ability to respond to gravistimulation directly depends on the relative concentration of DELLA proteins. In agreement with this, a quadruple DELLA KO mutant did not show any difference, compared to the wild type, in the speed or extent of reorientation after gravistimulation in darkness, when GAs are not in limiting concentrations (Fig. 3.).



**Fig. 3.5** Gravitropic reorientation of hypocotyls of quadruple *della* (*gai rga rgl1 rgl2*) knockout mutants.

Seedlings were grown for 3 days in darkness on MS, and the plates were turned 90° at time 0. In (A), seedlings were exposed to 150  $\mu\text{mol m}^{-2} \text{s}^{-1}$  of white light during 8 h prior to reorientation, while in (B), dark-grown seedlings were used. All values between 2 and 16 hours in (A) are statistically different ( $p < 0.01$ ) between the *della* mutant and the parental control. PAC concentration was 0.4  $\mu\text{M}$ . Error bars represent SD ( $n > 46$  individual seedlings). The behaviour of the *della* mutant was statistically significant ( $p < 0.05$ ) with respect to the wild type as indicated by a two-way ANOVA test.

However, a more critical consequence of the attenuation of auxin responsiveness by GAs is the increase in variance of the gravitropic response when large populations of *Arabidopsis* seedlings were examined right after germination. As shown in Fig. 3.6A, individual seedlings grown for 3 days in darkness displayed certain degree of

inclination (between 5 and 10°) with respect to the gravity vector under normal GA concentrations, but this variation practically disappeared when GA biosynthesis or GA signaling were compromised (with PAC, or in the *gai-ID* mutant). The mechanism by which GAs cause this variance is very likely equivalent to the one through which GAs control gravitropic reorientation (i.e., through IAA19/MSG2), given that *msg2-1* mutants showed a much larger variance compared to the wild type, which was consequently reduced by PAC (Fig. 3.6A).


To study whether the increase in variance of the gravitropic response caused by GAs could confer any adaptive advantage, we examined the behavior of seedling populations grown under two competing signals: a light source perpendicular to the gravity vector (i.e., photo- versus gravitropism). As shown in Fig. 3.6B, the increased gravitropic response caused by the presence of PAC, led to a reduction in phototropic orientation in wild-type seedlings. This effect was also observed when phototropic reorientation was monitored after illumination of dark-grown seedlings with lateral blue light (Fig. 3.6C). On the contrary, the aphototropic *phototropin1* (*phot1*) mutant, impaired in the main light receptor that regulates phototropism (Esmon et al., 2005), still responded to PAC with a severe increase in gravitropic response arguing against the direct regulation of phototropism by GAs (Tsuchida-Mayama et al., 2010). Competition between gravi- and phototropism has been proposed to be mediated by the phyA photoreceptor (Lariguet and Fankhauser, 2004; Whippo and Hangarter, 2004; Iino, 2006). Interestingly, the *phytochrome A* (*phyA*) mutant shows reduced response to PAC (Fig. 3.6B) suggesting a connection between GA action and the regulation of gravitropism by phyA. These observations highlight a specific role of GA-induced regulation of gravitropic response in a situation of competing environmental signals.

### **3.4. Discussion**

The work presented here reveals an unexpected role for GAs in the control of the response of plants to gravity, and highlights the physiological relevance of a novel interaction between DELLA proteins and the expression of *Aux/IAA* genes.

The main line of evidence that supports the relevance of IAA19/MSG2 in the control of gravitropism by GAs is the observation that the agravitropic phenotype caused by the dominant *msg2-1* allele was alleviated by inhibiting GA biosynthesis

(Fig. 3.4B). This result at least indicates that transcriptional regulation of *IAA19/MSG2* by GAs has a significant impact in the gravitropic response of etiolated seedlings. Our work also shows that the regulation of *IAA19/MSG2* expression by DELLA proteins does not require protein synthesis (Fig. 3.2C).



**Fig. 3.6 Physiological relevance of the regulation of gravitropism by GAs.**

(A) Distribution of hypocotyl orientation of 3-day-old seedlings in darkness in the presence and absence of PAC. Seedlings were grown on vertical MS plates and the angle with respect to a horizontal line was recorded, with 90° being perfectly vertical.

(B) Distribution of hypocotyl orientation of 3-day-old seedlings under continuous unilateral blue light (10 nmol m<sup>-2</sup>s<sup>-1</sup>), coming from the left as indicated by the arrow. The angle with respect to a horizontal line was recorded, with 90° being perfectly vertical. Boxes represent the middle quartiles around the median (bisecting line), while the whiskers represent the upper and lower quartiles. The data represent the values of 65-130 seedlings.

(C) Effect of GA deficiency on phototropic reorientation. Seedlings were grown for 3 days in darkness in the presence or absence of 0.4  $\mu$ M PAC and 10  $\mu$ M  $GA_4$ , and illuminated with 10  $\mu$ mol m<sup>-2</sup> s<sup>-1</sup> lateral blue light at time 0. Monitorization of reorientation was carried out as explained in Materials and Methods.

Since DELLA proteins do not bind DNA directly, it is highly likely that they regulate transcription of *IAA9/MSG2* through the interaction with other transcription factors. Two such mechanisms have been proposed so far: the inhibition of DNA binding of members of the PHYTOCHROME-INTERACTING FACTOR (PIF) family of bHLH transcription factors (de Lucas et al., 2008; Feng et al., 2008), and the inhibitory interaction with JASMONATE-ZIM-DOMAIN (JAZ) proteins (Hou et al., 2010), which in turn regulate MYC2 activity in jasmonic acid signaling. There are no indications that JA signaling is involved in gravitropic responses, but it is reasonable to think that DELLA proteins interact with PIFs already present at high levels in dark-grown seedlings to directly regulate the expression of the target *Aux/IAA* genes. In agreement with this model, the expression of *IAA19* is strongly reduced in etiolated *pifq* mutants (Leivar et al., 2009), although there is no experimental evidence to date showing direct interaction between any PIF transcription factor and the *IAA19* promoter.

A critical issue in the control of the gravitropic response is the spatial localization of the machinery that perceives gravity and directs reorientation. Starch-loaded amyloplasts have been shown to be an integral part of the mechanism that allows gravity perception (Boonsirichai et al., 2003), and cells accumulating amyloplasts are located in the tip of the roots and in the endodermis of aerial tissues, such as the hypocotyl. Given that GAs affect gravitropic responses in hypocotyls, it is tempting to suggest that GA signaling interferes with the early events after gravity perception in the endodermis, but this hypothesis requires additional experimental evidence.

In any case, our observation that GAs regulate the gravitropism at an early stage does not diminish the role of the auxin gradient as the driving force in the orientation of the plant with respect to the gravity vector. Rather, GAs would act by fine tuning the formation of this gradient and modulating the responsiveness to this gradient in auxin responding cells. However, we cannot rule out that GAs also affect auxin relocalization, therefore establishing a reinforcing mechanism.

Interestingly, our results do not support an important role of GAs in the execution of cell expansion during gravitropic reorientation, but highlight a function of GAs in the generation of variance in the response. A higher degree of variance or noise in biological responses is a trait often selected by nature, and proposed to be at the core

of the mechanisms that drive robust morphogenesis (Houchmandzadeh et al., 2002; Yucel and Small, 2006), and speciation (Braendle and Felix, 2008). In microorganisms, even adaptation of a population to the environment seems to be based, to some extent, on cell-to-cell variance within genetic circuits (Balaban et al., 2004; Sanchez and Kondev, 2008). Not many molecular mechanisms have been proposed that explain the generation of noise in cellular systems (Casal et al., 2004), and the attenuation of auxin response by GAs might represent one of such mechanisms to provide flexibility in situations under which plants face competing tropic signals. Such could be the case of plants that must optimize their access to light because of neighbors' proximity. The observation that DELLA protein concentration decreases in seedlings during the shade avoidance response (Djakovic-Petrovic et al., 2007) supports this scenario. In that case, part of the shade avoidance mechanism may involve the attenuation of gravitropism to allow bending against the gravity vector.

### 3.5. Materials and methods

#### Plants and growth conditions

*Arabidopsis thaliana* GA signaling dominant mutants *gai-1D* and *rga-Δ17*, and the quadruple loss-of-function *rga-24 gai-t6 rgl2-1 rgl1-1* (Cheng et al., 2004) are in the *Ler* background, while the other lines used in this work, such as *msg2-1* (Tatematsu et al., 2004), *phot1* (Huala et al., 1997), *phyA* (Nagatani et al., 1993), *HS::gai-1D* (Alabadí et al., 2008), *pMSG2::GUS* (Tatematsu et al., 2004), and *pMSG2::msg2-1::GFP* (Muto et al., 2007), are derived from Col-0 accession. For all experiments, seeds were surface sterilized and stratified for 4-7 days at 4°C in darkness. Germination was induced under continuous white fluorescent light (90–100  $\mu\text{mol m}^{-2}\text{s}^{-1}$ ) for 8 hours, and then the plates were kept in darkness at 22°C.

#### Construction of transgenic lines

To obtain the transgenic line pGAI::gai-1D:GR, the *gai-1D* coding sequence was amplified from genomic DNA of the *gai-1D* mutant with primers: GAId-BamHI-F (GGA TCC ATG AAG AGA GAT CAT CAT CAT CA) and GAId-SacI-R (GAG CTC ATT GGT GGA GAG TTT CCA AGC CGA) and cloned into the pCR2.1 vector

(Invitrogen), and the *Bam*HI-*Sac*I fragment was subcloned into pGreen0029-35S::GR (Hellens et al., 2000) to give rise to pG35::gai-1D:GR. The *GAI* promoter was PCR amplified from genomic DNA using oligos pGAI-KpnI-F (GGT ACC TGG GAC CAC AGT CTA AAT GGC GT) and pGAI-XbaI-R (TCT AGA GGT TGG TTT TTT TTC AGA GAT GGA), cloned into the pCR2.1 vector, and the *Kpn*I-*Xba*I fragment transferred into pG35::gai-1D:GR to construct pGAI::gai-1D:GR. *Agrobacterium* C58 pSOUP cells were transformed with pGAI::gai-1-GR and *Arabidopsis* Ler plants were transformed using the floral dip method (Clough and Bent, 1998). Transgenic seedlings with a 3:1 segregation ratio for kanamycin resistance conferred by the transgene marker were selected for further work.

### **Gene expression analysis**

Total RNA extraction, cDNA synthesis and quantitative PCR, were carried out as described previously (Frigerio et al., 2006) using *EF1- $\alpha$*  expression for normalization. The primers used were: *IAA19* (CTC GGG CTT GAG ATA ACG GA and CCA CAT CTC TCC CCG GAA), *IAA5* (AAC TAC GGC TAG GTC TTC CCG and AGA TGG ACT CAC CGG AGA CG) and *IAA6* (TGG CAA AGG AAG GTC TAG CAC and TGG AAG ACC CAA TCG AAG CT).

### **GUS staining was carried out as described previously (Frigerio et al., 2006).**

Detailed description of the microarray experiment will be published somewhere else. Briefly, *HS::gai-1D* and Col-0 seedlings were grown for 3 days at 22°C in darkness on half strength MS medium (Duchefa) with 0.8% w/v agar and 1% w/v sucrose, and then the plates were transferred to 37°C for 30 min. Samples were collected before, and 1, 2, 4 h after the beginning of the heat treatment. Three independent biological replicates were used for the analysis. RNA amplification, labeling, and hybridization of microarray slides (70-mer oligonucleotide arrays that represent the majority of the *Arabidopsis* genes: <http://www.ag.arizona.edu/microarray>) were carried out as previously described (Bueso et al., 2007). Hybridization was set up to isolate genes differentially expressed in *HS::gai-1D* compared to Col-0 at each time point.

### **Tropism tests**

To determine the angle of seedling emergence, seedlings were grown for 3 days in darkness at 22°C in a vertical orientation on plates containing half strength MS medium (Duchefa) with 0.8% w/v phytoagar and without sucrose, and supplemented with mock or 0.4 µM PAC. After 3 days the plates were photographed.

For reorientation experiments, the conditions were the same as above but after 3 days of growth the plates were reoriented 90° relative to the initial growth angle. Reorientation was recorded every hour under infrared light using CCD cameras coupled to Metamorph software as described by Schepens et al (Schepens et al., 2008).

To test blue light-induced phototropism, seedlings were grown for 3 days under 10 nmol m<sup>-2</sup>s<sup>-1</sup> of continuous unilateral blue light at 22°C in a vertical orientation on plates containing half strength MS medium (Duchefa) with 0.8% w/v phytoagar and without sucrose and supplemented with mock or 0.4 µM PAC. After 3 days, the plates were photographed.

In all cases, angles were measured using Image J software.

### **Acknowledgements**

We thank Kiyosi Tatematsu and the Arabidopsis Biological Resource Center for providing us with mutant and transgenic lines. We also thank François Parcy (PSV-CNRS, Grenoble) for fruitful discussions and hosting MAB, and Cristina Ferrándiz (IBMCP, Valencia) and Detlef Weigel (MPI for Developmental Biology, Tübingen) for very useful comments on the manuscript. JGB is the recipient of a CSIC JAE Fellowship.

## **3.6. Bibliography**

- Achard P, Liao L, Jiang C, Desnos T, Bartlett J, Fu X, Harberd NP (2007) DELLAs Contribute to Plant Photomorphogenesis. *Plant Physiol* **143**: 1163-1172
- Alabadí D, Gallego-Bartolomé J, García-Cárcel L, Orlando L, Rubio V, Martínez C, Frigerio M, Iglesias-Pedraz JM, Espinosa A, Deng XW, Blázquez MA (2008) Gibberellins modulate light signaling pathways to prevent Arabidopsis seedling de-etiolation in darkness. *Plant J* **53**: 324-335
- Aoyama T, Chua NH (1997) A glucocorticoid-mediated transcriptional induction system in transgenic plants. *Plant J* **11**: 605-612
- Balaban NQ, Merrin J, Chait R, Kowalik L, Leibler S (2004) Bacterial persistence as a phenotypic switch. *Science* **305**: 1622-1625



- Boonsirichai K, Sedbrook JC, Chen R, Gilroy S, Masson PH** (2003) ALTERED RESPONSE TO GRAVITY is a peripheral membrane protein that modulates gravity-induced cytoplasmic alkalization and lateral auxin transport in plant statocytes. *Plant Cell* **15**: 2612-2625
- Braendle C, Felix MA** (2008) Plasticity and errors of a robust developmental system in different environments. *Dev Cell* **15**: 714-724
- Bueso E, Alejandro S, Carbonell P, Perez-Amador MA, Fayos J, Belles JM, Rodriguez PL, Serrano R** (2007) The lithium tolerance of the *Arabidopsis* cat2 mutant reveals a cross-talk between oxidative stress and ethylene. *Plant J* **52**: 1052-1065
- Casal JJ, Fankhauser C, Coupland G, Blázquez MA** (2004) Signalling for developmental plasticity. *Trends Plant Sci* **9**: 309-314
- Cheng H, Qin L, Lee S, Fu X, Richards DE, Cao D, Luo D, Harberd NP, Peng J** (2004) Gibberellin regulates *Arabidopsis* floral development via suppression of DELLA protein function. *Development* **131**: 1055-1064
- Clough SJ, Bent AF** (1998) Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J* **16**: 735-743
- Cowling RJ, Harberd NP** (1999) Gibberellins control *Arabidopsis* hypocotyl growth via regulation of cellular elongation. *J Exp Bot* **50**: 1351-1357
- Darwin C** (1880) *The Power of Movement in Plants*. John Murray, London
- de Lucas M, Davière JM, Rodríguez-Falcón M, Pontin M, Iglesias-Pedraz JM, Lorrain S, Fankhauser C, Blázquez MA, Titarenko E, Prat S** (2008) A molecular framework for light and gibberellin control of cell elongation. *Nature* **451**: 480-484
- Dill A, Jung HS, Sun TP** (2001) The DELLA motif is essential for gibberellin-induced degradation of RGA. *Proc Natl Acad Sci U S A* **98**: 14162-14167
- Djakovic-Petrovic T, de Wit M, Voesenek LA, Pierik R** (2007) DELLA protein function in growth responses to canopy signals. *Plant J* **51**: 117-126
- Esmon CA, Pedmale UV, Liscum E** (2005) Plant tropisms: providing the power of movement to a sessile organism. *Int J Dev Biol* **49**: 665-674
- Esmon CA, Tinsley AG, Ljung K, Sandberg G, Hearne LB, Liscum E** (2006) A gradient of auxin and auxin-dependent transcription precedes tropic growth responses. *Proc Natl Acad Sci U S A* **103**: 236-241
- Feng S, Martinez C, Gusmaroli G, Wang Y, Zhou J, Wang F, Chen L, Yu L, Iglesias-Pedraz JM, Kircher S, Schafer E, Fu X, Fan LM, Deng XW** (2008) Coordinated regulation of *Arabidopsis thaliana* development by light and gibberellins. *Nature* **451**: 475-479
- Frigerio M, Alabadí D, Pérez-Gómez J, García-Cárcel L, Phillips AL, Hedden P, Blázquez MA** (2006) Transcriptional regulation of gibberellin metabolism genes by auxin signaling in *Arabidopsis*. *Plant Physiol* **142**: 553-563
- Friml J, Wisniewska J, Benkova E, Mendgen K, Palme K** (2002) Lateral relocation of auxin efflux regulator PIN3 mediates tropism in *Arabidopsis*. *Nature* **415**: 806-809
- Harper RM, Stowe-Evans EL, Luesse DR, Muto H, Tatematsu K, Watahiki MK, Yamamoto K, Liscum E** (2000) The NPH4 locus encodes the auxin response factor ARF7, a conditional regulator of differential growth in aerial *Arabidopsis* tissue. *Plant Cell* **12**: 757-770
- Hellens RP, Edwards EA, Leyland NR, Bean S, Mullineaux PM** (2000) pGreen: a versatile and flexible binary Ti vector for *Agrobacterium*-mediated plant transformation. *Plant Mol Biol* **42**: 819-832
- Hou X, Lee LY, Xia K, Yan Y, Yu H** (2010) DELLAs modulate jasmonate signaling via competitive binding to JAZs. *Dev Cell* **19**: 884-894
- Houchmandzadeh B, Wieschaus E, Leibler S** (2002) Establishment of developmental precision and proportions in the early *Drosophila* embryo. *Nature* **415**: 798-802
- Huala E, Oeller PW, Liscum E, Han IS, Larsen E, Briggs WR** (1997) *Arabidopsis* NPH1: a protein kinase with a putative redox-sensing domain. *Science* **278**: 2120-2123
- Iino M** (2006) Toward understanding the ecological functions of tropisms: interactions among and effects of light on tropisms. *Curr Opin Plant Biol* **9**: 89-93

- Kim TW, Lee SM, Joo SH, Yun HS, Lee Y, Kaufman PB, Kirakosyan A, Kim SH, Nam KH, Lee JS, Chang SC, Kim SK** (2007) Elongation and gravitropic responses of Arabidopsis roots are regulated by brassinolide and IAA. *Plant Cell Environ* **30**: 679-689
- Lariguet P, Fankhauser C** (2004) Hypocotyl growth orientation in blue light is determined by phytochrome A inhibition of gravitropism and phototropin promotion of phototropism. *Plant J* **40**: 826-834
- Leivar P, Tepperman JM, Monte E, Calderon RH, Liu TL, Quail PH** (2009) Definition of Early Transcriptional Circuitry Involved in Light-Induced Reversal of PIF-Imposed Repression of Photomorphogenesis in Young Arabidopsis Seedlings. *Plant Cell* **21**: 3535-3553
- Li L, Xu J, Xu ZH, Xue HW** (2005) Brassinosteroids stimulate plant tropisms through modulation of polar auxin transport in Brassica and Arabidopsis. *Plant Cell* **17**: 2738-2753
- Lloyd AM, Schena M, Walbot V, Davis RW** (1994) Epidermal cell fate determination in Arabidopsis: patterns defined by a steroid-inducible regulator. *Science* **266**: 436-439
- Meudt WJ** (1987) Investigations on the Mechanism of the Brassinosteroid Response: VI. Effect of Brassinolide on Gravitropism of Bean Hypocotyls. *Plant Physiol* **83**: 195-198
- Muto H, Watahiki MK, Nakamoto D, Kinjo M, Yamamoto KT** (2007) Specificity and similarity of functions of the Aux/IAA genes in auxin signaling of Arabidopsis revealed by promoter-exchange experiments among MSG2/IAA19, AXR2/IAA7, and SLR/IAA14. *Plant Physiol* **144**: 187-196
- Nagatani A, Reed JW, Chory J** (1993) Isolation and Initial Characterization of Arabidopsis Mutants That Are Deficient in Phytochrome A. *Plant Physiol* **102**: 269-277
- Overvoorde PJ, Okushima Y, Alonso JM, Chan A, Chang C, Ecker JR, Hughes B, Liu A, Onodera C, Quach H, Smith A, Yu G, Theologis A** (2005) Functional genomic analysis of the AUXIN/INDOLE-3-ACETIC ACID gene family members in Arabidopsis thaliana. *Plant Cell* **17**: 3282-3300
- Peng J, Carol P, Richards DE, King KE, Cowling RJ, Murphy GP, Harberd NP** (1997) The Arabidopsis GAI gene defines a signaling pathway that negatively regulates gibberellin responses. *Genes Dev* **11**: 3194-3205
- Rashotte AM, Brady SR, Reed RC, Ante SJ, Muday GK** (2000) Basipetal auxin transport is required for gravitropism in roots of Arabidopsis. *Plant Physiol* **122**: 481-490
- Sanchez A, Kondev J** (2008) Transcriptional control of noise in gene expression. *Proc Natl Acad Sci U S A* **105**: 5081-5086
- Schepens I, Boccalandro HE, Kami C, Casal JJ, Fankhauser C** (2008) PHYTOCHROME KINASE SUBSTRATE4 Modulates Phytochrome-Mediated Control of Hypocotyl Growth Orientation. *Plant Physiol* **147**: 661-671
- Schwechheimer C** (2008) Understanding gibberellic acid signaling--are we there yet? *Curr Opin Plant Biol* **11**: 9-15
- Tatematsu K, Kumagai S, Muto H, Sato A, Watahiki MK, Harper RM, Liscum E, Yamamoto KT** (2004) MASSUGU2 encodes Aux/IAA19, an auxin-regulated protein that functions together with the transcriptional activator NPH4/ARF7 to regulate differential growth responses of hypocotyl and formation of lateral roots in Arabidopsis thaliana. *Plant Cell* **16**: 379-393
- Tsuchida-Mayama T, Sakai T, Hanada A, Uehara Y, Asami T, Yamaguchi S** (2010) Role of the phytochrome and cryptochrome signaling pathways in hypocotyl phototropism. *Plant J*
- Úbeda-Tomás S, Swarup R, Coates J, Swarup K, Laplaze L, Beemster GT, Hedden P, Bhalerao R, Bennett MJ** (2008) Root growth in Arabidopsis requires gibberellin/DELLA signalling in the endodermis. *Nat Cell Biol* **10**: 625-628
- Whippo CW, Hangarter RP** (2004) Phytochrome modulation of blue-light phototropism. *Plant Cell Environ* **27**: 1223-1228
- Yucel G, Small S** (2006) Morphogens: precise outputs from a variable gradient. *Curr Biol* **16**: R29-31

**Zentella R, Zhang ZL, Park M, Thomas SG, Endo A, Murase K, Fleet CM, Jikumaru Y, Nambara E, Kamiya Y, Sun TP** (2007) Global analysis of della direct targets in early gibberellin signaling in Arabidopsis. *Plant Cell* **19**: 3037-3057



## Chapter 4



# Hierarchy of hormone action controlling apical hook development in *Arabidopsis*

Gallego-Bartolomé J, Arana MV, Vandenbussche F, Zádňíková P, Minguet EG, Guardiola V, Van Der Straeten D, Benkova E, Alabadí D, Blázquez MA.

**Plant J.** doi: 10.1111/j.1365-313X.2011.04621.x. (2011)



#### **4.1. Abstract**

The apical hook develops in the upper part of the hypocotyl when seeds buried in the soil germinate, and serves to protect cotyledons and the shoot apical meristem from possible damage caused by pushing through the soil. The curvature is formed through differential cell growth occurring at the two opposite sides of the hypocotyl, and it is established by a gradient of auxin activity and refined by the coordinated action of auxin and ethylene. Here we show that gibberellins (GAs) promote hook development through the transcriptional regulation of several genes of the ethylene and auxin pathways in *Arabidopsis*. The level of GA activity determines the speed of hook formation and the extent of the curvature during the formation phase independently of ethylene, likely by modulating auxin transport and response through *HLS1*, *PIN3*, and *PIN7*. Moreover, GAs cooperate with ethylene in preventing hook opening, in part through the induction of ethylene production mediated by *ACS5/ETO2* and *ACS8*.

## 4.2. Introduction

The acquisition of developmental innovations has accompanied the evolution of land plants (Langdale, 2008). A key innovation in seed plants is skotomorphogenesis (Wei *et al.*, 1994), an alternative to photomorphogenesis when seeds face germination in darkness, for example when they are buried in the soil. Importantly, skotomorphogenesis provides protection to emerging seedlings while pushing through the soil, especially to the shoot apical meristem (SAM) and cotyledons (Kami *et al.*, 2010). In dicotyledonous plants, these vital structures are protected by an apical hook in the hypocotyl that “pulls” them through the soil. Indeed, hookless mutants are not able to emerge when seeds germinate buried in the soil (Harpham, 1991).

The apical hook is mainly formed through differential elongation between the cells at opposite sides of the hypocotyl (Raz and Ecker, 1999). Hook development follows three phases: formation, maintenance, and opening (Raz and Ecker, 1999; Vandenbussche *et al.*, 2010; Zadnikova *et al.*, 2010). The formation phase extends from the time when germination is completed until the hook curvature reaches  $\sim 180^\circ$  and it usually takes  $\sim 24$  h in *Arabidopsis thaliana*. Then, the curvature is actively maintained in parallel to extensive hypocotyl elongation. Hook maintenance can be interrupted by light, and then full opening is completed typically in 6 h (Liscum and Hangarter, 1993; Wu *et al.*, 2010). If seedlings are kept in the dark, the hook is maintained for 24 h, and opening is completed 70-90 h later (Vandenbussche *et al.*, 2010; Zadnikova *et al.*, 2010).

The differential cell growth that underlies hook development is caused by an asymmetrical accumulation of auxin (Kuhn and Galston, 1992; Lehman *et al.*, 1996). Pharmacological treatments or mutations that affect either auxin accumulation (Boerjan *et al.*, 1995; Zhao *et al.*, 2001; Stepanova *et al.*, 2008), transport (Lehman *et al.*, 1996; Chaabouni *et al.*, 2009; Vandenbussche *et al.*, 2010; Zadnikova *et al.*, 2010), or signaling (Stowe-Evans *et al.*, 1998; Nagpal *et al.*, 2000; Li *et al.*, 2004; Tatematsu *et al.*, 2004; Yang *et al.*, 2004; Zadnikova *et al.*, 2010) influence apical hook development. Auxin accumulation marks the side with the lower growth rate in the apical hook (Kuhn and Galston, 1992; Raz and Ecker, 1999).



Besides auxin, other hormones participate in apical hook development. For example, exogenous treatment with ethylene induces the formation of exaggerated hooks whereas ethylene insensitive mutants are hookless (Guzman and Ecker, 1990). Similarly, gibberellins (GAs) are also required for correct hook development, given that a block in either GA synthesis or signaling results in a hookless phenotype (Achard *et al.*, 2003; Alabadí *et al.*, 2004; Vriezen *et al.*, 2004).

The concurrence of multiple hormones controlling a given output is a common theme in plant development (Alabadí and Blázquez, 2009), although their precise mode of action is not always clear. For instance, in the case of hook development, ethylene influences the auxin pathway (Li *et al.*, 2004; Stepanova *et al.*, 2008; Vandenbussche *et al.*, 2010; Zadnikova *et al.*, 2010), suggesting that ethylene requires auxin to control hook formation; but on the other hand, ethylene application is able to reverse the hook phenotype of the auxin mutant *nph4* (Harper *et al.*, 2000). Additionally, GAs act through ethylene in the control of hook development (Achard *et al.*, 2003; Vriezen *et al.*, 2004), but no molecular mechanism has been found yet.

To unveil the hierarchy of hormone action during hook development, we have investigated in detail the requirement for each hormone in a dynamic way from the time of hook formation to its opening phase, we have searched for gene targets downstream of GA action in the context of hook development, and we have tested the physiological relevance for this regulatory interactions *in vivo*.


### **4.3. Results**

#### **4.3.1. Dynamics of GA-regulated apical hook development**

To determine the phase of apical hook development in which GA activity is required, we performed a kinematic analysis of this process in *Ler* wild type plants untreated and treated with the GA biosynthesis inhibitor paclobutrazol (PAC), as well as in *gai-1* and quintuple *della* mutants. *gai-1* encodes a dominant version of the DELLA protein GAI that constitutively inhibits GA signaling; the *della* mutant, which lacks all DELLA proteins of *Arabidopsis*, shows a fully activated GA pathway (Peng *et al.*, 1997; Feng *et al.*, 2008). Untreated wild type seedlings displayed the three phases of

hook development (Figure 4.1a) (Vandenbussche et al., 2010; Zadnikova et al., 2010). On the contrary, seedlings were not able to form the apical hook when treated with 0.2  $\mu$ M PAC; instead, they gradually entered into the opening phase (Figure 4.1a). *gai-1* mutants behaved similarly to PAC-treated seedlings although they started to form the hook, reaching a maximum angle of  $121.4 \pm 9.5^\circ$  20 h after germination (Figure 4.1b). Notably, *della* seedlings showed exaggerated apical hooks (the maximum angle was  $241.8 \pm 7.9^\circ$ ) as a consequence of a faster kinetics of hook formation during the initial phase, whereas they behaved as the wild type during the other phases.

These results indicate that GA signaling is both necessary and limiting during the formation phase, and therefore the magnitude of hook curvature depends on this activity during the initial phase. In addition, GA activity is also necessary, yet not limiting, to delay hook opening.



**Figure 4.1 Regulation of apical hook development by GAs and ethylene.**

(a) Kinematic analysis of hook development in *Ler* wild type seedlings mock-treated and treated with 0.2 mM PAC, as well as in mock-treated *gai-1* and *della* seedlings.

(b) Kinematic analysis of hook development in *Ler* wild type and *gai-1* seedlings grown on control medium or with 10 mM ACC. Dotted vertical lines represent the transition between phases. All error bars represent s.e.m. ( $n > 20$ ).


#### 4.3.2. GA control on hook development is dependent and independent upon ethylene activity

Exaggerated apical hooks also appear when ethylene activity is high (Guzman and Ecker, 1990). The exaggerated curvature in response to the ethylene precursor 1-aminocyclopropane-1-carboxylate acid (ACC) was due to a delay in the transition between formation and maintenance phases (Figure 4.2a) (Vandenbussche et al., 2010; Zadnikova et al., 2010). Importantly, it was the level of GA activity, and not of

ethylene, which set up the speed of hook formation (Figures 4.2a,c and 4.1b). This suggests that both hormones act through different mechanisms during the initial phase, since ethylene is also necessary for hook formation (Vandenbussche *et al.*, 2010). To test if GA-mediated hook formation depends to some extent on ethylene activity, we analyzed hook development in the ethylene insensitive mutant *ein2-1* (Guzman and Ecker, 1990). *ein2-1* seedlings failed to complete hook formation (Vandenbussche *et al.*, 2010), whereas it was partially restored by GA-treatment (Figure 4.1c).

Analysis of mutants with low or null hormone activity suggested that both hormones are important to prevent hook opening (Figure 4.1a,c) (Vandenbussche *et al.*, 2010). The kinetics of hook opening was very similar in *della* and in wild type seedlings, and it remained unaltered when the latter were treated with a saturating amount of ACC (Figure 4.1b). Remarkably, the exaggerated hooks of *della* seedlings did not open after ACC-treatment (Figure 4.1b).

These results indicate that 1) GAs determine the rate of the hook formation and the extent of the curvature reached during this phase; 2) this role is partially independent of ethylene; 3) ethylene is necessary to complete this phase, although the response seems saturated; and 4) both hormones act jointly to prevent hook opening.



**Figure 4.2 Regulation of apical hook development by GAs and ethylene.**

(a,b) Kinematic analysis of hook development in *Ler* wild type and *della* seedlings grown on control medium or with 10  $\mu$ M ACC (b), as well as Col-0 wild type and *ein2-1* seedlings grown on control medium or with 50  $\mu$ M gibberellic acid ( $GA_3$ ) (c). Dotted vertical lines represent the transition between phases. All error bars represent s.e.m. ( $n>20$ ).

#### 4.3.3. The expression of *ACS5/ETO2*, *ACS8*, and *HLS1* genes is regulated by the GA pathway

To elucidate the molecular mechanism by which GAs regulate hook development, we searched through microarray analysis for genes that could be relevant for this process among those rapidly regulated by *gai-1* in 2-day-old *pHsp::gai-1* etiolated seedlings (Alabadí *et al.*, 2008) (Chapter 2: Gallego-Bartolomé, Alabadí, Blázquez, unpublished). We found that the ethylene biosynthesis genes *ACC SYNTHASE8* (*ACS8*) and *ACS5/ETO2* (Vogel *et al.*, 1998; Yamagami *et al.*, 2003), and the ethylene-induced gene *HOOKLESS1* (*HLS1*) (Lehman *et al.*, 1996), were downregulated by *gai-1*. Analyses in *pHsp::gai-1*, *ProRGA:GFP-(rga-Δ17)*, and *Pro35S:gai-1* lines (Alabadí *et al.*, 2008; Dill *et al.*, 2001), and in *gai-t6 rga-24* double loss-of-function mutants (Dill and Sun, 2001; King *et al.*, 2001) confirmed their regulation by DELLAs (Figure 4.3a,b).

Their rapid response to *gai-1* suggested that they might be direct targets. To confirm this, we examined their expression in *ProGAI:gai-1-GR* seedlings (Gallego-Bartolomé, Alabadí, Blázquez, submitted). As a control, we included the DELLA-induced gene *AtGA20ox2* gene in the analysis (Zentella *et al.*, 2007) (Figure 4.3b). Dexamethasone (DEX)-treatment repressed and induced *HLS1* and *AtGA20ox*, respectively, and this effect was not abolished by cycloheximide (CHX) indicating that regulation by *gai-1* is independent of protein synthesis (Figure 4.3c). However, downregulation of *ACS5/ETO2* and *ACS8* by *gai-1* requires the synthesis of a protein intermediate. The strong upregulation of *ACS8* by CHX could mask any effect of *gai-1*, and therefore we could not rule out the possibility of a direct effect of the DELLA protein. The transcription factor PIF5 promotes *ACS8* expression in etiolated seedlings (Khanna *et al.*, 2007). Since DELLAs regulate transcription by inhibiting several transcription factors of the PIF clade (de Lucas *et al.*, 2008; Feng *et al.*, 2008; Arnaud *et al.*, 2010), we tested whether this is the case for PIF5. GAI and PIF5 interacted *in vivo* in *Nicotiana benthamiana* leaves as shown by co-immunoprecipitation (Figure 4.3d) and bimolecular fluorescence complementation (BiFC) (Figure 4.S1). Remarkably, chromatin immunoprecipitation (ChIP) showed that PIF5 binds *in vivo* to a G-box in the *ACS8* promoter in a GA-dependent manner in *Arabidopsis* (Figure 4.3e), suggesting DELLAs may repress *ACS8* expression by inhibiting PIF5.


ACS5/ETO2- and ACS8-mediated ethylene production contributes to hook development (Vogel et al., 1998; Tsuchisaka et al., 2009), and the activity of *HLS1* is central to mediate fully this effect (Roman et al., 1995; Lehman et al., 1996). Thus, our gene expression analysis suggests that GAs regulate hook development through the control of *HLS1* gene expression through direct regulation by DELLA proteins and via ethylene biosynthesis (Figure 4.3f).

#### **4.3.4. GA-regulation of *ACS5/ETO2* and *ACS8* gene expression depends on the phase of hook development**

To examine the temporal and spatial distribution of *ACS5/ETO2* and *ACS8* expression during hook development and their response to GAs, we used the *ProACS5:GUS* and *ProACS8:GUS* reporters (Tsuchisaka and Theologis, 2004). Their spatial and temporal expression patterns were similar (Figures 4.4a,b and 4.S2). Staining was detected mainly in the hypocotyl vasculature, reaching the apical hook 36 h after germination. Both the timing and the extent of their response to GAs were somewhat different. The regulation of *ProACS5:GUS* expression upon GAs was evident 36 h after germination. Remarkably, GAs became limiting 36 h later, when GA-treatment resulted in augmented expression (Figure 4.4a). The dependence of *ProACS8:GUS* on GAs was also evident 36 h after germination (Figure 4.4b), although the response was already saturated. As expected, the PAC-effect on both reporter lines was reversed completely by simultaneous treatment with GAs (Figure 4.S2). Hence, both the basal expression and the responsiveness to GAs of *ACS5/ETO2* and *ACS8* are subject to developmental regulation in the apical hook.

#### **4.3.5. GAs support ethylene production in etiolated seedlings**

Staining patterns of *ProACS5:GUS* and, to a lesser extent, of *ProACS8:GUS* in response to GAs support the idea that GAs promote ethylene biosynthesis in etiolated seedlings. To test it we measured ethylene production in etiolated *Ler* wild type and *della* seedlings. The ability of wild type seedlings to produce ethylene decreased steadily during the first days after germination (Figure 4.4c). This trend was reversed in *della* seedlings, which produced more ethylene than the wild type after the second day. This timing is coincident with the dependence of *ACS8* and *ACS5/ETO2* expression



**Figure 4.3** GAs regulate the ethylene pathway in etiolated seedlings.

(a) Expression of *ACS5/ETO2*, *ACS8*, and *HLS1* in 2-d-old *pHsp::gai-1* seedlings subjected to a 30 min treatment at 37°C; control seedlings were kept at 20°C. Expression was determined by qRT-PCR and normalized to the respective control treatment.


(b) Thirty six-hour-old wild type *Ler* and *gai-t6 rga-24* seedlings were grown on control medium or with 0.2 μM PAC. Expression was determined by qRT-PCR. PAC, fold change between PAC- and mock-treated wild type *Ler* seedlings; *Pro35S:gai-1*, fold change between transgenic and wild type Col-0 seedlings; *rga-Δ17*, fold change between *ProRGF:GFP(rga-Δ17)* and wild type *Ler* seedlings; *gai-t6 rga-24* mock, fold change between *gai-t6 rga-24* and wild type *Ler* seedlings; *gai-t6 rga-24* PAC, fold change between PAC-treated and mock-treated *gai-t6 rga-24* seedlings.

(c) Two-day-old *ProGAI:gai-1-GR* etiolated seedlings were incubated for 5 h in water or in water supplemented with either 10 μM DEX, 10 μM CHX, or both. (a-c) Expression was determined by qRT-PCR and normalized to the respective control treatment. Data represent mean and standard deviation of three technical replicates. Experiments were repeated twice with similar results.

(d) co-IP showing the interaction between GAI and PIF5. YFP-GAI and HA-PIF5 were expressed either alone or together in leaves of *Nicotiana benthamiana*. Nuclear proteins were immunoprecipitated with anti-HA antibody-coated paramagnetic beads and detected by immunoblotting with either anti-HA or anti-GFP antibodies.

(e) qRT-PCR of a regulatory (G-box) or a control (ORF) sequence in the *ACS8* locus after ChIP with anti-HA. Analysis was performed in 36-hour-old Col-0 wild type and *Pro35S:PIF5-HA* seedlings grown on control medium or with 0.2 μM PAC. Enrichment of the regulatory and control ORF sequences is shown after normalization to the input value. Data represent mean and standard deviation of three technical replicates from a representative experiment out of three biological replicates.

(f) Model: GAs control hook development by transcriptional regulation of *HLS1*, either directly or indirectly through regulation of ethylene biosynthesis.



**Figure 4.4 Regulation of the ethylene pathway by GAs.**

(a,b) Expression patterns of *ProACS5::GUS* (a) and *ProACS8::GUS* (b) during hook development in seedlings grown on control medium or with 0.2  $\mu M$  PAC or 50  $\mu M$  GA<sub>3</sub>.


(c) GAs promote ethylene production in etiolated seedlings. The ability to produce ethylene per day was measured in wild type *Ler* and quintuple *della* etiolated seedlings. Three independent sets of biological material were used for calculating mean values. Error bars represent s.e.m. The experiments were done twice with similar results.

upon GA activity (Figure 4.4a,b). Thus, the GA pathway may contribute to reach the minimum threshold level of ethylene needed to sustain a proper transition to hook maintenance and to delay hook opening in the wild type.

#### 4.3.6. GAs regulate partly hook development by modulating PIF activity

The regulation of *ACS8* by the DELLA-PIF5 interaction (Figures 4.3d,e and 4.S1), together with the fact that PIF1, PIF3, and PIF5 promote hook development (Khanna et al., 2007; Leivar et al., 2008; Kim et al., 2011) suggests that PIFs could mediate the GA-regulation of this process. Indeed, *pif5* mutants showed a slight hypersensitivity in PAC-induced repression of *ACS8* and hook opening, whereas *Pro35S::PIF5-HA* seedlings were resistant (Figure 4.5a,b). In additional support of this hypothesis, *pif1 pif3 pif4 pif5* (*pif1/3/4/5*) seedlings (Leivar et al., 2008; Shin et al., 2009) did not form the apical hook and they immediately entered into the opening phase, whilst GA-treatment delayed hook opening for a few hours (Figure 4.5c). Analysis of the *pif3/4/5* mutant corroborated the significant role of PIF1 in this process,

since these seedlings were able to delay the opening phase (Figure 4.5d). Remarkably, PIF1 was able to restore the GA-responsiveness during the formation phase. These results indicate that PIF activity is necessary at least for hook formation and that there is a temporal coincidence in the need of GA and PIF activities, suggesting a functional relationship in the control of this process.



**Figure 4.5 The activity of PIF transcription factors mediate the GA control on hook development.**

(a) qRT-PCR analysis of *ACS8* expression in 3-day-old wild type Col-0, *pif5*, and *Pro35S:PIF5-HA* seedlings grown on control medium (M) or with 0.2  $\mu$ M PAC. All data were normalized to the expression value in the control wild type. Data represent mean and standard deviation of three technical replicates. Experiments were repeated twice with similar results.

(b) Hook angle of 3-day-old wild type Col-0, *pif5*, and *Pro35S:PIF5-HA* seedlings grown on control medium (M) or with 0.2  $\mu$ M PAC. Error bars represent s.e.m. ( $n > 20$ ).

(c,d) Kinematic analysis of hook development in Col-0 wild type and *pif1/3/4/5* (c) and *pif3/4/5* (d) seedlings grown on control medium or with 50  $\mu$ M GA<sub>3</sub>. Dotted vertical lines represent the transition between phases. Error bars represent s.e.m. ( $n > 20$ ).

#### 4.3.7. HLS1 activity mediates GA effect on hook development

The partially ethylene-independent control of GAs on hook formation (Figure 4.2a) is consistent with a model by which GAs regulate *HLS1* directly (Figure 4.3f), and with GA activity being necessary to allow ethylene to exert its control on apical hooking (Achard *et al.*, 2003; Vriezen *et al.*, 2004). One-day-resolution analysis of hook development indicated that *HLS1* is needed early after germination in the dark




(Raz and Ecker, 1999). Our kinematic analysis confirmed previous results showing that *hls1-1* mutation prevented hook formation (Figure 4.6a). The dynamics of hook development was very similar in *hls1-1* mutants and in PAC-treated seedlings (Figures 4.1a and 4.6a), indicating that there is a temporal coincidence in the requirement of both activities during hook development. Besides, the hook phenotype of *hls1-1* seedlings was not affected by exogenous GA-treatment, whereas the wild type showed exaggerated hooks (Figure 4.6a).

To confirm that GAs regulate hook development through HLS1, we analyzed the effect that uncoupling *HLS1* expression from GA-regulation had on the GA-control of hook development. For that purpose, we prepared *Pro35S:YFP-HLS1* transgenic lines and analyzed their response to PAC. As hypothesized, Figure 4.6b shows that apical hooks of *Pro35S:YFP-HLS1* seedlings were partially resistant to PAC-induced opening. Furthermore, time-course analysis of *HLS1* expression showed that GA activity is needed to sustain its expression during hook development (Figure 4.6c). Nonetheless, *HLS1* transcript level was not increased in *della* mutants indicating that its regulation by GAs is already saturated.

#### **4.3.8. GAs are needed to sustain differential auxin response during apical hook development**

Asymmetrical auxin accumulation and response is essential for the differential cell growth underlying apical hook development (Lehman et al., 1996; Li et al., 2004; Vandenbussche et al., 2010; Wu et al., 2010; Zadnikova et al., 2010). Moreover, HLS1 is critical to establish the auxin response in the hook, since the asymmetric distribution of *ProDR5:GUS* staining in the apical hook is lost in *hls1* (Li et al., 2004). Given the regulation of *HLS1* expression by GAs, we examined whether the *ProDR5:GUS* response was altered by GAs. By 18 h after germination, *ProDR5:GUS* staining was apparent at the concave side of the hook in control seedlings (Figure 4.7a,b) (Vandenbussche et al., 2010; Zadnikova et al., 2010).



**Figure 4.6** HLS1 activity mediates the GA control on hook development.

(a) Kinematic analysis of hook development in Col-0 wild type and *hls1-1* seedlings grown on control medium or with 50  $\mu$ M GA<sub>3</sub>. Dotted vertical lines represent the transition between phases. Error bars represent s.e.m. (n>20).

(b) Hook angle of 1-day-old wild type Col-0 and *Pro35S:HLS1* seedlings grown on control medium or with 0.05 or 0.2  $\mu$ M PAC. Error bars represent s.e.m. (n>20).


(c) qRT-PCR analysis of *HLS1* expression during hook development in wild type *Ler* seedlings grown on control medium (M) or with 0.2  $\mu$ M PAC, as well as in quintuple *della* seedlings. Thirty-six and 72 h data points were normalized to the expression value in the control wild type at the time point 18 h. Data represent mean and standard deviation of three technical replicates. Experiments were repeated twice with similar results.

Neither the intensity of the staining at the concave side nor the number of seedlings with differential staining was influenced by GA-treatment at this stage of development. Nevertheless, the percentage of seedlings with staining at the inner side of the hook was lower after treatment with 0.2  $\mu$ M PAC. This result suggests that GAs are necessary to support differential auxin response during the formation phase. Stronger GA-dependence was observed during the maintenance and opening phases. At these two stages no *ProDR5:GUS* expression was detected at the upper zone of the hypocotyl of any PAC-treated seedling, where the apical hook should form, whereas GA-treatment enhanced the differential *ProDR5:GUS* staining at the concave side of the hook (Figure 4.7a,b). As expected, the PAC-effect was reversed completely by simultaneous treatment with GAs (Figure 4.S2).

Remarkably, the *ProDR5:GUS* expression pattern is very similar in PAC-treated (Figure 4.7a) and in *hls1* seedlings (Li *et al.*, 2004). Despite the driving role proposed for HLS1 during apical hook development, its activity is not sufficient in the absence of polar auxin transport (Lehman *et al.*, 1996). In agreement, ACC-treatment does not revert the effects of the polar auxin transport inhibitor naphthylphthalamic acid (NPA) (Zadnikova *et al.*, 2010). Similarly, 50  $\mu$ M GA<sub>3</sub>-treatment did not revert either the hookless phenotype or the altered *ProDR5:GUS* staining pattern caused by NPA-treatment (Figure 4.7c,d), which suppressed the exaggerated hooks of *della* seedlings (Figure 4.7d). The effects of GA- and ethylene-treatments on *ProDR5:GUS* during maintenance and opening phases are similar (Vandenbussche *et al.*, 2010; Zadnikova *et al.*, 2010). Nevertheless, GAs might control auxin response independently of ethylene during the formation phase (Figure 4.2a). In fact, whereas indole-3-acetic acid (IAA)-treatment restores the apical hook to ethylene-insensitive mutants (Vandenbussche *et al.*, 2010), it was not able to restore it to PAC-treated seedlings and to *hls1-1* mutants (Figure 4.7e). In summary, these results draw new similarities between GAs and HLS1 activity, which suggests that they participate in the same pathway in the establishment and/or the interpretation of the auxin gradient during apical hook development.

#### **4.3.9. GAs participate in maintaining *PIN3* and *PIN7* expression in the apical hook**

Genetic analyses have implicated AUX1, LAX3, PIN1, PIN3, PIN4, and PIN7 in driving the auxin flux during apical hook development, and ethylene regulates the transcription of several of their genes (Vandenbussche *et al.*, 2010; Zadnikova *et al.*, 2010). We asked whether GAs would also influence the expression of these genes. Expression of *PIN1*, *PIN4*, and *AUX1* was not altered by GAs during hook development (data not shown). Sustained expression of *PIN3* was dependent upon GAs during the maintenance and opening phases, whereas this dependence was evident earlier for *PIN7* (Figure 4.8a,b). These results are consistent with results of Figure 4.7e, and suggest that GAs might also promote hook development by maintaining proper expression of *PIN* genes needed to distribute the auxin flux from cotyledons (Zadnikova *et al.*, 2010).



**Figure 4.7** GAs regulate the differential auxin response in the apical hook.

(a,b) Expression pattern of *ProDR5:GUS* during hook development in seedlings grown on control media or with 0.2  $\mu$ M PAC or 50  $\mu$ M GA<sub>3</sub>. Pictures of representative seedlings are shown (a). The percentage of seedlings showing DR5 signal at the inner side of the hook is represented in (b). Data are mean of three biological replicates,  $n > 25$  each. Error bars are s.d.

(c,d) Polar auxin transport mediates the GA regulation on hook development. Pictures of representative 1-day-old wild type Col-0 seedlings grown in control medium or with 50  $\mu$ M GA<sub>3</sub>, 5  $\mu$ M NPA, or both (c). Hook angle of 1-day-old *Ler* wild type and *della* seedlings grown in control medium or with 50  $\mu$ M GA<sub>3</sub>, 5  $\mu$ M NPA, or both (d).

(e) Hook angle of 1-day-old Col-0 wild type and *hls1-1* seedlings grown in control medium or with 0.1  $\mu$ M IAA, 0.2  $\mu$ M PAC, or both. All error bars represent s.e.m. ( $n > 20$ ).

To challenge this hypothesis, we investigated the response of *pin3 pin7* mutants to GAs. Double mutant seedlings were not able to complete hook formation and, importantly, they were resistant to GA-treatment (Figure 4.8e). Interestingly, single mutants had contrasting behaviors: *pin3* mutants showed a milder defect on hook formation than *pin7*, whereas their response to GAs was quite affected; *pin7* seedlings responded to GAs similarly to the wild type despite having more disturbed hook formation than *pin3* (Figure 4.8c,d).

#### **4.3.10. GA activity in the endodermis is required for apical hook development**


Misexpression approaches have shown that the context outlined by the cell type may be determinant to define the output of hormone pathways (Jaillais and Chory, 2010). For instance, DELLA activity in the endodermis controls meristem size and overall growth in the root (Úbeda-Tomás et al., 2008; Ubeda-Tomas et al., 2009), whereas the epidermis is the key tissue for brassinosteroids to control shoot growth (Savaldi-Goldstein et al., 2007). Thus, we examined whether GAs regulate hook development in a tissue-specific manner. We expressed *gai-1* exclusively in the endodermis under the control of the *SCARECROW* promoter (*ProSCR:gai-YFP-GR*) (Úbeda-Tomás et al., 2008), or in the epidermis under the control of the *MERISTEM LAYER1* promoter (*ProML1:GFP-gai-1*; Figure 4.S3).

Expression of *gai-1* in the endodermis but not in the epidermis impaired hook formation similar to the PAC-treatment or the *gai-1* mutation (Figure 4.9a). Since the *SCR* promoter is active in the hook endodermis starting 22 h after germination (Vandenbussche et al., 2010), our results indicate that GA activity is necessary in the endodermis for the correct progression of hook development at least during the late formation phase, whereas it is dispensable in the epidermis.

These results support further the functional relationship between GAs and PIFs sustaining hook development, since expression of *PIF1* only in the endodermis of the *pif1/3/4/5* mutant restores the hook (Kim et al., 2011), indicating there is also a spatial coincidence in the requirement of both activities.

Next, to place the transcriptional network regulated by GAs in the context of the endodermis, we examined the activity of *ProDR5:GUS*, *ProPIN3:GUS*, and *ProPIN7:GUS* in F1 seedlings from crosses between the reporter lines and *Ler* wild type, *ProML1:GFP-gai-1-11*, and *ProSCR:gai-YFP-GR* seedlings. Impairing GA signaling in the endodermis had the same effect on the expression of *ProDR5:GUS* and *ProPIN3:GUS* than PAC-treatment, whereas no effect was observed when GA signaling was blocked in the epidermis (Figure 4.9b). A tissue-independent effect was observed, however, when *ProPIN7:GUS* expression was examined. These results suggest that GAs control *PIN3* expression mainly from the endodermis and that confinement of its expression to the vascular bundle by PAC-treatment or *ProSCR:gai-YFP-GR* expression (see a magnification in Figure 4.9c), may impair to some extent the auxin

flux towards outer tissues, in agreement with the disappearance of *ProDR5:GUS* from the concave side.



**Figure 4.8 The contribution of PIN3 and PIN7 to GA-mediated hook development.**


(a,b) Expression patterns of *ProPIN3:GUS* (a) and *ProPIN7:GUS* (b) during hook development in seedlings grown on control medium or with 0.2  $\mu\text{M}$  PAC or 50  $\mu\text{M}$  GA<sub>3</sub>.

(c,d) Kinematic analysis of hook development in Col-0 wild type and *pin3-5* (c) and *pin7-1* (d) seedlings grown on control medium or with 50  $\mu\text{M}$  GA<sub>3</sub>. Dotted vertical lines represent the transition between phases. Error bars represent s.e.m. (n>15).

Kinematic analysis of hook development in Col-0 wild type and *pin3-3 pin7<sup>En</sup>* double mutant seedlings grown on control medium or with 50  $\mu\text{M}$  GA<sub>3</sub>. Dotted vertical lines represent the transition between phases. Error bars represent s.e.m. (n>15).

In support of this, PIN3 is present in endodermis, cortex, and epidermis, whereas PIN7 and PIN4 are predominant in outer tissues (Zadnikova et al., 2010). The mild hook phenotype of *pin3* mutants indicate that other efflux carriers are involved, although less relevant for the GA-control on the hook. Moreover, GAs may impinge on

other branches of the network, most likely *HLS1*, to regulate hook development from the endodermis.



**Figure 4.9 GA activity in the endodermis controls hook development.**

(a) Hook curvature was measured in 1-day-old *Ler* wild type seedlings grown on control medium or in medium with 10  $\mu$ M DEX or with 0.2  $\mu$ M PAC; in *gai-1*, *ProML1:GFP-gai-1-4* and *ProML1:GFP-gai-1-11* (*ML1:gai*) seedlings grown on control medium, and in *ProSCR:gai-YFP-GR* (*SCR:gai-GR*) seedlings grown on control medium or with 10  $\mu$ M DEX. All error bars represent s.e.m. ( $n > 20$ ). (b,c) GUS staining of 1-day-old F1 etiolated seedlings from the crosses indicated in the main text, grown on control medium or in medium with 10  $\mu$ M DEX or with 0.2  $\mu$ M PAC (b). See a magnification of regions within orange squares in (c). Pictures of representative seedlings are shown.

## 4.4. Discussion

The establishment of an apical hook is an intrinsic part of the skotomorphogenic developmental program and it depends on differential cell elongation on opposite sides of hypocotyls. The instructive molecular framework that guarantees this differential growth relies in the end on asymmetrical auxin response (Lehman *et al.*, 1996). Ethylene signaling represents one module of regulation that sustains this basic framework (Stepanova *et al.*, 2008; Vandenbussche *et al.*, 2010; Zadnikova *et al.*, 2010), in a large part targeting *HLS1* transcription (Li *et al.*, 2004; Chaabouni *et al.*, 2009). Our results show that GAs impinge both on the ethylene pathway and on auxin distribution and response, and therefore it represents a new layer of regulation that ensures proper progression through all phases of hook development (Figure 4.10).

### 4.4.1. GAs regulate hook formation independently of ethylene activity

Sustained asymmetric auxin activity is necessary during all phases for proper hook development (Lehman *et al.*, 1996; Chaabouni *et al.*, 2009; Wu *et al.*, 2010; Zadnikova *et al.*, 2010). Ethylene plays its major role in a time-window that encompasses maintenance and opening phases and overlaps with a period of augmented sensitivity to the hormone (Raz and Ecker, 1999), whereas its role during the formation phase is minor (Figure 4.10b) (Raz and Ecker, 1999; Knee *et al.*, 2000; Vandenbussche *et al.*, 2010; Zadnikova *et al.*, 2010). On the contrary, the GA pathway performs a prominent role during the initial phase, when the strength of its activity determines the speed of hook formation and the extent of hook curvature (Figure 4.1a). Importantly, this role of GAs is mostly independent of ethylene (Figure 4.1c). The high demand of GA activity for apical hooking is reminiscent of germination. The apical hook starts to form immediately after germination in darkness is completed. Germinating seeds require high levels of GAs to break dormancy (Ogawa *et al.*, 2003; Cao *et al.*, 2005; Penfield *et al.*, 2006). Our results suggest that this high GA activity might extend into the early stages of hook development to ensure a sustained GA response. Both processes may have similar mechanistic basis, the same GA response initiated in embryos during germination may continue later on in etiolated seedlings to promote apical hook development. In agreement, mutants with a hyperactive GA pathway show exaggerated growth of the embryo's axis (Cao *et al.*, 2005) and exaggerated hook



curvature (Figure 4.1a). Moreover, GA biosynthesis and response take place mainly in the hypocotyl endodermis and cortex during germination (Yamaguchi et al., 2001; Ogawa et al., 2003). Remarkably, sustained GA activity specifically in hypocotyl endodermis is required for proper progression through hook formation (Figure 4.9).

#### **4.4.2. GAs prevent hook opening in cooperation with the ethylene pathway**

GAs are also required to prevent hook opening. This task is performed jointly with ethylene, and the transition to this phase is prevented only when the two hormones become not limiting (Figure 4.2b). This response suggests that this process might be controlled by a signaling element whose activity is regulated in cooperation by both pathways. For example, DELLA proteins could inactivate an ethylene-regulated transcription factor that negatively regulates opening, similar to their negative effect on PIFs (de Lucas *et al.*, 2008; Feng *et al.*, 2008). The apical hook, on the other hand, is not a vital structure when seedlings grow *in vitro*. The timing and kinetics of hook opening may respond solely to endogenous cues under these conditions. The identification of GAs and ethylene as elements imposing a brake to hook opening suggests that both pathways are targets of light signaling during de-etiolation. In fact, the GA pathway is downregulated by light (Reid et al., 2002; Achard et al., 2007; Zhao et al., 2007; Alabadí et al., 2008), which might help to turn off the hormonal network that prevents hook opening (see below). The activity of ethylene is high in etiolated seedlings (Zhong *et al.*, 2009), so it is reasonable to think that it is also reduced during de-etiolation. Indeed, light impinges negatively on ethylene signaling rather on ethylene levels to promote hook opening in *Arabidopsis* (Knee *et al.*, 2000). Besides, the expression of the ethylene- and GA-induced gene *HLS1* is repressed by light, which surely contributes to hook opening (Li *et al.*, 2004).

#### **4.4.3. GAs regulate hook development by transcriptional regulation of auxin and ethylene pathways**


How do GAs regulate progression through hook development? Our results indicate that GAs exert this regulation, or at least part of it, by transcriptional regulation of several elements of the signaling network that controls apical hooking. First, GAs impinge on the core of the mechanism by regulating expression of auxin transporter

genes *PIN3* and *PIN7* (Figure 4.8). Second, GAs influence the expression of two *ACS* genes involved in ethylene biosynthesis, *ACS5/ETO2* and *ACS8* (Figures 4.3a-c and 4.3a,b), as well as the expression of the ethylene-induced gene *HLS1* (Figures 4.3a-c and 4.6c), whose activity is necessary to control auxin responses in the hook (Lehman *et al.*, 1996; Li *et al.*, 2004). The kinetics of their transcriptional response suggests that DELLAs operate through different regulatory mechanisms depending on each case. Regulation of *PIN3* and *PIN7* seems an indirect consequence of DELLAs' activity (data not shown). A similar case is found at the root meristem, where DELLAs downregulate *PIN* expression indirectly through *ARR1* and *SHY2* (Dello Ioio *et al.*, 2008; Moubayidin *et al.*, 2010). The downregulation of *HLS1* and *ACS8* is a direct consequence, whilst the fast regulation of *ACS5/ETO2* requires the synthesis of a protein intermediate (Figure 4.3c). Remarkably, DELLAs directly inhibit the activity of *PIF5* to repress the expression of *ACS8* (Figure 4.3d,e), as previously seen with *PIF3* and *PIF4* for light-regulated genes (de Lucas *et al.*, 2008; Feng *et al.*, 2008). The expression of both *HLS1* and *ACS5/ETO2* is lower in *pif1/3/4/5* mutants than in the wild type (Leivar *et al.*, 2009; Shin *et al.*, 2009), suggesting that PIFs mediate their regulation by DELLAs as well. Nonetheless, the influence of PIFs may be indirect given that there are no G-boxes in the upstream promoter region of both genes.

Several pieces of evidence support the idea that regulation of *ACS* genes by GAs is relevant for ethylene production in etiolated seedlings. First, the *della* mutant produces more ethylene than the wild type (Figure 4.4c). Second, the timing for increased ethylene production in *della* mutants correlates with the increased expression of *ACS5/ETO2* upon GA-treatments (Figure 4.4a); the contribution of *ACS8* activity to the extra ethylene in the *della* mutant may be lower. Third, this timing also coincides with the window of maximum ethylene sensitivity in the apical hook (Raz and Ecker, 1999). And fourth, *ACS5/ETO2* and *ACS8* contribute to ethylene-induced hook development (Vogel *et al.*, 1998; Tsuchisaka *et al.*, 2009).

The close connection of GAs with the auxin and ethylene pathways (Figure 4.10) is manifested by the strong hook phenotype observed when the GA activity is compromised. Despite the role of the GA-mediated ethylene production may be minor, the regulation of *HLS1* and the auxin transporters surely have a deep contribution to hook development. For instance, the hookless phenotype caused by low GA levels is alleviated by overexpressing *HLS1* (Figure 4.6b). This idea is supported further by the

staining patterns of *ProDR5::GUS* which are shared by PAC- or NPA-treatment (Figure 5a,c) and the *hls1* mutant (Li et al., 2004), and by the inability of IAA-treatment to restore the apical hook to PAC-treated and *hls1* seedlings (Figure 4.7e). We propose that GAs sustain differential auxin transport and response during the formation phase and that at least the latter might be mediated by HLS1 activity. This is based in three observations: first, there is a coincidence in the temporal requirement of HLS1 and GA activities during hook formation (Figures 4.1a and 4.6a). Second, *hls1* is epistatic over GA-application (Figure 4.6a). And third, *HLS1* expression is directly downregulated by DELLAs (Figure 4.3c). Notwithstanding, whereas GA activity is limiting to drive hook formation (Figure 1a), it is saturated to promote *HLS1* expression (Figure 4.6c). This suggests that there is another mechanism by which GAs regulate the formation phase besides transcriptional regulation of the *HLS1* gene.



**Figure 4. 10 Models explaining the pathway interactions and the timing of GA and ethylene action.**

(a,b) In etiolated seedlings proper activity of auxin is crucial for hook development (a). Its activity is sustained by GAs and ethylene at different levels, including auxin biosynthesis, transport, and response. Part of the GA control is exerted from the endodermis, for instance transcriptional regulation of *PIN3*. GAs and ethylene may exert this role independently or through common downstream signaling elements. Light act negatively on several branches of the hormonal network to promote hook opening. The contribution of the activity of the GA and ethylene pathways is different depending on the phase of hook development (b). GAs promote hook formation partly in a ethylene-independent manner, likely through *HLS1*, and the contribution of ethylene to this phase seems to be minor, whereas both pathways cooperate to prevent hook opening.

Note: References for Figure 4.10 are included separately at the end of the references section.

## 4.5. Experimental procedures

### Plant lines and growth conditions

*Arabidopsis thaliana* accessions *Ler* and *Col-0* were used as wild types. Mutants and transgenic lines used have been described: quintuple *della* (Feng *et al.*, 2008), *gai-1* (Peng *et al.*, 1997), *gai-t6 rga-24* (Dill and Sun, 2001; King *et al.*, 2001), *pRGA::GFP-(rga-Δ17)* (Dill *et al.*, 2001), *ProHsp:gai-1* and *Pro35S:gai-1* (Alabadí *et al.*, 2008), *ProSCR:gai-YFP-GR* (Úbeda-Tomás *et al.*, 2008), and *ProGAI:gai-1-GR* (Gallego-Bartolomé, Alabadí, Blázquez, submitted); *ein2-1* and *hls1-1* (Guzman and Ecker, 1990), and *pACS5::GUS* and *ProACS8:GUS* (Tsuchisaka and Theologis, 2004); *ProPIN7:GUS*, *pin7-1*, and *pin3-5* (Benkova *et al.*, 2003), *ProPIN3:GUS* (Friml *et al.*, 2002); *Pro35S:PIF5-HA* (Lorrain *et al.*, 2008), and *pif3/4/5* and *pif1/3/4/5* (Leivar *et al.*, 2008; Shin *et al.*, 2009). The *pin3-3 pin7<sup>En</sup>* double mutant has been kindly provided by Dr Ykä Helariutta (Helsinki University).

Seeds were sterilized and stratified for 6 days in water at 4°C. Germination took place under white fluorescent light (90–100  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ) at 22°C for 6 h in a Percival growth chamber E-30B (<http://www.percival-scientific.com>). Seeds were plated in plates of half-strength MS medium with 0.8% (w/v) agar and 1% (w/v) sucrose supplemented with either 0.2  $\mu\text{M}$  PAC, 50  $\mu\text{M}$  GA<sub>3</sub>, 10  $\mu\text{M}$  ACC, 10  $\mu\text{M}$  DEX, 0.1  $\mu\text{M}$  IAA or 5  $\mu\text{M}$  NPA and grown in darkness at 22°C. For exogenous GA-treatment, seeds were stratified in 50  $\mu\text{M}$  GA<sub>3</sub>. For short-term treatments, seedlings were incubated in the dark in water supplemented with 10  $\mu\text{M}$  CHX and/or 10  $\mu\text{M}$  DEX. MS, PAC, GA<sub>3</sub>, ACC, IAA and NPA were from Duchefa (<http://www.duchefa.com>). DEX and CHX were from Sigma (<http://www.sigmaaldrich.com>). Plates were placed vertically for kinematic analyses.

### Real-time analysis of apical hook development

Real-time imaging of apical hook development and hook angle measurement were performed as described (Vandenbussche *et al.*, 2010; Zadnikova *et al.*, 2010).

### **Analysis of reporter lines**

$\beta$ -glucuronidase (GUS) staining was performed as described (Zadnikova et al., 2010).

### **Construction of vectors and generation of transgenic lines**

The pENTR223 vector carrying the *HLS1* or ORF was obtained from the Arabidopsis Biological Resource Center (ABRC) and transferred into the *pEarleyGate104* vector (Earley et al., 2006) by Gateway technology using the LR clonase (Invitrogen, <http://www.invitrogen.com>) to create *pEG::HLS1ox*.

The construction of *ProML1:GFP-gai-1* was as follows. The *gai-1* coding sequence was amplified from genomic DNA of the *gai-1* mutant with primers GAIdf (ATGAAGAGAGATCATCATCATCA) and GAIdr (ATTGGTGGAGAGTTTCCAAGCCGA) that included the attB1 and attB2 Gateway recombination sites (not shown), respectively. The PCR product was cloned into *pDONR221* (Invitrogen) by BP reaction, and then into the binary vector *pSBright:GFP* (Bensmihen et al., 2005) by LR reaction to give rise to *pSBright:GFP-gai-1* construct. The *ML1* promoter was PCR-amplified using primers described (An et al., 2004) and that included the *HindIII* recognition site. The PCR product was cloned into the *pCR2.1* vector and sequenced. After digestion with *HindIII*, the *ML1* promoter was cloned into the *HindIII* site of *pSBright:GFP-gai-1*, to create *ProML1:GFP-gai-1*.

Constructs were introduced in *Agrobacterium* strain C58 and used to transform *Arabidopsis* Col-0 wild type plants, *pEG:HLS1ox*, or *Ler*, *ProML1:GFP-gai-1*. Transgenic seedlings in the T<sub>1</sub> and T<sub>2</sub> generations were selected on 50  $\mu$ M glufosinate ammonium (Sigma). Transgenic lines with a 3:1 (resistant:sensitive) segregation ratio were selected, and 10 homozygous lines were identified in the T<sub>3</sub> generation. Data from two representative lines are shown.

### **Real-time quantitative RT-PCR**

RNA extraction, cDNA synthesis, quantitative RT-PCR (qRT-PCR), analysis, and primer sequences for amplification of *AtGA20ox2* and *EF1- $\alpha$*  genes have been described (Frigerio et al., 2006). qRT-PCR oligonucleotides for *ACS5/ETO2*, *ACS8*, and *HLS1* genes were: qRT-ACS5f (GCTGGTTCGACATCTGCGA), qRT-ACS5r (AGGCTCTGCAAGGCAAAACAT), qRT-ACS8f (GGTGCTACTCCGGCTAACGA), qRT-ACS8r (TCCAGGATCAGCGAGACAAAA), qRT-HLS1f (CGATACCGTC

CGTTTTTCGAA), and qRT-HLS1r (GCCTTAGCCAAGTTATGCGC).

### **Ethylene measurements**

Ethylene measurements were performed as described (Thain *et al.*, 2004), with the following modifications. 150-200 seeds were sterilized and sown in a 10 ml chromatography vial containing 5 ml of half-strength MS with 1% (w/v) sucrose and 0.8% (w/v) agar. The vial was kept 5 days at 4°C in darkness and subsequently exposed to white light for 6 h at 21°C to stimulate germination. Seedlings were grown in darkness (capped vials wrapped in aluminium foil). Every 24 h, the vials were flushed with hydrocarbon free air (Air Liquide, <http://www.es.airliquide.com/>) and ethylene in the headspace was detected with an ETD-300 photo-acoustic ethylene detector (Sensor Sense, <http://www.sense.com.br>).

### **Confocal microscopy**

Images were taken using a Leica TCS SL confocal laser microscope (Leica Microsystems GmbH, <http://www.leica-microsystems.com/>) with excitation at 488 nm.

### **BIFC and co-IP assays**

BIFC vectors *pMDC43-YFN* and *pMDC43-YFC* were provided by Dr Alejandro Ferrando (IBMCP). *pENTR* vectors carrying the coding sequence of *PIF5* and *GAI* were generated by the REGIA project (Paz-Ares and The Regia, 2002). *PIF5* and *GAI* coding sequences were transferred into *pMDC43-YFC* and *pMDC43-YFN*, and into *pEarleyGate201* and *pEarleyGate104* (Earley *et al.*, 2006) for BIFC and co-IP, respectively, by Gateway using the LR clonase (Invitrogen). Each construct was introduced into *Agrobacterium* C58 cells, which were used subsequently to infiltrate leaves of *Nicotiana benthamiana*. BIFC analysis was performed as described (Scacchi *et al.*, 2009).

For co-IP, nuclear proteins were isolated from formaldehyde-fixed leaves. Immunoprecipitation was carried out with anti-HA antibody-coated paramagnetic beads (Miltenyi Biotec, <http://www.miltenyibiotec.com/en/default.aspx>) following manufacturer's instructions. HA- and YFP-tagged proteins in the input and immunoprecipitated were detected by immunoblotting using anti-HA (Roche, <https://www.roche-applied-science.com>) and anti-GFP (Clontech, <http://www.clontech.com/>) antibodies.

### ChIP and PCR amplification

Seedlings of *Arabidopsis* Col-0 and *Pro35S:PIF5-HA* transgenic line were grown at 22°C for 3 days in darkness before fixation. ChIP assays were performed as described (Hornitschek *et al.*, 2009). qPCR oligonucleotides to amplify the region around the G-box were pACS8-F-1 (ATGGAAATTTCACATCGTGCCTA) and pACS8-R-1 (GATGTCAGAGAAGAATGAGCACGT). The ORF region was amplified with the same oligonucleotides used to analyze *ACS8* gene expression by RT-qPCR.

### ACKNOWLEDGEMENTS

We thank Dr Antonella Locascio for the generation of the GAI construct for BiFC, and the ABRC, Malcolm Bennett, Christian Fankhauser, Nicholas Harberd, Ykã Helariutta, Peter Quail, Tai-pin Sun, and Susana Úbeda-Tomás for seeds. Work in the laboratory of M.A.B. and D.A. is supported by grants from the Spanish Ministry of Science and Innovation (BIO2007-60293 and Consolider-TRANSPLANTA). Work in the laboratories of D.V.D.S. and E.B. is funded by the Research Foundation Flanders (FWO, projects G.0524.07 and G.0298.09) and by the European Research Council (Starting Independent Research Grant ERC-2007-Stg-207362-HCPO), respectively. J.G.-B. was supported by a JAE pre-doctoral fellowship from the CSIC. F.V. is a post-doctoral fellow of the Research Foundation Flanders (FWO).

### 4.6. Bibliography

- Achard P, Liao L, Jiang C, Desnos T, Bartlett J, Fu X, Harberd NP (2007) DELLAs Contribute to Plant Photomorphogenesis. *Plant Physiol* **143**: 1163-1172
- Achard P, Vriezen WH, Van Der Straeten D, Harberd NP (2003) Ethylene regulates arabidopsis development via the modulation of DELLA protein growth repressor function. *Plant Cell* **15**: 2816-2825
- Alabadí D, Blázquez MA (2009) Molecular interactions between light and hormone signaling to control plant growth. *Plant Mol Biol* **69**: 409-417
- Alabadí D, Gallego-Bartolomé J, García-Cárcel L, Orlando L, Rubio V, Martínez C, Frigerio M, Iglesias-Pedraz JM, Espinosa A, Deng XW, Blázquez MA (2008) Gibberellins modulate light signaling pathways to prevent Arabidopsis seedling de-etiolation in darkness. *Plant J* **53**: 324-335
- Alabadí D, Gil J, Blázquez MA, García-Martínez JL (2004) Gibberellins repress photomorphogenesis in darkness. *Plant Physiol* **134**: 1050-1057
- An H, Roussot C, Suarez-Lopez P, Corbesier L, Vincent C, Pineiro M, Hepworth S, Mouradov A, Justin S, Turnbull C, Coupland G (2004) CONSTANS acts in the

- phloem to regulate a systemic signal that induces photoperiodic flowering of *Arabidopsis*. *Development* **131**: 3615-3626
- Arnaud N, Girin T, Sorefan K, Fuentes S, Wood TA, Lawrenson T, Sablowski R, Ostergaard L** (2010) Gibberellins control fruit patterning in *Arabidopsis thaliana*. *Genes Dev* **24**: 2127-2132
- Benkova E, Michniewicz M, Sauer M, Teichmann T, Seifertova D, Jurgens G, Friml J** (2003) Local, efflux-dependent auxin gradients as a common module for plant organ formation. *Cell* **115**: 591-602
- Bensmihen S, Giraudat J, Parcy F** (2005) Characterization of three homologous basic leucine zipper transcription factors (bZIP) of the ABI5 family during *Arabidopsis thaliana* embryo maturation. *J Exp Bot* **56**: 597-603
- Boerjan W, Cervera MT, Delarue M, Beeckman T, Dewitte W, Bellini C, Caboche M, Van Onckelen H, Van Montagu M, Inze D** (1995) Superroot, a recessive mutation in *Arabidopsis*, confers auxin overproduction. *Plant Cell* **7**: 1405-1419
- Cao D, Hussain A, Cheng H, Peng J** (2005) Loss of function of four DELLA genes leads to light- and gibberellin-independent seed germination in *Arabidopsis*. *Planta* **223**: 105-113
- Chaabouni S, Jones B, Delalande C, Wang H, Li Z, Mila I, Frasse P, Latche A, Pech JC, Bouzayen M** (2009) SI-IAA3, a tomato Aux/IAA at the crossroads of auxin and ethylene signalling involved in differential growth. *J Exp Bot* **60**: 1349-1362
- de Lucas M, Davière JM, Rodríguez-Falcón M, Pontin M, Iglesias-Pedraz JM, Lorrain S, Fankhauser C, Blázquez MA, Titarenko E, Prat S** (2008) A molecular framework for light and gibberellin control of cell elongation. *Nature* **451**: 480-484
- Dello Ioio R, Nakamura K, Moubayidin L, Perilli S, Taniguchi M, Morita MT, Aoyama T, Costantino P, Sabatini S** (2008) A genetic framework for the control of cell division and differentiation in the root meristem. *Science* **322**: 1380-1384
- Dill A, Jung HS, Sun TP** (2001) The DELLA motif is essential for gibberellin-induced degradation of RGA. *Proc Natl Acad Sci U S A* **98**: 14162-14167
- Dill A, Sun T** (2001) Synergistic derepression of gibberellin signaling by removing RGA and GAI function in *Arabidopsis thaliana*. *Genetics* **159**: 777-785
- Earley KW, Haag JR, Pontes O, Oppen K, Juehne T, Song K, Pikaard CS** (2006) Gateway-compatible vectors for plant functional genomics and proteomics. *Plant J* **45**: 616-629
- Feng S, Martinez C, Gusmaroli G, Wang Y, Zhou J, Wang F, Chen L, Yu L, Iglesias-Pedraz JM, Kircher S, Schafer E, Fu X, Fan LM, Deng XW** (2008) Coordinated regulation of *Arabidopsis thaliana* development by light and gibberellins. *Nature* **451**: 475-479
- Frigerio M, Alabadi D, Pérez-Gómez J, García-Cárcel L, Phillips AL, Hedden P, Blázquez MA** (2006) Transcriptional regulation of gibberellin metabolism genes by auxin signaling in *Arabidopsis*. *Plant Physiol* **142**: 553-563
- Friml J, Wisniewska J, Benkova E, Mendgen K, Palme K** (2002) Lateral relocation of auxin efflux regulator PIN3 mediates tropism in *Arabidopsis*. *Nature* **415**: 806-809
- Guzman P, Ecker JR** (1990) Exploiting the triple response of *Arabidopsis* to identify ethylene-related mutants. *Plant Cell* **2**: 513-523
- Harper RM, Stowe-Evans EL, Luesse DR, Muto H, Tatematsu K, Watahiki MK, Yamamoto K, Liscum E** (2000) The NPH4 locus encodes the auxin response factor ARF7, a conditional regulator of differential growth in aerial *Arabidopsis* tissue. *Plant Cell* **12**: 757-770
- Harpham NVJ** (1991) The Effect of Ethylene on the Growth and Development of Wild-type and Mutant *Arabidopsis thaliana* (L.) Heynh. *Ann Bot* **68**: 55-61
- Hornitschek P, Lorrain S, Zoete V, Michielin O, Fankhauser C** (2009) Inhibition of the shade avoidance response by formation of non-DNA binding bHLH heterodimers. *Embo J* **28**: 3893-3902
- Jaillais Y, Chory J** (2010) Unraveling the paradoxes of plant hormone signaling integration. *Nat Struct Mol Biol* **17**: 642-645



- Kami C, Lorrain S, Hornitschek P, Fankhauser C** (2010) Light-regulated plant growth and development. *Curr Top Dev Biol* **91**: 29-66
- Khanna R, Shen Y, Marion CM, Tsuchisaka A, Theologis A, Schafer E, Quail PH** (2007) The basic helix-loop-helix transcription factor PIF5 acts on ethylene biosynthesis and phytochrome signaling by distinct mechanisms. *Plant Cell* **19**: 3915-3929
- Kim K, Shin J, Lee SH, Kweon HS, Maloof JN, Choi G** (2011) Phytochromes inhibit hypocotyl negative gravitropism by regulating the development of endodermal amyloplasts through phytochrome-interacting factors. *Proc Natl Acad Sci U S A* **108**: 1729-1734
- King KE, Moritz T, Harberd NP** (2001) Gibberellins are not required for normal stem growth in *Arabidopsis thaliana* in the absence of GAI and RGA. *Genetics* **159**: 767-776
- Knee EM, Hangarter RP, Knee M** (2000) Interactions of light and ethylene in hypocotyl hook maintenance in *Arabidopsis thaliana* seedlings. *Physiol Plant* **108**: 208-215
- Kuhn H, Galston AW** (1992) Physiological asymmetry in etiolated pea epicotyls: relation to patterns of auxin distribution and phototropic behavior. *Photochem Photobiol* **55**: 313-318
- Langdale JA** (2008) Evolution of developmental mechanisms in plants. *Curr Opin Genet Dev* **18**: 368-373
- Lehman A, Black R, Ecker JR** (1996) HOOKLESS1, an ethylene response gene, is required for differential cell elongation in the *Arabidopsis* hypocotyl. *Cell* **85**: 183-194
- Leivar P, Monte E, Oka Y, Liu T, Carle C, Castillon A, Huq E, Quail PH** (2008) Multiple phytochrome-interacting bHLH transcription factors repress premature seedling photomorphogenesis in darkness. *Curr Biol* **18**: 1815-1823
- Leivar P, Tepperman JM, Monte E, Calderon RH, Liu TL, Quail PH** (2009) Definition of Early Transcriptional Circuitry Involved in Light-Induced Reversal of PIF-Imposed Repression of Photomorphogenesis in Young *Arabidopsis* Seedlings. *Plant Cell* **21**: 3535-3553
- Li H, Johnson P, Stepanova A, Alonso JM, Ecker JR** (2004) Convergence of signaling pathways in the control of differential cell growth in *Arabidopsis*. *Dev Cell* **7**: 193-204
- Liscum E, Hangarter RP** (1993) Light-Stimulated Apical Hook Opening in Wild-Type *Arabidopsis thaliana* Seedlings. *Plant Physiol* **101**: 567-572
- Lorrain S, Allen T, Duek P, Whitelam GC, Fankhauser C** (2008) Phytochrome-mediated inhibition of shade avoidance involves degradation of growth-promoting bHLH transcription factors. *Plant J* **53**: 312-323
- Moubayidin L, Perilli S, Dello Ioio R, Di Mambro R, Costantino P, Sabatini S** (2010) The rate of cell differentiation controls the *Arabidopsis* root meristem growth phase. *Curr Biol* **20**: 1138-1143
- Nagpal P, Walker LM, Young JC, Sonawala A, Timpte C, Estelle M, Reed JW** (2000) AXR2 encodes a member of the Aux/IAA protein family. *Plant Physiol* **123**: 563-574
- Ogawa M, Hanada A, Yamauchi Y, Kuwahara A, Kamiya Y, Yamaguchi S** (2003) Gibberellin biosynthesis and response during *Arabidopsis* seed germination. *Plant Cell* **15**: 1591-1604
- Paz-Ares J, The Regia C** (2002) REGIA, an EU project on functional genomics of transcription factors from *Arabidopsis Thaliana*. *Comp Funct Genomics* **3**: 102-108
- Penfield S, Gilday AD, Halliday KJ, Graham IA** (2006) DELLA-mediated cotyledon expansion breaks coat-imposed seed dormancy. *Curr Biol* **16**: 2366-2370
- Peng J, Carol P, Richards DE, King KE, Cowling RJ, Murphy GP, Harberd NP** (1997) The *Arabidopsis* GAI gene defines a signaling pathway that negatively regulates gibberellin responses. *Genes Dev* **11**: 3194-3205
- Raz V, Ecker JR** (1999) Regulation of differential growth in the apical hook of *Arabidopsis*. *Development* **126**: 3661-3668
- Reid JB, Botwright NA, Smith JJ, O'Neill DP, Kerckhoffs LH** (2002) Control of gibberellin levels and gene expression during de-etiolation in pea. *Plant Physiol* **128**: 734-741

- Roman G, Lubarsky B, Kieber JJ, Rothenberg M, Ecker JR** (1995) Genetic analysis of ethylene signal transduction in *Arabidopsis thaliana*: five novel mutant loci integrated into a stress response pathway. *Genetics* **139**: 1393-1409
- Savaldi-Goldstein S, Peto C, Chory J** (2007) The epidermis both drives and restricts plant shoot growth. *Nature* **446**: 199-202
- Scacchi E, Osmont KS, Beuchat J, Salinas P, Navarrete-Gomez M, Trigueros M, Ferrandiz C, Hardtke CS** (2009) Dynamic, auxin-responsive plasma membrane-to-nucleus movement of *Arabidopsis* BRX. *Development* **136**: 2059-2067
- Shin J, Kim K, Kang H, Zulfugarov IS, Bae G, Lee CH, Lee D, Choi G** (2009) Phytochromes promote seedling light responses by inhibiting four negatively-acting phytochrome-interacting factors. *Proc Natl Acad Sci U S A* **106**: 7660-7665
- Stepanova AN, Robertson-Hoyt J, Yun J, Benavente LM, Xie DY, Dolezal K, Schlereth A, Jurgens G, Alonso JM** (2008) TAA1-mediated auxin biosynthesis is essential for hormone crosstalk and plant development. *Cell* **133**: 177-191
- Stowe-Evans EL, Harper RM, Motchoulski AV, Liscum E** (1998) NPH4, a conditional modulator of auxin-dependent differential growth responses in *Arabidopsis*. *Plant Physiol* **118**: 1265-1275
- Tatematsu K, Kumagai S, Muto H, Sato A, Watahiki MK, Harper RM, Liscum E, Yamamoto KT** (2004) MASSUGU2 encodes Aux/IAA19, an auxin-regulated protein that functions together with the transcriptional activator NPH4/ARF7 to regulate differential growth responses of hypocotyl and formation of lateral roots in *Arabidopsis thaliana*. *Plant Cell* **16**: 379-393
- Thain SC, Vandenbussche F, Laarhoven LJ, Dowson-Day MJ, Wang ZY, Tobin EM, Harren FJ, Millar AJ, Van Der Straeten D** (2004) Circadian rhythms of ethylene emission in *Arabidopsis*. *Plant Physiol* **136**: 3751-3761
- Tsuchisaka A, Theologis A** (2004) Unique and overlapping expression patterns among the *Arabidopsis* 1-amino-cyclopropane-1-carboxylate synthase gene family members. *Plant Physiol* **136**: 2982-3000
- Tsuchisaka A, Yu G, Jin H, Alonso JM, Ecker JR, Zhang X, Gao S, Theologis A** (2009) A combinatorial interplay among the 1-aminocyclopropane-1-carboxylate isoforms regulates ethylene biosynthesis in *Arabidopsis thaliana*. *Genetics* **183**: 979-1003
- Ubeda-Tomas S, Federici F, Casimiro I, Beemster GT, Bhalerao R, Swarup R, Doerner P, Haseloff J, Bennett MJ** (2009) Gibberellin signaling in the endodermis controls *Arabidopsis* root meristem size. *Curr Biol* **19**: 1194-1199
- Úbeda-Tomás S, Swarup R, Coates J, Swarup K, Laplaze L, Beemster GT, Hedden P, Bhalerao R, Bennett MJ** (2008) Root growth in *Arabidopsis* requires gibberellin/DELLA signalling in the endodermis. *Nat Cell Biol* **10**: 625-628
- Vandenbussche F, Petrasek J, Zadnikova P, Hoyerova K, Pesek B, Raz V, Swarup R, Bennett M, Zazimalova E, Benkova E, Van Der Straeten D** (2010) The auxin influx carriers AUX1 and LAX3 are involved in auxin-ethylene interactions during apical hook development in *Arabidopsis thaliana* seedlings. *Development* **137**: 597-606
- Vogel JP, Woeste KE, Theologis A, Kieber JJ** (1998) Recessive and dominant mutations in the ethylene biosynthetic gene ACS5 of *Arabidopsis* confer cytokinin insensitivity and ethylene overproduction, respectively. *Proc Natl Acad Sci U S A* **95**: 4766-4771
- Vriezen WH, Achard P, Harberd NP, Van Der Straeten D** (2004) Ethylene-mediated enhancement of apical hook formation in etiolated *Arabidopsis thaliana* seedlings is gibberellin dependent. *Plant J* **37**: 505-516
- Wei N, Kwok SF, von Arnim AG, Lee A, McNellis TW, Piekos B, Deng XW** (1994) *Arabidopsis* COP8, COP10, and COP11 genes are involved in repression of photomorphogenic development in darkness. *Plant Cell* **6**: 629-643
- Wu G, Cameron JN, Ljung K, Spalding EP** (2010) A role for ABCB19-mediated polar auxin transport in seedling photomorphogenesis mediated by cryptochrome 1 and phytochrome B. *Plant J* **62**: 179-191

- Yamagami T, Tsuchisaka A, Yamada K, Haddon WF, Harden LA, Theologis A** (2003) Biochemical diversity among the 1-amino-cyclopropane-1-carboxylate synthase isozymes encoded by the Arabidopsis gene family. *J Biol Chem* **278**: 49102-49112
- Yamaguchi S, Kamiya Y, Sun T** (2001) Distinct cell-specific expression patterns of early and late gibberellin biosynthetic genes during Arabidopsis seed germination. *Plant J* **28**: 443-453
- Yang X, Lee S, So JH, Dharmasiri S, Dharmasiri N, Ge L, Jensen C, Hangarter R, Hobbie L, Estelle M** (2004) The IAA1 protein is encoded by AXR5 and is a substrate of SCF(TIR1). *Plant J* **40**: 772-782
- Zadnikova P, Petrasek J, Marhavy P, Raz V, Vandenbussche F, Ding Z, Schwarzerova K, Morita MT, Tasaka M, Hejatko J, Van Der Straeten D, Friml J, Benkova E** (2010) Role of PIN-mediated auxin efflux in apical hook development of Arabidopsis thaliana. *Development* **137**: 607-617
- Zentella R, Zhang ZL, Park M, Thomas SG, Endo A, Murase K, Fleet CM, Jikumaru Y, Nambara E, Kamiya Y, Sun TP** (2007) Global analysis of DELLA direct targets in early gibberellin signaling in Arabidopsis. *Plant Cell* **19**: 3037-3057
- Zhao X, Yu X, Foo E, Symons GM, Lopez J, Bendehakalu KT, Xiang J, Weller JL, Liu X, Reid JB, Lin C** (2007) A study of gibberellin homeostasis and cryptochrome-mediated blue light inhibition of hypocotyl elongation. *Plant Physiol* **145**: 106-118
- Zhao Y, Christensen SK, Fankhauser C, Cashman JR, Cohen JD, Weigel D, Chory J** (2001) A role for flavin monooxygenase-like enzymes in auxin biosynthesis. *Science* **291**: 306-309
- Zhong S, Zhao M, Shi T, Shi H, An F, Zhao Q, Guo H** (2009) EIN3/EIL1 cooperate with PIF1 to prevent photo-oxidation and to promote greening of Arabidopsis seedlings. *Proc Natl Acad Sci U S A* **106**: 21431-21436


**Reference list for Figure 4.10:**

1. **Li, H., Johnson, P., Stepanova, A., Alonso, J.M. and Ecker, J.R.** (2004) Convergence of signaling pathways in the control of differential cell growth in Arabidopsis. *Dev Cell*, **7**, 193-204.
2. **Knee, E.M., Hangarter, R.P. and Knee, M.** (2000) Interactions of light and ethylene in hypocotyl hook maintenance in Arabidopsis thaliana seedlings. *Physiol Plant*, **108**, 208-215.
3. **Achard, P., Liao, L., Jiang, C., Desnos, T., Bartlett, J., Fu, X. and Harberd, N.P.** (2007) DELLAs Contribute to Plant Photomorphogenesis. *Plant Physiol*, **143**, 1163-1172.
4. **Alabadí, D., Gallego-Bartolomé, J., García-Cárcel, L., Orlando, L., Rubio, V., Martínez, C., Frigerio, M., Iglesias-Pedraz, J.M., Espinosa, A., Deng, X.W. and Blázquez, M.A.** (2008) Gibberellins modulate light signaling pathways to prevent Arabidopsis seedling de-etiolation in darkness. *Plant J*, **53**, 324-335.
5. **Zhao, X., Yu, X., Foo, E., Symons, G.M., Lopez, J., Bendehakalu, K.T., Xiang, J., Weller, J.L., Liu, X., Reid, J.B. and Lin, C.** (2007) A study of gibberellin homeostasis and cryptochrome-mediated blue light inhibition of hypocotyl elongation. *Plant Physiol*, **145**, 106-118.
6. This work.
7. **Lehman, A., Black, R. and Ecker, J.R.** (1996) HOOKLESS1, an ethylene response gene, is required for differential cell elongation in the Arabidopsis hypocotyl. *Cell*, **85**, 183-194.
8. **Vandenbussche, F., Petrasek, J., Žádníková, P., Hoyerova, K., Pesek, B., Raz, V., Swarup, R., Bennett, M., Zazimalova, E., Benkova, E. and Van Der**

**Straeten, D.** (2010) The auxin influx carriers AUX1 and LAX3 are involved in auxin-ethylene interactions during apical hook development in *Arabidopsis thaliana* seedlings. *Development*, **137**, 597-606.


9. **Žádníková, P., Petrasek, J., Marhavy, P., Raz, V., Vandenbussche, F., Ding, Z., Schwarzerova, K., Morita, M.T., Tasaka, M., Hejatko, J., Van Der Straeten, D., Friml, J. and Benkova, E.** (2010) Role of PIN-mediated auxin efflux in apical hook development of *Arabidopsis thaliana*. *Development*, **137**, 607-617.

#### 4.7. Supporting figures




**Figure 4.S1. GAI and PIF5 interact in plant cells.**

BiFC analysis in tobacco leaves between GAI and PIF5 fusions to N- and C-terminal fragments of YFP, respectively. Left, visible; right, YFP fluorescence.



**Figure 4.S2 GAs revert the PAC-effect on *ProACS5:GUS*, *ProACS8:GUS*, and *ProDR5:GUS*.**

Expression patterns of *ProDR5:GUS*, *ProACS5:GUS*, and *ProACS8:GUS* in seedlings grown on control medium, or on medium supplemented with 0.2  $\mu$ M PAC or with 0.2  $\mu$ M PAC plus 50  $\mu$ M GA<sub>3</sub> for 36 h after germination.



**Figure 4.S3** Specific expression of GFP-gai-1 in the epidermis of etiolated seedlings.

Confocal image of a longitudinal section of the apical hook of a 1-day-old *ProML1:GFP-gai-1-11* seedling.



## **General discussion**





One of the major challenges during the transition from simple unicellular to complex multicellular organisms was the coordination between different cells in order to provide the appropriate response to internal and external cues. This issue is particularly important in plants, where continuous changes in the environment lead to phenotypical changes caused by plastic interpretation of developmental programs. Phytohormones play an essential role in this adaptation, both as instructive agents and as modifiers of existing programs (Alabadí et al., 2009, *Int. J. Dev. Biol.*). In this thesis we have attempted to shed some light on the mechanisms through which gibberellins (GAs) contribute to plant plastic behaviour. As stated in the introduction, we have addressed two general questions: (1) Where does the specificity of DELLA function reside in *Arabidopsis*? And (2) How do GAs modulate differential growth processes in *Arabidopsis*? In the following paragraphs, we will discuss the implications of the results presented in the previous chapters, trying to identify common threads in the mechanisms that we have elucidated, and in the context of plasticity in plant development.

### **D.1. Contribution of gibberellins to plant plasticity relies partially on DELLA protein subfuncionalization.**

As mentioned in the introduction, the *Arabidopsis* genome contains five genes encoding DELLA proteins, while many other plants only possess a single ortholog. This observation could be enough to deduce that the impact of DELLA multiplication on plant plasticity is only marginal, given that there does not seem to be an evident relationship between the number of *DELLA* genes and the degree of plasticity between different species. However, a different view is that the multiplication of *DELLA* genes in *Arabidopsis* (and other plants) may well be an alternative mechanism in these particular plants to achieve the plasticity that plants with single copies of *DELLA* genes generate by other means. Therefore, the investigation of a link between DELLA multiplication in *Arabidopsis* and plasticity may provide results that could perhaps be generalized to other similar situations.

If the previous idea is right, and *DELLA* multiplication confers advantages regarding plasticity in this species, it is reasonable to think that after diversification of expression, specialization would be the mean to improve the downstream signaling

events. Hence, each DELLA protein would be specialized in controlling processes related to their own expression domain. This hypothesis seems to hold true according to a search for genes coexpressed with each of the *DELLA* genes using the ATTED webtool (<http://atted.jp/>) at the TAIR website (<http://www.arabidopsis.org/>). Interestingly, the functional categories over-represented among the 300 best-scored coexpressed genes were different depending on the particular *DELLA* gene used in the analysis (Table D1), and these categories correlate with the known roles of each of the DELLA proteins. For instance, the main role of RGA and GAI as repressors of cell elongation (Dill and Sun, 2001) is suggested by the enrichment of “uni- and multi-dimensional growth” among the functional categories overrepresented in their set of coexpressed genes (Table D1). And the role of RGL1 and RGL2 together with RGA in flower development (Cheng et al., 2004) is also confirmed in this approach by the appearance of related functional categories overrepresented in their set of coexpressed genes (Table D1).

More importantly, this analysis may also reveal previously unknown roles of each DELLA protein, which of course should be confirmed or discarded on the basis of additional experimental evidence. Such would be the case for lateral root development, auxin signaling and transport, stomata movement or gravitropism (Table D1). Interestingly, our results (Chapters 3 and 4) confirm the involvement of DELLAs in two differential growth processes –hook and gravitropism- through their action on auxin signaling and transport, demonstrating again the potential use of this analysis, based on coexpression networks.

## **D.2. Regulation of transcriptional networks as a mechanism to improve plastic development**

A plausible mechanism through which a single hormone, GA, can trigger so many different developmental responses resides in a complex GA-GID-DELLA signaling pathway where the DELLA proteins, depending on the spatial or temporal context, will interact with different transcription factors (TFs) to trigger different subsets of GA responses. As shown in Chapter 2, one of the over-represented GO categories within the “molecular function” group in the *HS::gai-1* microarray was “transcription factor activity”.

**Table D1: Functional categories over-represented in the set of DELLA coregulated genes.**

<b>RGA</b>			<b>GAI</b>		
<b>GO term</b>	<b>list1_positive_ids</b>	<b>adj_pvalue</b>	<b>GO term</b>	<b>list1_positive_ids</b>	<b>adj_pvalue</b>
6833	water transport	2,21E-02	9639	response to red or far red light	3,85E-02
32880	regulation of protein localization	1,17E-03	9734	auxin mediated signaling pathway	7,65E-05
10014	meristem initiation	2,77E-02	51258	protein polymerization	3,62E-02
48443	stamen development	1,06E-02	7047	cell wall organization	4,33E-02
9965	leaf morphogenesis	7,64E-03	9825	multidimensional cell growth	4,43E-03
10015	root morphogenesis	1,84E-02	9826	unidimensional cell growth	3,20E-02
48527	lateral root development	1,78E-02	9926	auxin polar transport	3,22E-02
10540	basipetal auxin transport	2,21E-02	10268	brassinosteroids homeostasis	4,33E-02
	xylem and phloem pattern				
10051	formation	1,19E-03	48589	developmental growth	3,11E-02
9845	seed germination	3,02E-02	9911	positive regulation of flower	1,01E-02
9799	determination of symmetry	1,45E-02	48513	organ development	3,22E-02
				xylem and phloem pattern	
10152	pollen maturation	4,18E-02	10051	formation	2,36E-02
10075	regulation of meristem growth	1,27E-02	15994	chlorophyll biosynthetic process	3,95E-04
9825	multidimensional cell growth	2,30E-04			
9826	unidimensional cell growth	3,30E-03			
9913	epidermal cell differentiation	2,32E-03			
42538	hyperosmotic salinity response	1,20E-02			
9958	positive gravitropism	4,92E-03			
9637	response to blue light	1,95E-02			
10218	response to far red light	1,20E-02			
10161	red light signaling pathway	2,21E-02			
10118	regulation of stomata movement	1,35E-02			
	transmembrane receptor protein				
7169	tyrosine kinase signaling	3,60E-06			
	auxin mediated signaling				
9734	pathway	8,07E-12			
	brassinosteroids mediated				
9742	signaling	2,17E-04			
9664	plant-type cell wall organization	1,47E-02			
7010	cytoskeleton organization	4,55E-03			
51301	cell division	2,14E-02			
9832	plant-type cell wall biogenesis	3,30E-03			
7049	cell cycle	4,18E-02			
7017	microtubule-based process	1,86E-02			
6644	phospholipid metabolic process	2,26E-02			
30244	cellulose biosynthetic process	8,24E-03			

<b>RGL1</b>		
<b>GO term</b>	<b>list1_positive_ids</b>	<b>adj_pvalue</b>
8610	lipid catabolic process	6,14E-07
10166	wax metabolic process	2,61E-08
6633	fatty-acid metabolic process	7,17E-05
6011	UDP-glucose metabolic process	8,03E-03
9908	flower development	3,49E-02
9753	response to jasmonic acid stimulus	4,58E-02
10166	response to hormone stimulus	2,61E-08


  

<b>RGL2</b>		
<b>GO term</b>	<b>list1_positive_ids</b>	<b>adj_pvalue</b>
16042	lipid catabolic process	1,70E-03
10166	wax metabolic process	7,54E-04
6633	fatty acid biosynthetic process	3,57E-06
30154	cell differentiation	1,88E-05
9887	organ morphogenesis	9,43E-03
48440	carpel development	3,00E-09

<b>RGL3</b>			<b>GAI</b>		
<b>GO term</b>	<b>list1_positive_ids</b>	<b>adj_pvalue</b>	<b>GO term</b>	<b>list1_positive_ids</b>	<b>adj_pvalue</b>
9611	response to wounding	1,89E-10	9739	response to gibberellin stimulus	3,95E-02
6952	defense response	6,50E-03	9738	abscisic acid mediated signaling	1,71E-03
				regulation of ethylene mediated	
9651	response to salt stress	3,04E-05	10104	signaling pathway	9,43E-03
				jasmonic acid mediated signaling	
9414	response to water deprivation	1,95E-03	9867	pathway	9,43E-03
				negative regulation of signal	
10200	response to chitin	2,36E-04	9968	transduction	2,24E-03

If this is true, searching for DELLA-regulated TFs we might figure out by which means DELLAs trigger different downstream events. Looking at *HS::gai-1* microarray data (Chapter 2) it seems that *gai-1*, under the control of a heat-shock inducible promoter, preferentially modulates the expression of only a few transcription factor families: MYB, bHLH, GRAS, C2C2 or Homeobox (Fig D.1).




**Figure D.1. Transcription factor families regulated by *gai-1*.**

All *gai-1* targets annotate as TF in the *HS::gai-1* microarray dataset were grouped based on their TF families. Each family is shown as a percentage of the total families present in the dataset.

Thus, the set of transcription factors coexpressed with DELLA could reveal new connections between DELLAs and transcriptional regulation of GA targets. If the relative enrichment of a particular TF family is meaningful, the results shown in Fig D.2 suggest that bHLH proteins are at the core of DELLA regulated responses (only RGL3, probably specialized in defense, does not show coregulation with this TF family). But beyond the common points between all DELLAs, the different DELLA proteins seem to mobilize different TF families to exert their function. For instance, RGA and GAI, with a prominent role in general growth, are coexpressed with homeobox TF families, while RGL1 and RGL2, specialized partially in flower development, show better coexpression with MADS-box, MYB and Squamosa Binding Protein (SBP) TF families (also involved in this process).

In summary, it seems that after the diversification of DELLAs expression pattern, coexpression of specific DELLAs and certain TFs might eventually explain the different roles of each of the DELLA proteins in Arabidopsis. On the other hand, it could just indicate that different developmental processes might be triggered preferentially by different TF families (i.e. homeobox for vegetative growth vs MADS-box for flower development), and each DELLA protein would be mobilized by different developmental programs to cooperate with these TF families. In that case, coexpressed TF could be not

only DELLA targets but also modulators of DELLA expression, or even direct interactors that would mediate DELLA regulation of transcription.




**Figure D.2. Coexpressed transcription factor families**

Using the ATTEDII webtool, it was found the 100 best scored coregulated transcription factor for each *DELLA* gene. Only transcription factor families representing >5% of the total list in at least one coregulated set are shown. Transcription factor families representing < 5% are englobed as others.

### D.3. DELLA co-expressed as a source of new interactors.

So far, no DNA-binding domain has been found in DELLA proteins and it is currently accepted that they regulate gene transcription through the interaction with DNA binding transcription factors, such as PIF proteins (de Lucas et al., 2008; Feng et al., 2008), or modulators of the activity of transcription factor, such as JAZ proteins (Hou et al., 2010). Thus, an even more exciting possibility is that some of the DELLA co-expressed TF, apart from being regulated transcriptionally by DELLA, they could be regulated by physical interaction. In other words, DELLAs would had had bigger

chances to interact with the coexpressed TFs. Comparison of a list of all DELLA coexpressed TF with a list of all Arabidopsis TF shows that some TF families are over-represented within the DELLA coexpressed set such as bHLH, homeobox, SBP, TCP and Trihelix families (Fig D3). Interaction with bHLH TFs has been already shown (de Lucas et al., 2008; Feng et al., 2008; Arnaud et al., 2010; Josse et al., 2011). Although physical interaction with members of the other over-represented TF families has not been shown, there is a large overlap between GA-related responses and the functions of these DELLA-coexpressed TFs, (Hay et al., 2004; Zhang et al., 2007; Tatematsu et al., 2008; Breuer et al., 2009). For instance Squamosa-Binding Proteins (SBP) are related to several processes controlled by GAs such as juvenile to adult phase transition, flowering and male sterility (Chen et al., 2010). As an example, SPL8 over-expression induces GA constitutive response that leads to anther indehiscence and hence infertility (Zhang et al., 2007).



**Figure D.3. Comparison of representation of *DELLA* coexpressed TF with the overall TF list from Arabidopsis.**

All DELLA coexpressed TF were combined. Each TF family is represented as percentage of the total of families in this dataset. A comparison with the predicted list of all Arabidopsis TF (Qu and Zhu, 2006) was done to find over-represented families among the DELLA coexpressed TF.

Lastly, an additional source of information to find putative TFs involved in transcriptional regulation by DELLA proteins is the enrichment of certain *cis* elements in the promoters of genes differentially expressed in *HS::gai-1* plants (Chapter 2). For example, we found known binding sites for DOF (C2C2 family) (AAAG) or ARR1 (GARP family) (NGATT) TFs. These two families seem not to be over-represented in the previous analysis of co-expressed TF families, but this could reflect that the over-represented elements present in the promoter analysis of Chapter 2 belong to a certain context (2 day old etiolated seedlings) where GAI interacts with a certain pool of factors while the DELLA co-expressed genes are a compilation of the co-expressed genes through the whole *Arabidopsis* life. This points out to the importance of the context where a DELLA is expressed to adopt the control over a certain process and how the same proteins can control different processes through the interaction with different proteins.

#### **D.4. Gibberellin modulation of differential growth processes**

During this thesis we have studied the contribution of GAs to two differential growth processes: gravitropism and apical hook development. A comparative look at the mechanisms found in both processes reveals an interesting parallelism: GAs act as modulators of differential growth through their regulation upon auxin signaling and/or transport. This type of interaction between GAs and other signaling pathways is relatively novel, given that GAs have been previously shown to act on cell growth through direct regulation of genes encoding enzymes involved in cell wall modifications, such as expansins or xyloglucan endotransglucosylases (Lee and Kende, 2001; Vogler et al., 2003; Jan et al., 2004) often as a downstream subsidiary signal for the action of other hormones such as auxin or ethylene (Achard et al., 2003; Fu and Harberd, 2003; Frigerio et al., 2006). Conceptually, what we have found is novel in that during differential growth, GAs primarily modulate the activity of auxin, the master hormone in these processes, constituting a fine-tuning mechanism, rather than a *bona fide* instructive signal. Support for this claim not only comes from the molecular links found between DELLAs and auxin-signaling elements and their targets, but this mechanistic interaction may probably be extended to other processes beyond differential growth, based on the functional analysis of *DELLA*-coregulated genes discussed in the previous paragraphs (Table D1).

The parallelisms between the two regulatory mechanisms described in Chapters 3 and 4 are more than a simple conceptualization of the interaction between GAs and auxin during differential growth. Although at first look it seems like GAs regulate completely different steps of auxin activity in each process (gravitropism *vs* hook formation), a more careful analysis of the evidence suggests that GAs regulate both processes through the same mechanism. For instance, the repression of *IAA19/MSG2* expression by DELLAs has been proven essential for the modulation of hypocotyl gravitropism (Chapter 3), but this interaction may indeed also contribute to the formation of the apical hook, based on the hook-defective phenotype of *msg2-1* mutants (Tatematsu et al., 2004). Similarly, the repression of *PIN3* and *PIN7* by DELLAs is instrumental for the regulation of hook formation by GAs (Chapter 4), but one cannot forget that these two genes encode key auxin efflux carriers involved in the redistribution of auxins during the hypocotyl gravitropic response (Rakusova, Gallego-Bartolomé et al., 2011). And, finally, *HLS1* repression by DELLAs is important for hook unfolding (Chapter 4), but it may also be relevant for the regulation of gravitropism according to the reported phenotype of *hls1* mutants on gravitropic reorientation (Hamaguchi et al., 2008). Thus, it seems reasonable that the regulation by DELLAs of all these genes (*IAA19/MSG2*, *HLS1*, *PIN3* and *PIN7*) may represent a general core mechanism for the regulation of differential growth by GAs. Further studies should be done to confirm whether these interactions, although plausible, are actually happening in these contexts.

A part from the effect of GAs on auxin activity, we have also reported the positive effect of GAs on ethylene synthesis to prevent hook opening, through activation of ACC synthase genes. Interestingly, part of the ethylene effect on hook development is mediated by auxin and through regulation of the auxin efflux (Zadnikova et al., 2010) showing a parallelism with GAs action on auxin activity. The fact that both hormones, GAs and ethylene, show cross-regulation between them (1) GAs induce ethylene synthesis, (2) ethylene affects GAs response (Vriezen et al., 2004) and both together modulate auxins as common output to fine-tune the hook development, highlights the complexity of hormonal crosstalk and the importance of understanding the overall circuits and interactions among them.



One of the reasons why hormonal crosstalk is currently a very active field of plant research is that it may represent a common theme in plant development to fine-tune certain plastic responses. Achieving this fine-tuning may be the purpose of recruiting GAs/DELLAs as a modulator of the network that regulates differential growth. However, the demonstration that these new interactions constitute a selective advantage in natural environments still represents a big challenge.


### **D.5. Future perspectives**

Most of the previous work on GA signaling, including the work presented here, has been done analyzing GA responses in whole whole seedlings. However, plant organs are heterogeneous regarding the identity of different tissues and cell types. If this apparently obvious statement is taken into account, several questions arise that have hardly been addressed until now. For instance, how is growth coordinated through the different cell types that conform a given organ? Does GA signaling proceed through different circuits in the different tissues? When a particular whole-seedling response is analyzed, is GA signaling necessary only in a specific subset of cells, or does it happen homogeneously throughout the organ? All these questions represent only the initial motivation to analyze the spatial regulation of GA signaling.


A hint of the importance of spatial constraints in GA signaling has been beautifully exposed in recent work which showed that GA signaling in the root endodermis –and not in other tissues– is essential for coordinated growth of the root (Úbeda-Tomás et al., 2008). Importantly, this spatial component of GA signaling operates in other developmental contexts, as we have observed in our lab in the aerial part of vegetative tissues. For instance, results in Chapter 4 show that an active GA response in the endodermis is needed for the proper apical hook development, while a block of GA signaling in the epidermis does not affect this process. And in the course of this thesis we have generated unpublished results showing that, again, hypocotyl growth can be impaired by blocking GA signaling specifically in the endodermis, but not in the epidermis (Fig D.4A). Moreover, the attenuation of the gravitropic response conferred by GAs (Chapter 3) seems to happen, again, in the endodermis, given that forced expression of *gai-1D* in the endodermis alone, but not in the epidermis, was able to enhance gravitropic reorientation (Figure D.4B).

Of course, spatial constraints do not restrict to GA signaling, but have also been reported for other hormones. For instance, it has been shown that brassinosteroids control whole hypocotyl growth through the epidermis (Savaldi-Goldstein et al., 2007). Moreover, auxin signaling in root elongation epidermal cells is essential to develop a normal gravitropic response (Swarup et al., 2005).

A



B



**Figure D4. Cell-type specific effect of GA signaling upon gravitropic reorientation.**

A) Hypocotyl length of two day old Arabidopsis seedling expressing *gai-1D* from an endodermis-specific (*SCR*) or an epidermis-specific (*ML1*) promoter. *Ler*, *SCR::gai-1D:GR:YFP* and *ML1::gai-1D:GFP* seedlings were grown for 3 days in darkness in plates with either mock, 0.4  $\mu$ M PAC or 10  $\mu$ M dexamethasone solutions.

B) Gravitropic reorientation in seedlings expressing *gai-1D* from an endodermis-specific (*SCR*) or an epidermis-specific (*ML1*) promoter. *SCR::gai-1D:GR:YFP* and *ML1::gai-1D:GFP* seedlings were grown for 3 days in darkness on vertical MS plates and then transferred to plates with either mock or 10  $\mu$ M dexamethasone solutions for 15 h. Then the plates were turned 90° at time 0. The pictures illustrate *gai-1D:YFP* and *gai-1D:GFP* localization in the nuclei of endodermis and epidermis cells respectively.

As mentioned before, a previous study placed the epidermis as the main tissue controlling hypocotyl growth (Savaldi-Goldstein et al., 2007). Although GAs are also

mandatory for growth, we have observed that GA signaling in the epidermis does not produce any effect on hypocotyl growth while it is necessary in endodermal cells. Interestingly, the periclinal chimera of the grapevine Pinot Meunier points to the same direction. It expresses *gai-1* only in the epidermis and, however its size is indistinguishable from the nonchimeric variant (Boss and Thomas, 2002). However, plants regenerated from epidermal cells, but not from other tissues are dwarf. Notably, both Pinot Meunier mutant and *pML1::gai-1* Arabidopsis show defects on trichome development similar to GA deficient *gal* mutant.

The complexity that arises when both temporal and spatial regulation are combined is probably at the core of the generation of plastic developmental responses in plants, and points to an exciting direction for future investigations

## D.6. Bibliography

- Achard P, Vriezen WH, Van Der Straeten D, Harberd NP** (2003) Ethylene regulates arabidopsis development via the modulation of DELLA protein growth repressor function. *Plant Cell* **15**: 2816-2825
- Arnaud N, Girin T, Sorefan K, Fuentes S, Wood TA, Lawrenson T, Sablowski R, Ostergaard L** (2010) Gibberellins control fruit patterning in Arabidopsis thaliana. *Genes Dev* **24**: 2127-2132
- Boss PK, Thomas MR** (2002) Association of dwarfism and floral induction with a grape 'green revolution' mutation. *Nature* **416**: 847-850
- Breuer C, Kawamura A, Ichikawa T, Tominaga-Wada R, Wada T, Kondou Y, Muto S, Matsui M, Sugimoto K** (2009) The trihelix transcription factor GTL1 regulates ploidy-dependent cell growth in the Arabidopsis trichome. *Plant Cell* **21**: 2307-2322
- Chen X, Zhang Z, Liu D, Zhang K, Li A, Mao L** (2010) SQUAMOSA promoter-binding protein-like transcription factors: star players for plant growth and development. *J Integr Plant Biol* **52**: 946-951
- Cheng H, Qin L, Lee S, Fu X, Richards DE, Cao D, Luo D, Harberd NP, Peng J** (2004) Gibberellin regulates Arabidopsis floral development via suppression of DELLA protein function. *Development* **131**: 1055-1064
- de Lucas M, Davière JM, Rodríguez-Falcón M, Pontin M, Iglesias-Pedraz JM, Lorrain S, Fankhauser C, Blázquez MA, Titarenko E, Prat S** (2008) A molecular framework for light and gibberellin control of cell elongation. *Nature* **451**: 480-484
- Dill A, Sun T** (2001) Synergistic derepression of gibberellin signaling by removing RGA and GAI function in Arabidopsis thaliana. *Genetics* **159**: 777-785
- Feng S, Martinez C, Gusmaroli G, Wang Y, Zhou J, Wang F, Chen L, Yu L, Iglesias-Pedraz JM, Kircher S, Schafer E, Fu X, Fan LM, Deng XW** (2008) Coordinated regulation of Arabidopsis thaliana development by light and gibberellins. *Nature* **451**: 475-479
- Frigerio M, Alabadí D, Pérez-Gómez J, García-Cárcel L, Phillips AL, Hedden P, Blázquez MA** (2006) Transcriptional regulation of gibberellin metabolism genes by auxin signaling in Arabidopsis. *Plant Physiol* **142**: 553-563

- Fu X, Harberd NP** (2003) Auxin promotes Arabidopsis root growth by modulating gibberellin response. *Nature* **421**: 740-743
- Hamaguchi A, Yamashino T, Koizumi N, Kiba T, Kojima M, Sakakibara H, Mizuno T** (2008) A small subfamily of Arabidopsis RADIALIS-LIKE SANT/MYB genes: a link to HOOKLESS1-mediated signal transduction during early morphogenesis. *Biosci Biotechnol Biochem* **72**: 2687-2696
- Hay A, Craft J, Tsiantis M** (2004) Plant hormones and homeoboxes: bridging the gap? *Bioessays* **26**: 395-404
- Hou X, Lee LY, Xia K, Yan Y, Yu H** (2010) DELLAs modulate jasmonate signaling via competitive binding to JAZs. *Dev Cell* **19**: 884-894
- Jan A, Yang G, Nakamura H, Ichikawa H, Kitano H, Matsuoka M, Matsumoto H, Komatsu S** (2004) Characterization of a xyloglucan endotransglucosylase gene that is up-regulated by gibberellin in rice. *Plant Physiol* **136**: 3670-3681
- Josse EM, Gan Y, Bou-Torrent J, Stewart KL, Gilday AD, Jeffree CE, Vaistij FE, Martinez-Garcia JF, Nagy F, Graham IA, Halliday KJ** (2011) A DELLA in Disguise: SPATULA Restrains the Growth of the Developing Arabidopsis Seedling. *Plant Cell*
- Lee Y, Kende H** (2001) Expression of beta-expansins is correlated with internodal elongation in deepwater rice. *Plant Physiol* **127**: 645-654
- Rakusova H, Gallego-Bartolomé J, Vanstraelen M, Robert H, Alabadí D, Blázquez M, Benkova E, Friml J** (2011) Polarization of PIN3-dependent auxin transport for hypocotyl gravitropic response in Arabidopsis thaliana. *Plant J.*
- Savaldi-Goldstein S, Peto C, Chory J** (2007) The epidermis both drives and restricts plant shoot growth. *Nature* **446**: 199-202
- Swarup R, Kramer EM, Perry P, Knox K, Leyser HM, Haseloff J, Beemster GT, Bhalerao R, Bennett MJ** (2005) Root gravitropism requires lateral root cap and epidermal cells for transport and response to a mobile auxin signal. *Nat Cell Biol* **7**: 1057-1065
- Tatematsu K, Kumagai S, Muto H, Sato A, Watahiki MK, Harper RM, Liscum E, Yamamoto KT** (2004) MASSUGU2 encodes Aux/IAA19, an auxin-regulated protein that functions together with the transcriptional activator NPH4/ARF7 to regulate differential growth responses of hypocotyl and formation of lateral roots in Arabidopsis thaliana. *Plant Cell* **16**: 379-393
- Tatematsu K, Nakabayashi K, Kamiya Y, Nambara E** (2008) Transcription factor AtTCP14 regulates embryonic growth potential during seed germination in Arabidopsis thaliana. *Plant J* **53**: 42-52
- Úbeda-Tomás S, Swarup R, Coates J, Swarup K, Laplaze L, Beemster GT, Hedden P, Bhalerao R, Bennett MJ** (2008) Root growth in Arabidopsis requires gibberellin/DELTA signalling in the endodermis. *Nat Cell Biol* **10**: 625-628
- Vogler H, Caderas D, Mandel T, Kuhlemeier C** (2003) Domains of expansin gene expression define growth regions in the shoot apex of tomato. *Plant Mol Biol* **53**: 267-272
- Vriezen WH, Achard P, Harberd NP, Van Der Straeten D** (2004) Ethylene-mediated enhancement of apical hook formation in etiolated Arabidopsis thaliana seedlings is gibberellin dependent. *Plant J* **37**: 505-516
- Zadnikova P, Petrasek J, Marhavy P, Raz V, Vandenbussche F, Ding Z, Schwarzerova K, Morita MT, Tasaka M, Hejatko J, Van Der Straeten D, Friml J, Benkova E** (2010) Role of PIN-mediated auxin efflux in apical hook development of Arabidopsis thaliana. *Development* **137**: 607-617
- Zhang Y, Schwarz S, Saedler H, Huijser P** (2007) SPL8, a local regulator in a subset of gibberellin-mediated developmental processes in Arabidopsis. *Plant Mol Biol* **63**: 429-439

## **Conclusions**



- Diversification of DELLAs' roles in Arabidopsis relies mainly on their differential expression patterns, although certain degree of biochemical specialization due to the exposure to different protein contexts cannot be discarded.
- The identity of early targets of GAI in darkness confirms that DELLAs mobilize different transcriptomes depending on the developmental condition. HY5 and PIF transcription factors mediate the regulation of a large subset of DELLA targets. However, our results also indicate that DELLAs act through additional, yet unidentified, transcription factors.
- Gibberellins decrease auxin sensitivity through the induction of *IAA19/MSG2* expression in hypocotyls. As a result, gibberellins increase the variance in the gravitropic response of the hypocotyl in a population of seedlings, and they attenuate gravitropism in case of conflicting tropic stimuli, such as during shade avoidance.
- The mechanism by which gibberellins promote the formation of the apical hook is through upregulation of key genes for auxin homeostasis and transport – *HLS1*, *PIN3*, and *PIN7*–, most likely in the endodermis. Moreover gibberellins cooperate with ethylene during the maintenance and opening phases to prevent hook opening. This is achieved through the activation of ethylene biosynthetic genes, *ACS5* and *ACS8*, that leads to enhanced ethylene levels.